

On general Eulerian integral of certain products of Aleph-function, the multivariable I-function and a class of polynomials II

F.Y. AYANT¹

¹ Teacher in High School , France

ABSTRACT

The object of this paper is to establish an general Eulerian integral involving the product of the Aleph-function, the multivariable I-function defined by Nambisan et al [2], a general class of multivariable polynomials and a generalized hypergeometric function which provide unification and extension of numerous results. We will study the particular case concerning the multivariable H-function defined by Srivastava et al [8] and the Srivastava-Daoust polynomial [5].

Keywords: Eulerian integral, multivariable I-function, Lauricella function of several variables, multivariable Aleph-function, generalized hypergeometric function, multivariable H-function, Srivastava-Daoust polynomial

2010 Mathematics Subject Classification. 33C45, 33C60, 26D20

1. Introduction

In this paper, we evaluate a general Eulerian integral concerning the product the multivariable I-functions defined by Nambisan et al [2], the multivariable Aleph-function a generalized hypergeometric function and a class of multivariable polynomials. We will give a serie expansion of a multivariable Aleph-function. The multivariable I-function is an extension of the multivariable H-function defined by Srivastava et al [8]. We will given a contracted form. The Aleph-function of several variables is a generalisation of G and H-functions of several variables defined by Srivastava et al [8]. The multiple Mellin-Barnes integral occuring in this paper will be referred to as the multivariable Aleph-function throughout our present study and will be defined and represented as follows.

$$\text{We define : } \aleph(z_1''', \dots, z_v''') = \aleph_{p_i, q_i, \tau_i; R; p_i(1), q_i(1), \tau_i(1); R^{(1)}; \dots; p_i(r), q_i(r), \tau_i(r); R^{(v)}}^{0, n; m_1, n_1, \dots, m_v, n_v} \left(\begin{matrix} z_1''' \\ \vdots \\ z_v''' \end{matrix} \right)$$

$$[(a_j; \alpha_j^{(1)}, \dots, \alpha_j^{(v)})_{1, n}] \quad , [\tau_i(a_{ji}; \alpha_{ji}^{(1)}, \dots, \alpha_{ji}^{(v)})_{n+1, p_i}] :$$

$$\dots \dots \dots \quad , [\tau_i(b_{ji}; \beta_{ji}^{(1)}, \dots, \beta_{ji}^{(v)})_{m+1, q_i}] :$$

$$\left[(c_j^{(1)}; \gamma_j^{(1)})_{1, n_1} \right], [\tau_{i(1)}(c_{ji(1)}; \gamma_{ji(1)})_{n_1+1, p_i(1)}]; \dots ; [(c_j^{(v)}; \gamma_j^{(v)})_{1, n_v}], [\tau_{i(v)}(c_{ji(v)}; \gamma_{ji(v)})_{n_v+1, p_i(v)}]$$

$$\left[(d_j^{(1)}; \delta_j^{(1)})_{1, m_1} \right], [\tau_{i(1)}(d_{ji(1)}; \delta_{ji(1)})_{m_1+1, q_i(1)}]; \dots ; [(d_j^{(v)}; \delta_j^{(v)})_{1, m_v}], [\tau_{i(v)}(d_{ji(v)}; \delta_{ji(v)})_{m_v+1, q_i(v)}]$$

$$= \frac{1}{(2\pi\omega)^v} \int_{L_1} \dots \int_{L_v} \psi_1(s_1, \dots, s_v) \prod_{k=1}^v \xi_k(s_k) z_k''' s_k \, ds_1 \dots ds_r \tag{1.1}$$

with $\omega = \sqrt{-1}$

$$\psi_1(s_1, \dots, s_v) = \frac{\prod_{j=1}^n \Gamma(1 - a_j + \sum_{k=1}^v \alpha_j^{(k)} s_k)}{\sum_{i=1}^R [\tau_i \prod_{j=n+1}^{p_i} \Gamma(a_{ji} - \sum_{k=1}^v \alpha_{ji}^{(k)} s_k) \prod_{j=1}^{q_i} \Gamma(1 - b_{ji} + \sum_{k=1}^v \beta_{ji}^{(k)} s_k)]} \tag{1.2}$$

$$\text{and } \xi_k(s_k) = \frac{\prod_{j=1}^{m_k} \Gamma(d_j^{(k)} - \delta_j^{(k)} s_k) \prod_{j=1}^{n_k} \Gamma(1 - c_j^{(k)} + \gamma_j^{(k)} s_k)}{\sum_{i^{(k)}=1}^{R^{(k)}} [\tau_{i^{(k)}} \prod_{j=m_k+1}^{q_i^{(k)}} \Gamma(1 - d_{j i^{(k)}}^{(k)} + \delta_{j i^{(k)}}^{(k)} s_k) \prod_{j=n_k+1}^{p_i^{(k)}} \Gamma(c_{j i^{(k)}}^{(k)} - \gamma_{j i^{(k)}}^{(k)} s_k)]} \quad (1.3)$$

Suppose , as usual , that the parameters

$$a_j, j = 1, \dots, p; b_j, j = 1, \dots, q;$$

$$c_j^{(k)}, j = 1, \dots, n_k; c_{j i^{(k)}}^{(k)}, j = n_k + 1, \dots, p_i^{(k)};$$

$$d_j^{(k)}, j = 1, \dots, m_k; d_{j i^{(k)}}^{(k)}, j = m_k + 1, \dots, q_i^{(k)};$$

$$\text{with } k = 1 \dots, r, i = 1, \dots, R, i^{(k)} = 1, \dots, R^{(k)}$$

are complex numbers , and the $\alpha's, \beta's, \gamma's$ and $\delta's$ are assumed to be positive real numbers for standardization purpose such that

$$U_i^{(k)} = \sum_{j=1}^n \alpha_j^{(k)} + \tau_i \sum_{j=n+1}^{p_i} \alpha_{j i}^{(k)} + \sum_{j=1}^{n_k} \gamma_j^{(k)} + \tau_{i^{(k)}} \sum_{j=n_k+1}^{p_i^{(k)}} \gamma_{j i^{(k)}}^{(k)} - \tau_i \sum_{j=1}^{q_i} \beta_{j i}^{(k)} - \sum_{j=1}^{m_k} \delta_j^{(k)} - \tau_{i^{(k)}} \sum_{j=m_k+1}^{q_i^{(k)}} \delta_{j i^{(k)}}^{(k)} \leq 0 \quad (1.4)$$

The reals numbers τ_i are positives for $i = 1$ to R , $\tau_{i^{(k)}}$ are positives for $i^{(k)} = 1$ to $R^{(k)}$

The contour L_k is in the s_k -p lane and run from $\sigma - i\infty$ to $\sigma + i\infty$ where σ is a real number with loop , if necessary ,ensure that the poles of $\Gamma(d_j^{(k)} - \delta_j^{(k)} s_k)$ with $j = 1$ to m_k are separated from those of $\Gamma(1 - a_j + \sum_{i=1}^r \alpha_j^{(k)} s_k)$ with $j = 1$ to n and $\Gamma(1 - c_j^{(k)} + \gamma_j^{(k)} s_k)$ with $j = 1$ to n_k to the left of the

contour L_k . The condition for absolute convergence of multiple Mellin-Barnes type contour (1.9) can be obtained by extension of the corresponding conditions for multivariable H-function given by as :

$$|arg z_k'''| < \frac{1}{2} A_i^{(k)} \pi, \text{ where}$$

$$A_i^{(k)} = \sum_{j=1}^n \alpha_j^{(k)} - \tau_i \sum_{j=n+1}^{p_i} \alpha_{j i}^{(k)} - \tau_i \sum_{j=1}^{q_i} \beta_{j i}^{(k)} + \sum_{j=1}^{n_k} \gamma_j^{(k)} - \tau_{i^{(k)}} \sum_{j=n_k+1}^{p_i^{(k)}} \gamma_{j i^{(k)}}^{(k)} + \sum_{j=1}^{m_k} \delta_j^{(k)} - \tau_{i^{(k)}} \sum_{j=m_k+1}^{q_i^{(k)}} \delta_{j i^{(k)}}^{(k)} > 0, \text{ with } k = 1, \dots, v, i = 1, \dots, R, i^{(k)} = 1, \dots, R^{(k)} \quad (1.5)$$

The complex numbers z_i are not zero. Throughout this document , we assume the existence and absolute convergence conditions of the multivariable Aleph-function.

We may establish the the asymptotic expansion in the following convenient form :

$$\aleph(z_1''', \dots, z_v''') = O(|z_1'''|^{\alpha_1}, \dots, |z_r'''|^{\alpha_r}), \max(|z_1'''|, \dots, |z_v'''|) \rightarrow 0$$

$$\aleph(z_1''', \dots, z_v''') = O(|z_1'''|^{\beta_1}, \dots, |z_r'''|^{\beta_r}), \min(|z_1'''|, \dots, |z_v'''|) \rightarrow \infty$$

where $k = 1, \dots, r : \alpha_k = \min[Re(d_j^{(k)} / \delta_j^{(k)})], j = 1, \dots, m_k$ and

$$\beta_k = \max[\operatorname{Re}((c_j^{(k)} - 1)/\gamma_j^{(k)})], j = 1, \dots, n_k$$

Series representation of Aleph-function of u -variables is given by

$$\begin{aligned} \aleph(z_1''', \dots, z_v''') &= \sum_{G_1, \dots, G_v=0}^{\infty} \sum_{g_1=0}^{m_1} \dots \sum_{g_v=0}^{m_v} \frac{(-)^{G_1+\dots+G_v}}{\delta_{g_1}^{G_1!} \dots \delta_{g_v}^{G_v!}} \psi_1(\eta_{G_1, g_1}, \dots, \eta_{G_v, g_v}) \\ &\times \xi_1(\eta_{G_1, g_1}) \dots \xi_v(\eta_{G_v, g_v}) z_1^{-\eta_{G_1, g_1}} \dots z_v^{-\eta_{G_v, g_v}} \end{aligned} \tag{1.6}$$

Where $\psi(\dots, \dots), \theta_i(\dots), i = 1, \dots, r$ are given respectively in (1.2), (1.3) and

$$\eta_{G_1, g_1} = \frac{d_{g_1}^{(1)} + G_1}{\delta_{g_1}^{(1)}}, \dots, \eta_{G_v, g_v} = \frac{d_{g_v}^{(v)} + G_v}{\delta_{g_v}^{(v)}}$$

which is valid under the conditions $\delta_{g_i}^{(i)}[d_j^i + p_i] \neq \delta_{g_i}^{(i)}[d_{g_i}^i + G_i]$ (1.7)

for $j \neq m_i, m_i = 1, \dots, \eta_{G_i, g_i}; p_i, n_i = 0, 1, 2, \dots; y_i \neq 0, i = 1, \dots, v$ (1.8)

The multivariable I-function of r -variables is defined in term of multiple Mellin-Barnes type integral :

$$I(z_1, \dots, z_r) = I_{p, q; p'_1, q'_1; \dots; p'_r, q'_r}^{0, n; m'_1, n'_1; \dots; m'_r, n'_r} \left(\begin{matrix} z_1 \\ \vdots \\ z_r \end{matrix} \middle| \begin{matrix} (a_j; \alpha_j^{(1)}, \dots, \alpha_j^{(r)}; A_j)_{1, p} : \\ \\ (b_j; \beta_j^{(1)}, \dots, \beta_j^{(r)}; B_j)_{1, q} : \end{matrix} \right)$$

$$\left(\begin{matrix} (c_j^{(1)}, \gamma_j^{(1)}; C_j^{(1)})_{1, p_1}; \dots; (c_j^{(r)}, \gamma_j^{(r)}; C_j^{(r)})_{1, p_r} \\ \\ (d_j^{(1)}, \delta_j^{(1)}; D_j^{(1)})_{1, q_1}; \dots; (d_j^{(r)}, \delta_j^{(r)}; D_j^{(r)})_{1, q_r} \end{matrix} \right) \tag{1.9}$$

$$= \frac{1}{(2\pi\omega)^r} \int_{L_1} \dots \int_{L_r} \phi(s_1, \dots, s_r) \prod_{i=1}^r \theta_i(s_i) z_i^{s_i} ds_1 \dots ds_r \tag{1.10}$$

where $\phi(s_1, \dots, s_r), \theta_i(s_i), i = 1, \dots, r$ are given by :

$$\phi(s_1, \dots, s_r) = \frac{\prod_{j=1}^n \Gamma^{A_j} (1 - a_j + \sum_{i=1}^r \alpha_j^{(i)} s_j)}{\prod_{j=n+1}^p \Gamma^{A_j} (a_j - \sum_{i=1}^r \alpha_j^{(i)} s_j) \prod_{j=1}^q \Gamma^{B_j} (1 - b_j + \sum_{i=1}^r \beta_j^{(i)} s_j)} \tag{1.11}$$

$$\theta_i(s_i) = \frac{\prod_{j=1}^{n'_i} \Gamma^{C_j^{(i)}} (1 - c_j^{(i)} + \gamma_j^{(i)} s_i) \prod_{j=1}^{m'_i} \Gamma^{D_j^{(i)}} (d_j^{(i)} - \delta_j^{(i)} s_i)}{\prod_{j=n'_i+1}^{p'_i} \Gamma^{C_j^{(i)}} (c_j^{(i)} - \gamma_j^{(i)} s_i) \prod_{j=m'_i+1}^{q'_i} \Gamma^{D_j^{(i)}} (1 - d_j^{(i)} + \delta_j^{(i)} s_i)} \tag{1.12}$$

For more details, see Nambisan et al [2].

Following the result of Braaksma [1] the I-function of r variables is analytic if :

$$U_i = \sum_{j=1}^p A_j \alpha_j^{(i)} - \sum_{j=1}^q B_j \beta_j^{(i)} + \sum_{j=1}^{p'_i} C_j^{(i)} \gamma_j^{(i)} - \sum_{j=1}^{q'_i} D_j^{(i)} \delta_j^{(i)} \leq 0, i = 1, \dots, r \tag{1.13}$$

The integral (2.1) converges absolutely if

$$|arg(z_k)| < \frac{1}{2} \Delta_k \pi, k = 1, \dots, r \text{ where}$$

$$\Delta_k = - \sum_{j=n+1}^p A_j \alpha_j^{(k)} - \sum_{j=1}^q B_j \beta_j^{(k)} + \sum_{j=1}^{m'_k} D_j^{(k)} \delta_j^{(k)} - \sum_{j=m'_k+1}^{q_k} D_j^{(k)} \delta_j^{(k)} + \sum_{j=1}^{n'_k} C_j^{(k)} \gamma_j^{(k)} - \sum_{j=n'_k+1}^{p_k} C_j^{(k)} \gamma_j^{(k)} > 0 \tag{1.14}$$

Srivastava and Garg [6] introduced and defined a general class of multivariable polynomials as follows

$$S_L^{h_1, \dots, h_u} [z_1, \dots, z_u] = \sum_{R_1, \dots, R_u=0}^{h_1 R_1 + \dots + h_u R_u \leq L} (-L)_{h_1 R_1 + \dots + h_u R_u} B(E; R_1, \dots, R_u) \frac{z_1^{R_1} \dots z_u^{R_u}}{R_1! \dots R_u!} \tag{1.15}$$

The coefficients are $B[E; R_1, \dots, R_u]$ arbitrary constants, real or complex.

2. Integral representation of generalized Lauricella function of several variables

The following generalized hypergeometric function in terms of multiple contour integrals is also required [7, page 39 eq.30]

$$\frac{\prod_{j=1}^P \Gamma(A_j)}{\prod_{j=1}^Q \Gamma(B_j)} {}_P F_Q [(A_P); (B_Q); -(x_1 + \dots + x_r)]$$

$$= \frac{1}{(2\pi\omega)^r} \int_{L_1} \dots \int_{L_r} \frac{\prod_{j=1}^P \Gamma(A_j + s_1 + \dots + s_r)}{\prod_{j=1}^Q \Gamma(B_j + s_1 + \dots + s_r)} \Gamma(-s_1) \dots \Gamma(-s_r) x_1^{s_1} \dots x_r^{s_r} ds_1 \dots ds_r \tag{2.1}$$

where the contours are of Barnes type with indentations, if necessary, to ensure that the poles of $\Gamma(A_j + s_1 + \dots + s_r)$ are separated from those of $\Gamma(-s_j), j = 1, \dots, r$. The above result (1.23) can be easily established by an appeal to the calculus of residues by calculating the residues at the poles of $\Gamma(-s_j), j = 1, \dots, r$

In order to evaluate a number of integrals of multivariable I-function, we first establish the formula

$$\int_a^b (t-a)^{\alpha-1} (b-t)^{\beta-1} \prod_{j=1}^l [1 - \tau_j (t-a)^{h_j}]^{-\lambda_j} \prod_{j=1}^k (f_j t + g_j)^{\sigma_j} dt = (b-a)^{\alpha+\beta-1} B(\alpha, \beta) \prod_{j=1}^k (af_j + g_j)^{\sigma_j}$$

$$F_{1:0, \dots, 0; 0, \dots, 0}^{1:1, \dots, 1; 1, \dots, 1} \left(\begin{matrix} (\alpha : h_1, \dots, h_l, 1, \dots, 1) : (\lambda_1 : 1), \dots, (\lambda_l : 1); (-\sigma_1 : 1), \dots, (-\sigma_k : 1) \\ \dots \\ (\alpha + \beta : h_1, \dots, h_l, 1, \dots, 1) : -, \dots, -; -, \dots, - \end{matrix} \right)$$

$$; \tau_1 (b-a)^{h_1}, \dots, \tau_l (b-a)^{h_l}, -\frac{(b-a)f_1}{af_1 + g_1}, \dots, -\frac{(b-a)f_k}{af_k + g_k} \tag{2.2}$$

where $a, b \in \mathbb{R}(a < b), \alpha, \beta, f_i, g_i, \sigma_i, \tau_j, h_j \in \mathbb{C}, \lambda_j \in \mathbb{R}^+(i = 1, \dots, k; j = 1, \dots, l)$

$$\min(\operatorname{Re}(\alpha), \operatorname{Re}(\beta)) > 0, \max_{1 \leq j \leq l} \{|\tau_j(b-a)^{h_j}|\} < 1, \max_{1 \leq j \leq k} \left\{ \left| \frac{(b-a)f_i}{af_i + g_i} \right| \right\} < 1,$$

and $F_{1:0, \dots, 0; 0, \dots, 0}^{1:1, \dots, 1; 1, \dots, 1}$ is a particular case of the generalized Lauricella function introduced by Srivastava-Daoust [5, page 454] given by :

$$F_{1:0, \dots, 0; 0, \dots, 0}^{1:1, \dots, 1; 1, \dots, 1} \left(\begin{matrix} (\alpha : h_1, \dots, h_l, 1, \dots, 1) : (\lambda_1 : 1), \dots, (\lambda_l : 1); (-\sigma_1 : 1), \dots, (-\sigma_k : 1) \\ \dots \\ (\alpha + \beta : h_1, \dots, h_l, 1, \dots, 1) : -, \dots, -; -, \dots, - \end{matrix} \right) ; \tau_1(b-a)^{h_1}, \dots, \tau_l(b-a)^{h_l}, -\frac{(b-a)f_1}{af_1 + g_1}, \dots, -\frac{(b-a)f_k}{af_k + g_k} = \frac{\Gamma(\alpha + \beta)}{\Gamma(\alpha) \prod_{j=1}^l \Gamma(\lambda_j) \prod_{j=1}^k \Gamma(-\sigma_j)}$$

$$\frac{1}{(2\pi\omega)^{l+k}} \int_{L_1} \dots \int_{L_{l+k}} \frac{\Gamma(\alpha + \sum_{j=1}^l h_j w_j + \sum_{j=1}^k w_{l+j})}{\Gamma(\alpha + \beta + \sum_{j=1}^l h_j w_j + \sum_{j=1}^k w_{l+j})} \prod_{j=1}^l \Gamma(\lambda_j + w_j) \prod_{j=1}^k \Gamma(-\sigma_j + w_{l+j})$$

$$\prod_{j=1}^{l+k} \Gamma(-w_j) z_1^{w_1} \dots z_l^{w_l} z_{l+1}^{w_{l+1}} \dots, w_{l+k}^{w_{l+k}} dw_1 \dots dw_{l+k} \tag{2.3}$$

Here the contour L'_j s are defined by $L_j = L_{w\zeta_j\infty}(\operatorname{Re}(\zeta_j) = v''_j)$ starting at the point $v''_j - \omega\infty$ and terminating at the point $v''_j + \omega\infty$ with $v''_j \in \mathbb{R}(j = 1, \dots, l)$ and each of the remaining contour L_{l+1}, \dots, L_{l+k} run from $-\omega\infty$ to $\omega\infty$

(2.2) can be easily established by expanding $\prod_{j=1}^l [1 - \tau_j(t-a)^{h_j}]^{-\lambda_j}$ by means of the formula :

$$(1-z)^{-\alpha} = \sum_{r=0}^{\infty} \frac{(\alpha)_r}{r!} z^r (|z| < 1) \tag{2.4}$$

integrating term by term with the help of the integral given by Saigo and Saxena [3, page 93, eq.(3.2)] and applying the definition of the generalized Lauricella function [5, page 454].

3. Eulerian integral

In this section , we evaluate a general Eulerian integral with the product of the multivariable Aleph-function, the multivariable I-function defined by Nambisan et al [2], a class of multivariable polynomials and generalized hypergeometric function. We note

$$B_u = \frac{(-L)_{h_1 R_1 + \dots + h_u R_u} B(E; R_1, \dots, R_u)}{R_1! \dots R_u!} \tag{3.1}$$

$$\text{and } B_{u,v} = (b-a)^{\sum_{i=1}^v (a'_i + b'_i) \eta_{G_i, g_i} + \sum_{i=1}^u (a_i + b_i) R_i} \left\{ \prod_{j=1}^h (af_j + g_j)^{-\sum_{i=1}^v \lambda''_i \eta_{g_i, h_i} - \sum_{i=1}^u \lambda'_i R_i} \right\} G_v \tag{3.2}$$

$$\text{where } G_v = \psi(\eta_{G_1, g_1}, \dots, \eta_{G_v, g_v}) \times \xi_1(\eta_{G_1, g_1}) \dots \xi_v(\eta_{G_v, g_v}) \tag{3.3}$$

$\psi_1, \xi_i, i = 1, \dots, v$ are defined respectively by (1.2) and (1.3)

$$\theta_i = \prod_{j=1}^l [1 - \tau_j(t-a)^{h_i}]^{-\zeta_j^{(i)}}, \zeta_j^{(i)} > 0 (i = 1, \dots, r); \theta'_i = \prod_{j=1}^l [1 - \tau_j(t-a)^{h_i}]^{-\zeta_j'^{(i)}}, \zeta_j'^{(i)} > 0 (i = 1, \dots, s)$$

$$\theta''_i = \prod_{j=1}^l [1 - \tau_j(t-a)^{h_i}]^{-\zeta_j''^{(i)}}, \zeta_j''^{(i)} > 0 (i = 1, \dots, u)$$

$$\theta'''_i = \prod_{j=1}^l [1 - \tau_j(t-a)^{h_i}]^{-\zeta_j'''^{(i)}}, \zeta_j'''^{(i)} > 0 (i = 1, \dots, v) \tag{3.4}$$

$$X = m'_1, n'_1; \dots; m'_r, n'_r; 1, 0; \dots; 1, 0; 1, 0; \dots; 1, 0; 1, 0; \dots; 1, 0 \tag{3.5}$$

$$Y = p'_1, q'_1; \dots; p'_r, q'_r; 0, 1; \dots; 0, 1; 0, 1; \dots; 0, 1; 0, 1; \dots; 0, 1 \tag{3.6}$$

$$A = (a_j; \alpha_j^{(1)}, \dots, \alpha_j^{(r)}, 0, \dots, 0, 0, \dots, 0, 0, \dots, 0; A_j)_{1,p} \tag{3.7}$$

$$B = (b_j; \beta_j^{(1)}, \dots, \beta_j^{(r)}, 0, \dots, 0, 0, \dots, 0, 0, \dots, 0; B_j)_{1,q} \tag{3.8}$$

$$C = (c_j^{(1)}, \gamma_j^{(1)}; C_j^{(1)})_{1,p_1}; \dots; (c_j^{(r)}, \gamma_j^{(r)}; C_j^{(r)})_{1,p_r}; (1, 0; 1) \dots; (1, 0; 1)$$

$$(1, 0; 1); \dots; (1, 0; 1); (1, 0; 1); \dots; (1, 0; 1) \tag{3.9}$$

$$D = (d_j^{(1)}, \delta_j^{(1)}; D_j^{(1)})_{1,q_1}; \dots; (d_j^{(r)}, \delta_j^{(r)}; D_j^{(r)})_{1,q_r}; (0, 1; 1); \dots; (0, 1; 1);$$

$$(0, 1; 1); \dots; (0, 1; 1); (0, 1; 1); \dots; (0, 1; 1) \tag{3.10}$$

$$K_1 = (1 - \alpha - \sum_{i=1}^u R_i a_i - \sum_{i=1}^v \eta_{G_i, g_i} a'_i; \mu_1, \dots, \mu_r, \mu'_1, \dots, \mu'_s, h_1, \dots, h_l, 1, \dots, 1; 1) \tag{3.11}$$

$$K_2 = (1 - \beta - \sum_{i=1}^u R_i b_i - \sum_{i=1}^v \eta_{G_i, g_i} b'_i; \rho_1, \dots, \rho_r, \rho'_1, \dots, \rho'_s, 0, \dots, 0, 0, \dots, 0; 1) \tag{3.12}$$

$$K_P = [1 - A_j; 0, \dots, 0, 1, \dots, 1, 0, \dots, 0, 0, \dots, 0; 1]_{1,P} \tag{3.13}$$

$$K_j = [1 - \lambda_j - \sum_{i=1}^u R_i \zeta_j''^{(i)} - \sum_{i=1}^v \eta_{G_i, g_i} \zeta_j'''^{(i)}; \zeta_j^{(1)}, \dots, \zeta_j^{(r)}, \zeta_j'^{(1)} \dots, \zeta_j'^{(s)},$$

$$0, \dots, 1, \dots, 0, 0, \dots, 0; 1]_{1,l} \tag{3.14}$$

$$K'_j = [1 + \sigma_j - \sum_{i=1}^u R_i \lambda_j''^{(i)} - \sum_{i=1}^v \eta_{G_i, g_i} \lambda_j'''^{(i)}; \lambda_j^{(1)}, \dots, \lambda_j^{(r)}, \lambda_j'^{(1)} \dots, \lambda_j'^{(s)},$$

$$0, \dots, 0, 0, \dots, 1, \dots, 0; 1]_{1,k} \tag{3.15}$$

$$L_1 = (1 - \alpha - \beta - \sum_{i=1}^u R_i(a_i + b_i) - \sum_{i=1}^v (a'_i + b'_i)\eta_{G_i, g_i}; \mu_1 + \rho_1, \dots, \mu_r + \rho_r, \mu'_1 + \rho'_1, \dots, \mu'_r + \rho'_r, h_1, \dots, h_u, 1, \dots, 1; 1) \tag{3.16}$$

$$L_Q = [1 - B_j; 0, \dots, 0, 1, \dots, 1, 0, \dots, 0, 0 \dots, 0; 1]_{1, Q} \tag{3.17}$$

$$L_j = [1 - \lambda_j - \sum_{i=1}^u R_i \zeta_j^{(i)} - \sum_{i=1}^s \zeta_j^{(i)} \eta_{G_i, g_i}; \zeta_j^{(1)}, \dots, \zeta_j^{(r)}, \zeta_j^{(1)}, \dots, \zeta_j^{(s)}, 0, \dots, 0, 0 \dots, 0; 1]_{1, l} \tag{3.18}$$

$$L'_j = [1 + \sigma_j - \sum_{i=1}^u R_i \lambda_j^{(i)} - \sum_{i=1}^v \lambda_j^{(i)} \eta_{G_i, g_i}; \lambda_j^{(1)}, \dots, \lambda_j^{(r)}, \lambda_j^{(1)}, \dots, \lambda_j^{(s)}, 0, \dots, 0, 0, \dots, 0; 1]_{1, k} \tag{3.19}$$

We have the general Eulerian integral

$$\int_a^b (t-a)^{\alpha-1} (b-t)^{\beta-1} \prod_{j=1}^l [1 - \tau_j(t-a)^{h_j}]^{-\lambda_j} \prod_{j=1}^k (f_j t + g_j)^{\sigma_j} S_L^{h_1, \dots, h_u} \left(\begin{matrix} z_1'' \theta_1'' (t-a)^{a_1} (b-t)^{b_1} \prod_{j=1}^k (f_j t + g_j)^{-\lambda_j^{(1)}} \\ \vdots \\ z_u'' \theta_u'' (t-a)^{a_u} (b-t)^{b_u} \prod_{j=1}^k (f_j t + g_j)^{-\lambda_j^{(u)}} \end{matrix} \right) \mathfrak{N} \left(\begin{matrix} z_1''' \theta_1''' (t-a)^{a'_1} (b-t)^{b'_1} \prod_{j=1}^k (f_j t + g_j)^{-\lambda_j^{(1)'}} \\ \vdots \\ z_v''' \theta_v''' (t-a)^{a'_v} (b-t)^{b'_v} \prod_{j=1}^k (f_j t + g_j)^{-\lambda_j^{(v)'}} \end{matrix} \right) I \left(\begin{matrix} z_1 \theta_1 (t-a)^{\mu_1} (b-t)^{\rho_1} \prod_{j=1}^k (f_j t + g_j)^{-\lambda_j^{(1)}} \\ \vdots \\ z_r \theta_r (t-a)^{\mu_r} (b-t)^{\rho_r} \prod_{j=1}^k (f_j t + g_j)^{-\lambda_j^{(r)}} \end{matrix} \right)$$

$${}_p F_Q \left[(A_P); (B_Q); - \sum_{i=1}^s z'_i \theta'_i (t-a)^{\mu'_i} (b-t)^{\rho'_i} \prod_{j=1}^k (f_j t + g_j)^{-\lambda_j^{(i)'}} \right] dt =$$

$$(b-a)^{\alpha+\beta-1} \frac{\prod_{j=1}^Q \Gamma(B_j)}{\prod_{j=1}^P \Gamma(A_j)} \prod_{j=1}^l (a f_j + g_j)^{\sigma_j} \sum_{G_1, \dots, G_v=0}^{\infty} \sum_{g_1=0}^{m_1} \dots \sum_{g_v=0}^{m_v} \sum_{R_1, \dots, R_u=0}^{h_1 R_1 + \dots + h_u R_u \leq L} \prod_{i=1}^v z_i^{\eta_{h_i, k_i}} \prod_{k=1}^u z''^{K_k} B_u B_{u,v}$$

$$I_{p+P+l+k+2,q+Q+l+k+1;Y}^{0,n+P+l+k+2;X} \left(\begin{array}{c|c} \frac{z_1(b-a)^{\mu_1+\rho_1}}{\prod_{j=1}^k (af_j+g_j)^{\lambda_j^{(1)}}} & \vdots \\ \dots & \vdots \\ \frac{z_r(b-a)^{\mu_r+\rho_r}}{\prod_{j=1}^k (af_j+g_j)^{\lambda_j^{(r)}}} & \vdots \\ \frac{z'_1(b-a)^{\mu'_1+\rho'_1}}{\prod_{j=1}^k (af_j+g_j)^{\lambda_j'^{(1)}}} & \vdots \\ \dots & \vdots \\ \frac{z'_s(b-a)^{\mu'_s+\rho'_s}}{\prod_{j=1}^k (af_j+g_j)^{\lambda_j'^{(s)}}} & \vdots \\ \tau_1(b-a)^{h_1} & \vdots \\ \dots & \vdots \\ \tau_l(b-a)^{h_l} & \vdots \\ \frac{(b-a)f_1}{af_1+g_1} & \vdots \\ \dots & \vdots \\ \frac{(b-a)f_k}{af_k+g_k} & \vdots \end{array} \right) \begin{array}{l} A ; K_1, K_2, K_3, K_j, K'_j : C \\ \\ \\ \\ \\ \\ \\ \\ B , L_1, L_2, L_j, L'_j : D \end{array} \quad (3.20)$$

This result is an extension the formula given by Saxena et al [4].

Provided that

- (A) $a, b \in \mathbb{R}(a < b); \mu_i, \mu'_u, \rho_i, \rho'_u, \lambda_j^{(i)}, \lambda_j^{(u)}, h_v \in \mathbb{R}^+, f_i, g_j, \tau_v, \sigma_j, \lambda_v \in \mathbb{C} (i = 1, \dots, r; j = 1, \dots, k; u = 1, \dots, s; v = 1, \dots, l), a_i, b_i, \lambda_j^{(i)}, \zeta_j^{(i)} \in \mathbb{R}^+, (i = 1, \dots, u; j = 1, \dots, k)$
- $a'_i, b'_i, \lambda_j^{(i)}, \zeta_j^{(i)} \in \mathbb{R}^+, (i = 1, \dots, v; j = 1, \dots, k)$
- (B) $m'_j, n'_j, p'_j, q'_j(j = 1, \dots, r), n, p, q \in \mathbb{N}^*; \delta_j^{(i)} \in \mathbb{R}_+(j = 1, \dots, q_i; i = 1, \dots, r)$
- $\alpha_j^{(i)} \in \mathbb{R}_+(j = 1, \dots, p; i = 1, \dots, r), \beta_j^{(i)} \in \mathbb{R}_+(j = 1, \dots, q; i = 1, \dots, r), \gamma_j^{(i)} \in \mathbb{R}_+(j = 1, \dots, p_i; i = 1, \dots, r)$
- $a_j(j = 1, \dots, p), b_j(j = 1, \dots, q), c_j^{(i)}(j = 1, \dots, p_i, i = 1, \dots, r), d_j^{(i)}(j = 1, \dots, q_i, i = 1, \dots, r) \in \mathbb{C}$

The exponents $A_j(j = 1, \dots, p), B_j(j = 1, \dots, q), C_j^{(i)}(j = 1, \dots, p_i; i = 1, \dots, r), D_j^{(i)}(j = 1, \dots, q_i; i = 1, \dots, r)$ of various gamma function involved in (1.3) and (1.4) may take non integer values.

- (C) $\max_{1 \leq j \leq k} \left\{ \left| \frac{(b-a)f_i}{af_i+g_i} \right| \right\} < 1$
- (D) $Re \left[\alpha + \sum_{i=1}^v a'_i \min_{1 \leq j \leq m_i} \frac{d_j^{(i)}}{\delta_j^{(i)}} + \sum_{i=1}^r \mu_i \min_{1 \leq j \leq m'_i} \frac{d_j^{(i)}}{\delta_j^{(i)}} \right] > 0$ and

$$Re \left[\beta + \sum_{i=1}^v b'_i \min_{1 \leq j \leq m_i} \frac{d_j^{(i)}}{\delta_j^{(i)}} + \sum_{i=1}^r \rho_i \min_{1 \leq j \leq m'_i} \frac{d'_j{}^{(i)}}{\delta'_j{}^{(i)}} \right] > 0$$

$$(E) Re \left(\alpha + \sum_{i=1}^v \eta_{G_i, g_i} a'_i + \sum_{i=1}^u R_i a_i + \sum_{i=1}^r \mu_i s_i + \sum_{i=1}^l h_i w_i \right) > 0 ; Re \left(\beta + \sum_{i=1}^v \eta_{G_i, g_i} b'_i + \sum_{i=1}^u R_i b_i + \sum_{i=1}^r \rho_i s_i \right) > 0$$

$$Re \left(\lambda_j + \sum_{i=1}^v \eta_{G_i, g_i} \lambda_j'''^{(i)} + \sum_{i=1}^u R_i \lambda_j''^{(i)} + \sum_{i=1}^r s_i \zeta_j^{(i)} \right) > 0 (j = 1, \dots, l)$$

$$Re \left(-\sigma_j + \sum_{i=1}^v \eta_{G_i, g_i} \lambda_j'''^{(i)} + \sum_{i=1}^u R_i \lambda_j''^{(i)} + \sum_{i=1}^r s_i \lambda_j^{(i)} \right) > 0 (j = 1, \dots, k)$$

$$(F) U_i = \sum_{j=1}^p A_j \alpha_j^{(i)} - \sum_{j=1}^q B_j \beta_j^{(i)} + \sum_{j=1}^{p'_i} C_j^{(i)} \gamma_j^{(i)} - \sum_{j=1}^{q'_i} D_j^{(i)} \delta_j^{(i)} \leq 0, i = 1, \dots, r$$

$$(G) \Delta_k = - \sum_{j=n+1}^p A_j \alpha_j^{(k)} - \sum_{j=1}^q B_j \beta_j^{(k)} + \sum_{j=1}^{m'_k} D_j^{(k)} \delta_j'^{(k)} - \sum_{j=m_k+1}^{q'_k} D_j^{(k)} \delta_j'^{(k)} + \sum_{j=1}^{n'_k} C_j^{(k)} \gamma_j^{(k)} - \sum_{j=n_k+1}^{p'_k} C_j^{(k)} \gamma_j^{(k)}$$

$$- \mu_i - \rho_i - \sum_{l=1}^k \lambda_l^{(i)} > 0 \quad (i = 1, \dots, r)$$

$$(H) \left| arg \left(z_i \prod_{j=1}^l [1 - \tau_j(t-a)^{h_i}]^{-\zeta_j^{(i)}} \prod_{j=1}^k (f_j t + g_j)^{-\lambda_j^{(i)}} \right) \right| < \frac{1}{2} \Delta_i \pi \quad (a \leq t \leq b; i = 1, \dots, r)$$

(I) $P \leq Q + 1$. The equality holds, when , in addition,

$$\text{either } P > Q \text{ and } \left| \left(z'_i \sum_{i=1}^l [1 - \tau_j(t-a)^{h_i}]^{-\zeta_j'^{(i)}} \prod_{j=1}^k (f_j t + g_j)^{-\lambda_j'^{(i)}} \right) \right|^{\frac{1}{Q-P}} < 1 \quad (a \leq t \leq b)$$

$$\text{or } P \leq Q \text{ and } \max_{1 \leq i \leq k} \left[\left| \left(z'_i \sum_{j=1}^l [1 - \tau_j(t-a)^{h_i}]^{-\zeta_j'^{(i)}} \prod_{j=1}^k (f_j t + g_j)^{-\lambda_j'^{(i)}} \right) \right| \right] < 1 \quad (a \leq t \leq b)$$

(J) The multiple series occurring on the right-hand side of (3.20) is absolutely and uniformly convergent.

Proof

To prove (3.20), first, we express in serie the multivariable Aleph-function with the help of (1.6), a class of multivariable polynomials defined by Srivastava et al [6] $S_L^{h_1, \dots, h_u} [.]$ in serie with the help of (1.15), the I-functions of r-variables defined by Nambisan et al [2] and in terms of Mellin-Barnes type contour integral with the help of (1.10), the generalized hypergeometric function ${}_pF_q(\cdot)$ in Mellin-Barnes contour integral with the help of (2.1). Now collect the power of $[1 - \tau_j(t-a)^{h_i}]$ with $(i = 1, \dots, r; j = 1, \dots, l)$ and collect the power of $(f_j t + g_j)$ with $j = 1, \dots, k$. Use the equations (2.2) and (2.3) and express the result in Mellin-Barnes contour integral. Interpreting the $(r + s + k + l)$ dimensional Mellin-Barnes integral to multivariable I-function, we obtain the equation (3.20).

4. Particular cases

a) If $A_j = B_j = C_j^{(i)} = D_j^{(i)} = 1$, the multivariable I-function defined by Nambisan et al [2] reduces to multivariable H-function defined by Srivastava et al [8]. We have the following result.

$$\int_a^b (t-a)^{\alpha-1} (b-t)^{\beta-1} \prod_{j=1}^l [1 - \tau_j(t-a)^{h_i}]^{-\lambda_j} \prod_{j=1}^k (f_j t + g_j)^{\sigma_j}$$

$$\begin{aligned}
 & S_L^{h_1, \dots, h_u} \begin{pmatrix} z_1'' \theta_1'' (t-a)^{a_1} (b-t)^{b_1} \prod_{j=1}^k (f_j t + g_j)^{-\lambda_j^{(1)}} \\ \vdots \\ z_u'' \theta_u'' (t-a)^{a_u} (b-t)^{b_u} \prod_{j=1}^k (f_j t + g_j)^{-\lambda_j^{(u)}} \end{pmatrix} \\
 & \mathfrak{N} \begin{pmatrix} z_1''' \theta_1''' (t-a)^{a'_1} (b-t)^{b'_1} \prod_{j=1}^k (f_j t + g_j)^{-\lambda_j^{(1)'}} \\ \vdots \\ z_v''' \theta_v''' (t-a)^{a'_v} (b-t)^{b'_v} \prod_{j=1}^k (f_j t + g_j)^{-\lambda_j^{(v)'}} \end{pmatrix} H \begin{pmatrix} z_1 \theta_1 (t-a)^{\mu_1} (b-t)^{\rho_1} \prod_{j=1}^k (f_j t + g_j)^{-\lambda_j^{(1)}} \\ \vdots \\ z_r \theta_r (t-a)^{\mu_r} (b-t)^{\rho_r} \prod_{j=1}^k (f_j t + g_j)^{-\lambda_j^{(r)}} \end{pmatrix} \\
 & = (b-a)^{\alpha+\beta-1} \frac{\prod_{j=1}^Q \Gamma(B_j)}{\prod_{j=1}^P \Gamma(A_j)} \prod_{j=1}^k (af_j + g_j)^{\sigma_j} \sum_{G_1, \dots, G_v=0}^{\infty} \sum_{g_1=0}^{m_1} \cdots \sum_{g_v=0}^{m_v} \sum_{R_1, \dots, R_u=0}^{h_1 R_1 + \dots + h_u R_u \leq L} \prod_{i=1}^v z_i^{\eta_{h_i, k_i}} \prod_{k=1}^u z''^{K_k} B_u B_{u,v} \\
 & H_{p+P+l+k+2, q+Q+l+k+1; Y}^{0, n+P+l+k+2; X} \left(\begin{array}{c|c} \frac{z_1 (b-a)^{\mu_1 + \rho_1}}{\prod_{j=1}^k (af_j + g_j)^{\lambda_j^{(1)}}} & \vdots \\ \vdots & \vdots \\ \frac{z_r (b-a)^{\mu_r + \rho_r}}{\prod_{j=1}^k (af_j + g_j)^{\lambda_j^{(r)}}} & \vdots \\ \frac{z'_1 (b-a)^{\mu'_1 + \rho'_1}}{\prod_{j=1}^k (af_j + g_j)^{\lambda_j^{(1)'}}} & \vdots \\ \vdots & \vdots \\ \frac{z'_s (b-a)^{\mu'_s + \rho'_s}}{\prod_{j=1}^k (af_j + g_j)^{\lambda_j^{(s)'}}} & \vdots \\ \tau_1 (b-a)^{h_1} & \vdots \\ \vdots & \vdots \\ \tau_l (b-a)^{h_l} & \vdots \\ \frac{(b-a)f_1}{af_1 + g_1} & \vdots \\ \vdots & \vdots \\ \frac{(b-a)f_k}{af_k + g_k} & \vdots \end{array} \right) \begin{array}{l} A ; K_1, K_2, K_3, K_j, K'_j : C \\ \\ \\ \\ \\ \\ \\ B , L_1, L_2, L_j, L'_j : D \end{array} \tag{4.1}
 \end{aligned}$$

under the same notations and conditions that (3.20) with $A_j = B_j = C_j^{(i)} = D_j^{(i)} = 1$:

$$b) \text{ If } B(L; R_1, \dots, R_u) = \frac{\prod_{j=1}^{\bar{A}} (a_j)_{R_1 \theta'_j + \dots + R_r \theta_j^{(r)}} \prod_{j=1}^{B'} (b'_j)_{R_1 \phi'_j} \dots \prod_{j=1}^{B^{(r)}} (b_j^{(r)})_{R_r \phi_j^{(r)}}}{\prod_{j=1}^{\bar{C}} (c_j)_{R_1 \psi'_j + \dots + R_r \psi_j^{(r)}} \prod_{j=1}^{D'} (d'_j)_{R_1 \delta'_j} \dots \prod_{j=1}^{D^{(r)}} (d_j^{(r)})_{R_r \delta_j^{(r)}}} \quad (4.2)$$

then the general class of multivariable polynomial $S_L^{h_1, \dots, h_u} [z_1, \dots, z_u]$ reduces to generalized Lauricella function defined by Srivastava et al [5]. We have

$$F_{\bar{C}:D'; \dots; D^{(u)}}^{1+\bar{A}:B'; \dots; B^{(u)}} \left(\begin{matrix} y_1 \\ \dots \\ y_u \end{matrix} \middle| \begin{matrix} [(-L); R_1, \dots, R_u] [(a); \theta', \dots, \theta^{(u)}] : [(b'); \phi']; \dots; [(b^{(u)}); \phi^{(u)}] \\ [(c); \psi', \dots, \psi^{(u)}] : [(d'); \delta']; \dots; [(d^{(u)}); \delta^{(u)}] \end{matrix} \right) \quad (4.3)$$

and we have the two following formulas

$$\int_a^b (t-a)^{\alpha-1} (b-t)^{\beta-1} \prod_{j=1}^l [1 - \tau_j (t-a)^{h_j}]^{-\lambda_j} \prod_{j=1}^k (f_j t + g_j)^{\sigma_j}$$

$$F_{\bar{C}:D'; \dots; D^{(u)}}^{1+\bar{A}:B'; \dots; B^{(u)}} \left(\begin{matrix} z_1'' \theta_1'' (t-a)^{a_1} (b-t)^{b_1} \prod_{j=1}^k (f_j t + g_j)^{-\lambda_j''(1)} \\ \vdots \\ z_u'' \theta_u'' (t-a)^{a_u} (b-t)^{b_u} \prod_{j=1}^k (f_j t + g_j)^{-\lambda_j''(u)} \end{matrix} \middle| \begin{matrix} [(-L); R_1, \dots, R_u] [(a); \theta', \dots, \theta^{(u)}] : [(b'); \phi']; \dots; [(b^{(u)}); \phi^{(u)}] \\ [(c); \psi', \dots, \psi^{(u)}] : [(d'); \delta']; \dots; [(d^{(u)}); \delta^{(u)}] \end{matrix} \right)$$

$$\aleph \left(\begin{matrix} z_1''' \theta_1''' (t-a)^{a'_1} (b-t)^{b'_1} \prod_{j=1}^k (f_j t + g_j)^{-\lambda_j'''(1)} \\ \vdots \\ z_v''' \theta_v''' (t-a)^{a'_v} (b-t)^{b'_v} \prod_{j=1}^k (f_j t + g_j)^{-\lambda_j'''(v)} \end{matrix} \right)$$

$$I \left(\begin{matrix} z_1 \theta_1 (t-a)^{\mu_1} (b-t)^{\rho_1} \prod_{j=1}^k (f_j t + g_j)^{-\lambda_j^{(1)}} \\ \vdots \\ z_r \theta_r (t-a)^{\mu_r} (b-t)^{\rho_r} \prod_{j=1}^k (f_j t + g_j)^{-\lambda_j^{(r)}} \end{matrix} \right)$$

$${}^pF_Q \left[(A_P); (B_Q); - \sum_{i=1}^s z_i' \theta_i'(t-a)^{\mu_i'} (b-t)^{\rho_i'} \prod_{j=1}^k (f_j t + g_j)^{-\lambda_j^{(i)}} \right] dt =$$

$$(b-a)^{\alpha+\beta-1} \frac{\prod_{j=1}^Q \Gamma(B_j)}{\prod_{j=1}^P \Gamma(A_j)} \prod_{j=1}^k (af_j + g_j)^{\sigma_j} \sum_{G_1, \dots, G_v=0}^{\infty} \sum_{g_1=0}^{m_1} \cdots \sum_{g_v=0}^{m_v} \sum_{R_1, \dots, R_u=0}^{h_1 R_1 + \dots + h_u R_u \leq L} \prod_{i=1}^v z_i''' \eta_{h_i, k_i} \prod_{k=1}^u z_i'' K_k B_u' B_{u,v}$$

$$I_{p+P+l+k+2, q+Q+l+k+1; Y}^{0, n+P+l+k+2; X} \left(\begin{array}{c|c} \frac{z_1 (b-a)^{\mu_1+\rho_1}}{\prod_{j=1}^k (af_j+g_j)^{\lambda_j^{(1)}}} & \vdots \\ \dots & \vdots \\ \frac{z_r (b-a)^{\mu_r+\rho_r}}{\prod_{j=1}^k (af_j+g_j)^{\lambda_j^{(r)}}} & \vdots \\ \frac{z'_1 (b-a)^{\mu'_1+\rho'_1}}{\prod_{j=1}^k (af_j+g_j)^{\lambda_j'^{(1)}}} & \vdots \\ \dots & \vdots \\ \frac{z'_s (b-a)^{\mu'_s+\rho'_s}}{\prod_{j=1}^k (af_j+g_j)^{\lambda_j'^{(s)}}} & \vdots \\ \tau_1 (b-a)^{h_1} & \vdots \\ \dots & \vdots \\ \tau_l (b-a)^{h_l} & \vdots \\ \frac{(b-a)f_1}{af_1+g_1} & \vdots \\ \dots & \vdots \\ \frac{(b-a)f_k}{af_k+g_k} & \vdots \end{array} \right) \begin{array}{l} A ; K_1, K_2, K_3, K_j, K'_j : C \\ \\ B , L_1, L_2, L_j, L'_j : D \end{array} \tag{4.4}$$

under the same notations and conditions that (3.20)

where $B'_u = \frac{(-L)_{h_1 R_1 + \dots + h_u R_u} B(E; R_1, \dots, R_u)}{R_1! \cdots R_u!}$, $B[E; R_1, \dots, R_u]$ is defined by (4.2)

Remark:

By the following similar procedure, the results of this document can be extended to product of any finite number of multivariable I-functions and a class of multivariable polynomials defined by Srivastava et al [5].

5. Conclusion

In this paper we have evaluated a generalized Eulerian integral involving the product of the multivariable Aleph-function, the multivariable I-function defined by Nambisan et al [2], a class of multivariable polynomials and generalized hypergeometric function with general arguments. The formulae established in this paper is very general nature. Thus, the results established in this research work would serve as a key formula from which, upon specializing the parameters, as many as desired results involving the special functions of one and several variables can be obtained.

REFERENCES

- [1] B. L. J. Braaksma, "Asymptotic expansions and analytic continuations for a class of Barnes integrals," *Compositio Mathematica*, vol. 15, pp. 239–341, 1964.
- [2] Prathima J. Nambisan V. and Kurumujji S.K. A Study of I-function of Several Complex Variables, *International Journal of Engineering Mathematics* Vol(2014) , 2014 page 1-12
- [3] Saigo M. and Saxena R.K. Unified fractional integral formulas for the multivariable H-function I. *J.Fractional Calculus* 15 (1999), page 91-107.
- [4] Saigo M. and Saxena R.K. Unified fractional integral formulas for the multivariable H-function III. *J.Fractional Calculus* 20 (2001), page 45-68.
- [5] Srivastava H.M. and Daoust M.C. Certain generalized Neumann expansions associated with Kampé de Fériet function. *Nederl. Akad. Wetensch. Proc. Ser A72 = Indag Math* 31(1969) page 449-457.
- [6] Srivastava H.M. And Garg M. Some integral involving a general class of polynomials and multivariable H-function. *Rev. Roumaine Phys.* 32(1987), page 685-692.
- [7] Srivastava H.M. and Karlsson P.W. *Multiple Gaussian Hypergeometric series*. Ellis.Horwood. Limited. New-York, Chichester. Brisbane. Toronto , 1985.
- [8] H.M. Srivastava And R.Panda. Some expansion theorems and generating relations for the H-function of several complex variables. *Comment. Math. Univ. St. Paul.* 24(1975), p.119-137.

Personal adress : 411 Avenue Joseph Raynaud
Le parc Fleuri , Bat B
83140 , Six-Fours les plages
Tel : 06-83-12-49-68
Department : VAR
Country : FRANCE