

On general Eulerian integral of certain products of I-function, the multivariable Aleph-function and a class of polynomials

F.Y. AYANT¹

¹ Teacher in High School , France

ABSTRACT

The object of this paper is to establish a general Eulerian integral involving the product of the I-function defined by Nambisan et al [1], the multivariable Aleph-function, a general class of multivariable polynomials and a generalized hypergeometric function which provide unification and extension of numerous results. We will study the particular case concerning the multivariable H-function defined by Srivastava et al [8] and the Srivastava-Daoust polynomial [5].

Keywords: Eulerian integral, multivariable Aleph-function, Lauricella function of several variables, multivariable I-function, generalized hypergeometric function, Srivastava-Daoust polynomial.

2010 Mathematics Subject Classification. 33C45, 33C60, 26D20

1.Introduction

In this paper, we evaluate a general Eulerian integral concerning the product of the multivariable Aleph-functions, the \bar{I} -function of several variables defined by Nambisan et al [1], a generalized hypergeometric function and a class of multivariable polynomials. We will give a series expansion of a multivariable \bar{I} -function.

The Aleph-function of several variables generalizes the multivariable I-function defined by Sharma and Ahmad [4], itself is a generalization of G and H-functions of several variables defined by Srivastava et al [8]. The multiple Mellin-Barnes integral occurring in this paper will be referred to as the multivariable Aleph-function throughout our present study and will be defined and represented as follows.

First time, we define the multivariable \bar{I} -function by :

$$I(z_1''', \dots, z_v''') = \bar{I}_{P,Q:P_1,Q_1;\dots;P_v,Q_v}^{0,N:M_1,N_1;\dots;M_v,N_v} \left(\begin{matrix} z_1''' \\ \cdot \\ \cdot \\ z_v''' \end{matrix} \middle| \begin{matrix} (a_j; \alpha_j^{(1)}, \dots, \alpha_j^{(v)}; A_j)_{N+1,P} : \\ \\ (b_j; \beta_j^{(1)}, \dots, \beta_j^{(v)}; B_j)_{M+1,Q} : \end{matrix} \right.$$

$$\left. \begin{matrix} (c_j^{(1)}, \gamma_j^{(1)}; 1)_{1,N_1}, (c_j^{(1)}, \gamma_j^{(1)}; C_j^{(1)})_{N_1+1,P_1}; \dots; (c_j^{(v)}, \gamma_j^{(v)}; 1)_{1,N_v}, (c_j^{(v)}, \gamma_j^{(v)}; C_j^{(v)})_{N_v+1,P_v} \\ (d_j^{(1)}, \delta_j^{(1)}; 1)_{1,M_1}, (\bar{d}_j^{(1)}, \delta_j^{(1)}; D_j^{(1)})_{M_1+1,Q_1}; \dots; (\bar{d}_j^{(v)}, \delta_j^{(v)}; 1)_{1,M_v}, (\bar{d}_j^{(v)}, \delta_j^{(v)}; D_j^{(v)})_{M_v+1,Q_v} \end{matrix} \right) \quad (1.1)$$

$$= \frac{1}{(2\pi\omega)^v} \int_{L_1} \dots \int_{L_v} \phi_1(s_1, \dots, s_v) \prod_{i=1}^v \xi_i'(s_i) z_i''' s_i ds_1 \dots ds_v \quad (1.2)$$

where $\phi_1(s_1, \dots, s_v), \xi_i'(s_i), i = 1, \dots, v$ are given by :

$$\phi_1(s_1, \dots, s_v) = \frac{1}{\prod_{j=N+1}^P \Gamma^{A_j} \left(a_j - \sum_{i=1}^v \alpha_j^{(i)} s_j \right) \prod_{j=M+1}^Q \Gamma^{B_j} \left(1 - b_j + \sum_{i=1}^v \beta_j^{(i)} s_j \right)} \quad (1.3)$$

$$\xi'_i(s_i) = \frac{\prod_{j=1}^{N_i} \Gamma(1 - c_j^{(i)} + \gamma_j^{(i)} s_i) \prod_{j=1}^{M_i} \Gamma(\bar{d}_j^{(i)} - \delta_j^{(i)} s_i)}{\prod_{j=N_i+1}^{P_i} \Gamma C_j^{(i)} (c_j^{(i)} - \gamma_j^{(i)} s_i) \prod_{j=M_i+1}^{Q_i} \Gamma D_j^{(i)} (1 - \bar{d}_j^{(i)} + \delta_j^{(i)} s_i)} \tag{1.4}$$

$i = 1, \dots, v$

Series representation

If $z'''_i \neq 0; i = 1, \dots, v$

$\delta_{h_i}^{(i)} (\bar{d}_j^{(i)} + k_i) \neq \bar{d}_j^{(i)} (\delta_{h_i}^{(i)} + \eta_i)$ for $j \neq h_i, j, h_i = 1, \dots, m_i (i = 1, \dots, v), k_i, \eta_i = 0, 1, 2, \dots (i = 1, \dots, v)$, then

$$\bar{I}(z'''_1, \dots, z'''_v) = \sum_{h_1=1}^{M_1} \dots \sum_{h_v=1}^{M_v} \sum_{k_1=0}^{\infty} \dots \sum_{k_v=0}^{\infty} \left[\phi_1 \left(\frac{dh_1^{(1)} + k_1}{\delta h_1^{(1)}}, \dots, \frac{dh_v^{(v)} + k_v}{\delta h_v^{(v)}} \right) \right] \prod_{j \neq h_i, i=1}^r \frac{(-)^{k_i}}{\delta h_i^{(i)} k_i!} z'''_i \frac{dh_i + k_i}{\delta h_i} \tag{1.5}$$

This result can be proved on computing the residues at the poles :

$$s_i = \frac{dh_i^{(i)} + k_i}{\delta h_i^{(i)}}, (h_i = 1, \dots, m_i, k_i = 0, 1, 2, \dots) \text{ for } i = 1, \dots, v \tag{1.6}$$

We may establish the asymptotic expansion in the following convenient form :

$$\bar{I}(z'''_1, \dots, z'''_v) = O(|z'''_1|^{\alpha_1}, \dots, |z'''_v|^{\alpha_v}), \max(|z'''_1|, \dots, |z'''_v|) \rightarrow 0$$

$$I(z'''_1, \dots, z'''_v) = O(|z'''_1|^{\beta_1}, \dots, |z'''_v|^{\beta_v}), \min(|z'''_1|, \dots, |z'''_v|) \rightarrow \infty$$

where $k = 1, \dots, v : \alpha_k = \min[Re(\bar{d}_j^{(k)} / \delta_j^{(k)})], j = 1, \dots, m_k$ and

$$\beta_k = \max[Re((c_j^{(k)} - 1) / \gamma_j^{(k)})], j = 1, \dots, n_k$$

$$\text{We will note } \eta_{h_i, k_i} = \frac{dh_i^{(i)} + k_i}{\delta h_i^{(i)}}, (h_i = 1, \dots, m_i, k_i = 0, 1, 2, \dots) \text{ for } i = 1, \dots, v \tag{1.7}$$

We define : $\aleph(z_1, \dots, z_r) = \aleph_{p'_i, q'_i, \tau'_i; R': p'_{i(1)}, q'_{i(1)}, \tau'_{i(1)}; R^{(1)}; \dots; p'_{i(r)}, q'_{i(r)}, \tau'_{i(r)}; R^{(r)}}^{0, n': m'_1, n'_1, \dots, m'_r, n'_r} \left(\begin{array}{c} z_1 \\ \vdots \\ z_r \end{array} \right)$

$$[(a'_j; \alpha'_j)^{(1)}, \dots, \alpha'_j)^{(r)}]_{1, n'} : [\tau'_i(a_{ji}; \alpha'_{ji})^{(1)}, \dots, \alpha'_{ji})^{(r)}]_{n'+1, p'_i} : \\ \dots \dots \dots [\tau'_i(b_{ji}; \beta'_{ji})^{(1)}, \dots, \beta'_{ji})^{(r)}]_{m'+1, q'_i} :$$

$$\left[(c_j^{(1)}; \gamma_j^{(1)})_{1, n'_1}, [\tau'_{i(1)}(c'_{ji(1)}; \gamma'_{ji(1)})_{n'_1+1, p'_{i(1)}}]; \dots; [(c_j^{(r)}; \gamma_j^{(r)})_{1, n'_r}, [\tau'_{i(r)}(c'_{ji(r)}; \gamma'_{ji(r)})_{n'_r+1, p'_{i(r)}}] \right] \\ \left[(d_j^{(1)}; \delta_j^{(1)})_{1, m'_1}, [\tau'_{i(1)}(d'_{ji(1)}; \delta'_{ji(1)})_{m'_1+1, q'_{i(1)}}]; \dots; [(d_j^{(r)}; \delta_j^{(r)})_{1, m'_r}, [\tau'_{i(r)}(d'_{ji(r)}; \delta'_{ji(r)})_{m'_r+1, q'_{i(r)}}] \right]$$

$$= \frac{1}{(2\pi\omega)^r} \int_{L_1} \dots \int_{L_r} \psi(s_1, \dots, s_r) \prod_{k=1}^r \theta_k(s_k) z_k^{s_k} ds_1 \dots ds_r \tag{1.8}$$

with $\omega = \sqrt{-1}$

$$\psi(s_1, \dots, s_r) = \frac{\prod_{j=1}^{n'} \Gamma(1 - a'_j + \sum_{k=1}^r \alpha'_{j(k)} s_k)}{\sum_{i=1}^R [\tau'_i \prod_{j=n'+1}^{p'_i} \Gamma(a'_{ji} - \sum_{k=1}^r \alpha'_{ji(k)} s_k) \prod_{j=1}^{q'_i} \Gamma(1 - b'_{ji} + \sum_{k=1}^r \beta'_{ji(k)} s_k)]} \quad (1.9)$$

and

$$\theta_k(s_k) = \frac{\prod_{j=1}^{m'_k} \Gamma(d'_j(k) - \delta'_j(k) s_k) \prod_{j=1}^{n'_k} \Gamma(1 - c'_j(k) + \gamma'_j(k) s_k)}{\sum_{i^{(k)}=1}^{R^{(k)}} [\tau'_{i^{(k)}} \prod_{j=m'_k+1}^{q'_{i^{(k)}}} \Gamma(1 - d'_{ji^{(k)}}(k) + \delta'_{ji^{(k)}}(k) s_k) \prod_{j=n'_k+1}^{p'_{i^{(k)}}} \Gamma(c'_{ji^{(k)}}(k) - \gamma'_{ji^{(k)}}(k) s_k)]} \quad (1.10)$$

Suppose, as usual, that the parameters

$$a'_j, j = 1, \dots, p'; b_j, j = 1, \dots, q';$$

$$c'_j(k), j = 1, \dots, n'_k; c'_{ji^{(k)}}, j = n'_k + 1, \dots, p'_{i^{(k)}};$$

$$d'_{ji^{(k)}}, j = m'_k + 1, \dots, q_{i^{(k)}}; d'_j(k), j = 1, \dots, m'_k;$$

$$\text{with } k = 1 \dots, r, i = 1, \dots, R', i^{(k)} = 1, \dots, R^{(k)}$$

are complex numbers, and the $\alpha' s, \beta' s, \gamma' s$ and $\delta' s$ are assumed to be positive real numbers for standardization purpose such that

$$U_i^{(k)} = \sum_{j=1}^{n'} \alpha'_j(k) + \tau'_i \sum_{j=n'+1}^{p'_i} \alpha'_{ji}(k) + \sum_{j=1}^{n'_k} \gamma'_j(k) + \tau'_{i^{(k)}} \sum_{j=n'_k+1}^{p'_{i^{(k)}}} \gamma'_{ji^{(k)}}(k) - \tau'_i \sum_{j=1}^{q'_i} \beta'_{ji}(k) - \sum_{j=1}^{m'_k} \delta'_j(k) - \tau'_{i^{(k)}} \sum_{j=m'_k+1}^{q'_{i^{(k)}}} \delta'_{ji^{(k)}}(k) \leq 0 \quad (1.11)$$

The reals numbers τ_i are positives for $i = 1$ to R' , $\tau_{i^{(k)}}$ are positives for $i^{(k)} = 1$ to $R^{(k)}$

The contour L_k is in the s_k -p lane and run from $\sigma - i\infty$ to $\sigma + i\infty$ where σ is a real number with loop, if necessary, ensure that the poles of $\Gamma(d'_j(k) - \delta'_j(k) s_k)$ with $j = 1$ to m_k are separated from those $\Gamma(1 - a'_j + \sum_{i=1}^r \alpha'_{ji}(k) s_k)$ of with $j = 1$ to n and $\Gamma(1 - c'_j(k) + \gamma'_j(k) s_k)$ with $j = 1$ to n_k to the left of the contour L_k . The condition for absolute convergence of multiple Mellin-Barnes type contour (1.9) can be obtained by extension of the corresponding conditions for multivariable H-function given by as :

$$|arg z_k| < \frac{1}{2} A_i^{(k)} \pi, \text{ where}$$

$$A_i^{(k)} = \sum_{j=1}^{n'} \alpha'_j(k) - \tau'_i \sum_{j=n'+1}^{p'_i} \alpha'_{ji}(k) - \tau'_{i^{(k)}} \sum_{j=1}^{n'_k} \beta'_{ji}(k) + \sum_{j=1}^{n'_k} \gamma'_j(k) - \tau'_{i^{(k)}} \sum_{j=n'_k+1}^{p'_{i^{(k)}}} \gamma'_{ji^{(k)}}(k) + \sum_{j=1}^{m'_k} \delta'_j(k) - \tau'_{i^{(k)}} \sum_{j=m'_k+1}^{q'_{i^{(k)}}} \delta'_{ji^{(k)}}(k) > 0, \text{ with } k = 1, \dots, r, i = 1, \dots, R', i^{(k)} = 1, \dots, R^{(k)} \quad (1.12)$$

The complex numbers z_i are not zero. Throughout this document, we assume the existence and absolute convergence conditions of the multivariable Aleph-function.

We may establish the asymptotic expansion in the following convenient form :

$$\aleph(z_1, \dots, z_r) = O(|z_1|^{\alpha_1}, \dots, |z_r|^{\alpha_r}), \max(|z_1|, \dots, |z_r|) \rightarrow 0$$

$$\aleph(z_1, \dots, z_r) = O(|z_1|^{\beta_1}, \dots, |z_r|^{\beta_r}), \min(|z_1|, \dots, |z_r|) \rightarrow \infty$$

where $k = 1, \dots, r : \alpha'_k = \min[Re(d'_j^{(k)} / \delta'_j^{(k)})], j = 1, \dots, m'_k$ and

$$\beta'_k = \max[Re((c'_j)^{(k)} - 1) / \gamma'_j^{(k)}], j = 1, \dots, n'_k$$

We will use these following notations in this paper

$$U = p'_i, q'_i, \tau'_i; R' ; V = m'_1, n'_1; \dots; m'_r, n'_r \tag{1.13}$$

$$W = p'_{i(1)}, q'_{i(1)}, \tau'_{i(1)}; R^{(1)}, \dots, p'_{i(r)}, q'_{i(r)}, \tau'_{i(r)}; R^{(r)} \tag{1.14}$$

$$A = \{(a'_j; \alpha'_j, \dots, \alpha'_j)^{(1)}\}_{1, n'_1}, \{\tau'_i(a'_{ji}; \alpha'_{ji}^{(1)}, \dots, \alpha'_{ji}^{(r)})\}_{n'_1+1, p'_{i(1)}} \tag{1.15}$$

$$B = \{\tau'_i(b'_{ji}; \beta'_{ji}^{(1)}, \dots, \beta'_{ji}^{(r)})\}_{m'_1+1, q'_i} \tag{1.16}$$

$$C = \{(c'_j)^{(1)}; \gamma'_j^{(1)}\}_{1, n'_1}, \tau'_{i(1)}(c'_{ji(1)}; \gamma'_{ji(1)})_{n'_1+1, p'_{i(1)}} \tag{1.17}$$

$$\{(c'_j)^{(r)}; \gamma'_j^{(r)}\}_{1, n'_r}, \tau'_{i(r)}(c'_{ji(r)}; \gamma'_{ji(r)})_{n'_r+1, p'_{i(r)}}$$

$$D = \{(d'_j)^{(1)}; \delta'_j^{(1)}\}_{1, m'_1}, \tau'_{i(1)}(d'_{ji(1)}; \delta'_{ji(1)})_{m'_1+1, q'_{i(1)}}, \dots \tag{1.18}$$

$$, \{(d'_j)^{(r)}; \delta'_j^{(r)}\}_{1, m'_r}, \tau'_{i(r)}(d'_{ji(r)}; \delta'_{ji(r)})_{m'_r+1, q'_{i(r)}}$$

The multivariable Aleph-function write :

$$\aleph(z_1, \dots, z_r) = \aleph_{U; V}^{0, n'; V} \left(\begin{matrix} z_1 \\ \cdot \\ \cdot \\ \cdot \\ z_r \end{matrix} \middle| \begin{matrix} A : C \\ \cdot \\ \cdot \\ B : D \end{matrix} \right) \tag{1.19}$$

Srivastava and Garg [6] introduced and defined a general class of multivariable polynomials as follows

$$S_L^{h_1, \dots, h_u} [z_1, \dots, z_u] = \sum_{R_1, \dots, R_u=0}^{h_1 R_1 + \dots + h_u R_u \leq L} (-L)_{h_1 R_1 + \dots + h_u R_u} B(E; R_1, \dots, R_u) \frac{z_1^{R_1} \dots z_u^{R_u}}{R_1! \dots R_u!} \tag{1.20}$$

The coefficients are $B[E; R_1, \dots, R_u]$ arbitrary constants, real or complex.

2. Integral representation of generalized Lauricella function of several variables

The following generalized hypergeometric function in terms of multiple contour integrals is also required [7, page 39 eq .30]

$$\frac{\prod_{j=1}^P \Gamma(A_j)}{\prod_{j=1}^Q \Gamma(B_j)} {}_P F_Q [(A_P); (B_Q); -(x_1 + \dots + x_r)]$$

$$= \frac{1}{(2\pi\omega)^r} \int_{L_1} \dots \int_{L_r} \frac{\prod_{j=1}^P \Gamma(A_j + s_1 + \dots + s_r)}{\prod_{j=1}^Q \Gamma(B_j + s_1 + \dots + s_r)} \Gamma(-s_1) \dots \Gamma(-s_r) x_1^{s_1} \dots x_r^{s_r} ds_1 \dots ds_r \quad (2.1)$$

where the contours are of Barnes type with indentations, if necessary, to ensure that the poles of $\Gamma(A_j + s_1 + \dots + s_r)$ are separated from those of $\Gamma(-s_j), j = 1, \dots, r$. The above result (1.23) can be easily established by an appeal to the calculus of residues by calculating the residues at the poles of $\Gamma(-s_j), j = 1, \dots, r$

In order to evaluate a number of integrals of multivariable I-function, we first establish the formula

$$\int_a^b (t-a)^{\alpha-1} (b-t)^{\beta-1} \prod_{j=1}^l [1 - \tau_j (t-a)^{h_j}]^{-\lambda_j} \prod_{j=1}^k (f_j t + g_j)^{\sigma_j} dt = (b-a)^{\alpha+\beta-1} B(\alpha, \beta) \prod_{j=1}^k (af_j + g_j)^{\sigma_j}$$

$$F_{1:0, \dots, 0; 0, \dots, 0}^{1:1, \dots, 1; 1, \dots, 1} \left(\begin{matrix} (\alpha : h_1, \dots, h_l, 1, \dots, 1) : (\lambda_1 : 1), \dots, (\lambda_l : 1); (-\sigma_1 : 1), \dots, (-\sigma_k : 1) \\ \dots \\ (\alpha + \beta : h_1, \dots, h_l, 1, \dots, 1) : -, \dots, -; -, \dots, - \end{matrix} \right)$$

$$; \tau_1 (b-a)^{h_1}, \dots, \tau_l (b-a)^{h_l}, -\frac{(b-a)f_1}{af_1 + g_1}, \dots, -\frac{(b-a)f_k}{af_k + g_k} \quad (2.2)$$

where $a, b \in \mathbb{R} (a < b), \alpha, \beta, f_i, g_i, \sigma_i, \tau_j, h_j \in \mathbb{C}, \lambda_j \in \mathbb{R}^+ (i = 1, \dots, k; j = 1, \dots, l)$

$$\min(Re(\alpha), Re(\beta)) > 0, \max_{1 \leq j \leq l} \{|\tau_j (b-a)^{h_j}|\} < 1, \max_{1 \leq j \leq k} \left\{ \left| \frac{(b-a)f_j}{af_j + g_j} \right| \right\} < 1,$$

and $F_{1:0, \dots, 0; 0, \dots, 0}^{1:1, \dots, 1; 1, \dots, 1}$ is a particular case of the generalized Lauricella function introduced by Srivastava-Daoust[5, page 454] given by :

$$F_{1:0, \dots, 0; 0, \dots, 0}^{1:1, \dots, 1; 1, \dots, 1} \left(\begin{matrix} (\alpha : h_1, \dots, h_l, 1, \dots, 1) : (\lambda_1 : 1), \dots, (\lambda_l : 1); (-\sigma_1 : 1), \dots, (-\sigma_k : 1) \\ \dots \\ (\alpha + \beta : h_1, \dots, h_l, 1, \dots, 1) : -, \dots, -; -, \dots, - \end{matrix} \right)$$

$$; \tau_1 (b-a)^{h_1}, \dots, \tau_l (b-a)^{h_l}, -\frac{(b-a)f_1}{af_1 + g_1}, \dots, -\frac{(b-a)f_k}{af_k + g_k} \Bigg) = \frac{\Gamma(\alpha + \beta)}{\Gamma(\alpha) \prod_{j=1}^l \Gamma(\lambda_j) \prod_{j=1}^k \Gamma(-\sigma_j)}$$

$$\frac{1}{(2\pi\omega)^{l+k}} \int_{L_1} \dots \int_{L_{l+k}} \frac{\Gamma(\alpha + \sum_{j=1}^l h_j s_j + \sum_{j=1}^k s_{l+j})}{\Gamma(\alpha + \beta + \sum_{j=1}^l h_j s_j + \sum_{j=1}^k s_{l+j})} \prod_{j=1}^l \Gamma(\lambda_j + s_j) \prod_{j=1}^k \Gamma(-\sigma_j + s_{l+j})$$

$$\prod_{j=1}^{l+k} \Gamma(-s_j) z_1^{s_1} \dots z_l^{s_l} z_{l+1}^{s_{l+1}} \dots, z_{l+k}^{s_{l+k}} ds_1 \dots ds_{l+k} \quad (2.3)$$

Here the contour L'_j s are defined by $L_j = L_{w\zeta_j\infty}(Re(\zeta_j) = v''_j)$ starting at the point $v''_j - \omega\infty$ and terminating at the point $v''_j + \omega\infty$ with $v''_j \in \mathbb{R}(j = 1, \dots, l)$ and each of the remaining contour L_{l+1}, \dots, L_{l+k} run from $-\omega\infty$ to $\omega\infty$

(2.2) can be easily established by expanding $\prod_{j=1}^l [1 - \tau_j(t-a)^{h_j}]^{-\lambda_j}$ by means of the formula :

$$(1-z)^{-\alpha} = \sum_{r=0}^{\infty} \frac{(\alpha)_r}{r!} z^r (|z| < 1) \tag{2.4}$$

integrating term by term with the help of the integral given by Saigo and Saxena [2, page 93, eq.(3.2)] and applying the definition of the generalized Lauricella function [5, page 454].

3. Eulerian integral

In this section , we evaluate a general Eulerian integral with the product of the multivariable Aleph-function, the multivariable \bar{I} -function of Nambisan, a class of multivariable polynomials and generalized hypergeometric function; We note

$$B_u = \frac{(-L)_{h_1 R_1 + \dots + h_u R_u} B(E; R_1, \dots, R_u)}{R_1! \dots R_u!} \tag{3.1}$$

$$\text{and } B_{u,v} = (b-a)^{\sum_{i=1}^v (a'_i + b'_i) \eta_{G_i, g_i} + \sum_{i=1}^u (a_i + b_i) R_i} \left\{ \prod_{j=1}^h (af_j + g_j)^{-\sum_{i=1}^v \lambda_i''' \eta_{g_i, h_i} - \sum_{i=1}^u \lambda_i'' R_i} \right\} \tag{3.2}$$

$$\theta_i = \prod_{j=1}^l [1 - \tau_j(t-a)^{h_j}]^{-\zeta_j^{(i)}}, \zeta_j^{(i)} > 0 (i = 1, \dots, r); \theta'_i = \prod_{j=1}^l [1 - \tau_j(t-a)^{h_j}]^{-\zeta_j'^{(i)}}, \zeta_j'^{(i)} > 0 (i = 1, \dots, s) \tag{3.3}$$

$$\theta''_i = \prod_{j=1}^l [1 - \tau_j(t-a)^{h_j}]^{-\zeta_j''^{(i)}}, \zeta_j''^{(i)} > 0 (i = 1, \dots, u) \tag{3.4}$$

$$\theta'''_i = \prod_{j=1}^l [1 - \tau_j(t-a)^{h_j}]^{-\zeta_j'''^{(i)}}, \zeta_j'''^{(i)} > 0 (i = 1, \dots, v) \tag{3.5}$$

$$K_1 = (1 - \alpha - \sum_{i=1}^u R_i a_i - \sum_{i=1}^v \eta_{G_i, g_i} a'_i; \mu_1, \dots, \mu_r, \mu'_1, \dots, \mu'_s, h_1, \dots, h_l, 1, \dots, 1) \tag{3.6}$$

$$K_2 = (1 - \beta - \sum_{i=1}^u R_i b_i - \sum_{i=1}^v \eta_{G_i, g_i} b'_i; \rho_1, \dots, \rho_r, \rho'_1, \dots, \rho'_s, 0, \dots, 0, 0 \dots, 0) \tag{3.7}$$

$$K_P = [1 - A_j; 0, \dots, 0, 1, \dots, 1, 0, \dots, 0, 0, \dots, 0]_{1,P} \tag{3.8}$$

$$K_j = [1 - \lambda_j - \sum_{i=1}^u R_i \zeta_j''^{(i)} - \sum_{i=1}^v \eta_{G_i, g_i} \zeta_j'''^{(i)}; \zeta_j^{(1)}, \dots, \zeta_j^{(r)}, \zeta_j'^{(1)}, \dots, \zeta_j'^{(s)}, 0, \dots, 1, \dots, 0, 0 \dots, 0]_{1,l} \tag{3.9}$$

$$K'_j = [1 + \sigma_j - \sum_{i=1}^u R_i \lambda_j''^{(i)} - \sum_{i=1}^v \eta_{G_i, g_i} \lambda_j'''^{(i)}; \lambda_j^{(1)}, \dots, \lambda_j^{(r)}, \lambda_j'^{(1)}, \dots, \lambda_j'^{(s)}, 0, \dots, 0, 0 \dots, 1, \dots, 0]_{1,k} \tag{3.9}$$

$$L_1 = (1 - \alpha - \beta - \sum_{i=1}^u R_i(a_i + b_i) - \sum_{i=1}^v (a'_i + b'_i)\eta_{G_i, g_i}; \mu_1 + \rho_1, \dots, \mu_r + \rho_r, \mu'_1 + \rho'_1, \dots, \mu'_r + \rho'_r, h_1, \dots, h_l, 1, \dots, 1) \tag{3.10}$$

$$L_Q = [1 - B_j; 0, \dots, 0, 1, \dots, 1, 0, \dots, 0, 0 \dots, 0]_{1, Q} \tag{3.11}$$

$$L_j = [1 - \lambda_j - \sum_{i=1}^u R_i \zeta_j^{(i)} - \sum_{i=1}^s \zeta_j^{(i)} \eta_{G_i, g_i}; \zeta_j^{(1)}, \dots, \zeta_j^{(r)}, \zeta_j^{(1)} \dots, \zeta_j^{(s)}, 0, \dots, 0, 0 \dots, 0]_{1, l} \tag{3.12}$$

$$L'_j = [1 + \sigma_j - \sum_{i=1}^u R_i \lambda_j^{(i)} - \sum_{i=1}^v \lambda_j^{(i)} \eta_{G_i, g_i}; \lambda_j^{(1)}, \dots, \lambda_j^{(r)}, \lambda_j^{(1)} \dots, \lambda_j^{(s)}, 0, \dots, 0, 0, \dots, 0]_{1, k} \tag{3.13}$$

$$V_1 = V; 1, 0; \dots; 1, 0; 1, 0; \dots; 1, 0; 1, 0; \dots; 1, 0; W_1 = W; 0, 1; \dots; 0, 1; 0, 1; \dots; 0, 1; 0, 1; \dots; 0, 1 \tag{3.14}$$

$$C_1 = C; (1, 0), \dots, (1, 0); (1, 0), \dots, (1, 0); (1, 0); \dots; (1, 0);$$

$$D_1 = D; (0, 1), \dots, (0, 1); (0, 1), \dots, (0, 1); (0, 1); \dots; (0, 1) \tag{3.15}$$

V, W, C and D are defined by (1.13), (1.14), (1.17) and (1.18) respectively

We have the general Eulerian integral

$$\int_a^b (t-a)^{\alpha-1} (b-t)^{\beta-1} \prod_{j=1}^l [1 - \tau_j(t-a)^{h_j}]^{-\lambda_j} \prod_{j=1}^k (f_j t + g_j)^{\sigma_j} S_L^{h_1, \dots, h_u} \left(\begin{matrix} z_1'' \theta_1'' (t-a)^{a_1} (b-t)^{b_1} \prod_{j=1}^k (f_j t + g_j)^{-\lambda_j^{(1)}} \\ \vdots \\ z_u'' \theta_u'' (t-a)^{a_u} (b-t)^{b_u} \prod_{j=1}^k (f_j t + g_j)^{-\lambda_j^{(u)}} \end{matrix} \right) \bar{I} \left(\begin{matrix} z_1''' \theta_1''' (t-a)^{a'_1} (b-t)^{b'_1} \prod_{j=1}^k (f_j t + g_j)^{-\lambda_j^{(1)'}} \\ \vdots \\ z_v''' \theta_v''' (t-a)^{a'_v} (b-t)^{b'_v} \prod_{j=1}^k (f_j t + g_j)^{-\lambda_j^{(v)'}} \end{matrix} \right) \aleph \left(\begin{matrix} z_1 \theta_1 (t-a)^{\mu_1} (b-t)^{\rho_1} \prod_{j=1}^k (f_j t + g_j)^{-\lambda_j^{(1)}} \\ \vdots \\ z_r \theta_r (t-a)^{\mu_r} (b-t)^{\rho_r} \prod_{j=1}^k (f_j t + g_j)^{-\lambda_j^{(r)}} \end{matrix} \right)$$

$${}_P F_Q \left[(A_P); (B_Q); - \sum_{i=1}^s z'_i \theta'_i (t-a)^{\mu'_i} (b-t)^{\rho'_i} \prod_{j=1}^k (f_j t + g_j)^{-\lambda'_j(i)} \right] dt = (b-a)^{\alpha+\beta-1} \frac{\prod_{j=1}^Q \Gamma(B_j)}{\prod_{j=1}^P \Gamma(A_j)} \prod_{j=1}^k (af_j + g_j)^{\sigma_j}$$

$$\sum_{h_1=1}^{M_1} \cdots \sum_{h_v=1}^{M_v} \sum_{k_1=0}^{\infty} \cdots \sum_{k_v=0}^{\infty} \sum_{R_1, \dots, R_u=0}^{h_1 R_1 + \dots + h_u R_u \leq L} \prod_{i=1}^v \frac{(-)^{k_i}}{\delta h_i^{(i)} k_i!} z_i^{m_i} \eta_{i, k_i} \prod_{k=1}^u z''_{R_k} B_u B_{u,v} [\phi_1 (\eta_{h_1, k_1}, \dots, \eta_{h_r, k_r})]_{j \neq h_i}$$

$$\mathfrak{N}_{U_{P+l+k+2, P+k+l+1}; W_1}^{0, n+P+k+k+2; V_1} \left(\begin{array}{c} \frac{z_1 (b-a)^{\mu_1 + \rho_1}}{\prod_{j=1}^k (af_j + g_j)^{\lambda_j^{(1)}}} \\ \dots \\ \frac{z_r (b-a)^{\mu_r + \rho_r}}{\prod_{j=1}^k (af_j + g_j)^{\lambda_j^{(r)}}} \\ \frac{z'_1 (b-a)^{\mu'_1 + \rho'_1}}{\prod_{j=1}^k (af_j + g_j)^{\lambda_j'^{(1)}}} \\ \dots \\ \frac{z'_s (b-a)^{\mu'_s + \rho'_s}}{\prod_{j=1}^k (af_j + g_j)^{\lambda_j'^{(s)}}} \\ \tau_1 (b-a)^{h_1} \\ \dots \\ \tau_l (b-a)^{h_l} \\ \frac{(b-a)f_1}{af_1 + g_1} \\ \dots \\ \frac{(b-a)f_k}{af_k + g_k} \end{array} \middle| \begin{array}{l} \mathbf{A} ; K_1, K_2, K_P, K_j, K'_j : C \\ \dots \\ \mathbf{B} ; L_1, L_Q, L_j, L'_j : D \end{array} \right) \quad (3.16)$$

where $U_{P+k+l+2, P+k+l+1} = p'_i + P + k + l + 2, q'_i + P + k + l + 1, \tau'_i; R'$

This result is an extension the formula given by Saxena et al [3].

Provided that

(A) $\max_{1 \leq j \leq k} \left\{ \left| \frac{(b-a)f_i}{af_i + g_i} \right| \right\} < 1, \max_{1 \leq j \leq l} \{ |\tau_j (b-a)^{h_j}| \} < 1,$

(B) $Re \left[\alpha + \sum_{i=1}^v a'_i \min_{1 \leq j \leq M_i} \frac{d_j^{(i)}}{\delta_j^{(i)}} + \sum_{i=1}^r \mu_i \min_{1 \leq j \leq m'_i} \frac{d_j'^{(i)}}{\delta_j'^{(i)}} \right] > 0$ and

$$Re \left[\beta + \sum_{i=1}^v b'_i \min_{1 \leq j \leq M_i} \frac{d_j^{(i)}}{\delta_j^{(i)}} + \sum_{i=1}^r \rho_i \min_{1 \leq j \leq m'_i} \frac{d'_j{}^{(i)}}{\delta'_j{}^{(i)}} \right] > 0$$

$$(C) Re \left(\alpha + \sum_{i=1}^v \eta_{G_i, g_i} a'_i + \sum_{i=1}^u R_i a_i + \sum_{i=1}^r \mu_i s_i + \sum_{i=1}^l h_i w_i \right) > 0; Re \left(\beta + \sum_{i=1}^v \eta_{G_i, g_i} b'_i + \sum_{i=1}^u R_i b_i + \sum_{i=1}^r \rho_i s_i \right) > 0$$

$$Re \left(\lambda_j + \sum_{i=1}^v \eta_{G_i, g_i} \lambda_j''^{(i)} + \sum_{i=1}^u R_i \lambda_j''^{(i)} + \sum_{i=1}^r s_i \zeta_j^{(i)} \right) > 0 (j = 1, \dots, l)$$

$$Re \left(-\sigma_j + \sum_{i=1}^v \eta_{G_i, g_i} \lambda_j'''^{(i)} + \sum_{i=1}^u R_i \lambda_j'''^{(i)} + \sum_{i=1}^r s_i \lambda_j^{(i)} \right) > 0 (j = 1, \dots, k)$$

$$(D) A_i'^{(k)} = \sum_{j=1}^{n'} \alpha_j'^{(k)} - \tau_i' \sum_{j=n'+1}^{p'_i} \alpha_{ji}^{(k)} - \tau_i' \sum_{j=1}^{q'_i} \beta_{ji}'^{(k)} + \sum_{j=1}^{n'_k} \gamma_j'^{(k)} - \tau_{i(k)}' \sum_{j=n'_k+1}^{p'_i(k)} \gamma_{ji}'^{(k)}$$

$$+ \sum_{j=1}^{m'_k} \delta_j'^{(k)} - \tau_{i(k)}' \sum_{j=m'_k+1}^{q'_i(k)} \delta_{ji}^{(k)} - \mu_i - \rho_i - \sum_{l=1}^k \lambda_l^{(i)} > 0 \quad (i = 1, \dots, r)$$

$$(E) \left| arg \left(z_i \prod_{j=1}^l [1 - \tau_j(t-a)^{h_i}]^{-\zeta_j^{(i)}} \prod_{j=1}^k (f_j t + g_j)^{-\lambda_j^{(i)}} \right) \right| < \frac{1}{2} A_i'^{(k)} \pi \quad (a \leq t \leq b; i = 1, \dots, r)$$

(F) $P \leq Q + 1$. The equality holds, when , in addition,

$$\text{either } P > Q \text{ and } \left| \left(z_i' \sum_{i=1}^l [1 - \tau_j(t-a)^{h_i}]^{-\zeta_j^{(i)}} \prod_{j=1}^k (f_j t + g_j)^{-\lambda_j^{(i)}} \right) \right|^{\frac{1}{Q-P}} < 1 \quad (a \leq t \leq b)$$

$$\text{or } P \leq Q \text{ and } \max_{1 \leq i \leq k} \left[\left| \left(z_i' \sum_{j=1}^l [1 - \tau_j(t-a)^{h_i}]^{-\zeta_j^{(i)}} \prod_{j=1}^k (f_j t + g_j)^{-\lambda_j^{(i)}} \right) \right| \right] < 1 \quad (a \leq t \leq b)$$

(G) The multiple series occurring on the right-hand side of (3.16) is absolutely and uniformly convergent.

Proof

To prove (3.16), first, we express in serie the multivariable \bar{I} -function with the help of (1.5), a class of multivariable polynomials defined by Srivastava et al [6] $S_L^{h_1, \dots, h_u} [.]$ in serie with the help of (1.20), the Aleph-functions of r-variables and in terms of Mellin-Barnes type contour integral with the help of (1.8), the generalized hypergeometric function ${}_pF_Q(\cdot)$ in Mellin-Barnes contour integral with the help of (2.1). Now collect the power of $[1 - \tau_j(t-a)^{h_i}]$ with $(i = 1, \dots, r; j = 1, \dots, l)$ and collect the power of $(f_j t + g_j)$ with $j = 1, \dots, k$. Use the equations (2.2) and (2.3) and express the result in Mellin-Barnes contour integral. Interpreting the $(r + s + k + l)$ dimensional Mellin-Barnes integral to multivariable Aleph-function, we obtain the equation (3.16).

4. Particular cases

a) If $\tau'_i, \tau'_{i(1)}, \dots, \tau'_{i(r)} \rightarrow 1$, the multivariable Aleph-function of s-variables reduces to multivariable I-function of s-variables defined by Sharma and al [4] and we have the following integral under the same conditions and notations that (3.16) with $\tau'_i, \tau'_{i(1)}, \dots, \tau'_{i(r)} \rightarrow 1$

$$\int_a^b (t-a)^{\alpha-1} (b-t)^{\beta-1} \prod_{j=1}^l [1-\tau_j(t-a)^{h_j}]^{-\lambda_j} \prod_{j=1}^k (f_j t + g_j)^{\sigma_j}$$

$$S_L^{h_1, \dots, h_u} \left(\begin{matrix} z_1'' \theta_1'' (t-a)^{a_1} (b-t)^{b_1} \prod_{j=1}^k (f_j t + g_j)^{-\lambda_j^{(1)}} \\ \vdots \\ z_u'' \theta_u'' (t-a)^{a_u} (b-t)^{b_u} \prod_{j=1}^k (f_j t + g_j)^{-\lambda_j^{(u)}} \end{matrix} \right)$$

$$\bar{I} \left(\begin{matrix} z_1''' \theta_1''' (t-a)^{a'_1} (b-t)^{b'_1} \prod_{j=1}^k (f_j t + g_j)^{-\lambda_j^{(1)'}} \\ \vdots \\ z_v''' \theta_v''' (t-a)^{a'_v} (b-t)^{b'_v} \prod_{j=1}^k (f_j t + g_j)^{-\lambda_j^{(v)'}} \end{matrix} \right)$$

$$I \left(\begin{matrix} z_1 \theta_1 (t-a)^{\mu_1} (b-t)^{\rho_1} \prod_{j=1}^k (f_j t + g_j)^{-\lambda_j^{(1)}} \\ \vdots \\ z_r \theta_r (t-a)^{\mu_r} (b-t)^{\rho_r} \prod_{j=1}^k (f_j t + g_j)^{-\lambda_j^{(r)}} \end{matrix} \right)$$

$${}_P F_Q \left[(A_P); (B_Q); - \sum_{i=1}^s z_i' \theta_i' (t-a)^{\mu_i'} (b-t)^{\rho_i'} \prod_{j=1}^k (f_j t + g_j)^{-\lambda_j^{(i)'}} \right] dt = (b-a)^{\alpha+\beta-1} \frac{\prod_{j=1}^Q \Gamma(B_j)}{\prod_{j=1}^P \Gamma(A_j)} \prod_{j=1}^k (a f_j + g_j)^{\sigma_j}$$

$$\sum_{h_1=1}^{M_1} \dots \sum_{h_v=1}^{M_v} \sum_{k_1=0}^{\infty} \dots \sum_{k_v=0}^{\infty} \sum_{R_1, \dots, R_u=0}^{h_1 R_1 + \dots + h_u R_u \leq L} \sum_{i=1}^v \frac{(-)^{k_i}}{\delta h_i^{(i)} k_i!} z_i''' \eta_{h_i, k_i} \prod_{k=1}^u z''^{R_k} B_u B_{u,v} [\phi_1 (\eta_{h_1, k_1}, \dots, \eta_{h_r, k_r})]_{j \neq h_i}$$

$$I_{U_{P+l+k+2, Q+k+l+1}:W_1}^{0, n+P+k+k+2:V_1} \left(\begin{matrix} \frac{z_1(b-a)^{\mu_1+\rho_1}}{\prod_{j=1}^k (af_j+g_j)^{\lambda_j^{(1)}}} \\ \dots \\ \frac{z_r(b-a)^{\mu_r+\rho_r}}{\prod_{j=1}^k (af_j+g_j)^{\lambda_j^{(r)}}} \\ \frac{z'_1(b-a)^{\mu'_1+\rho'_1}}{\prod_{j=1}^k (af_j+g_j)^{\lambda'_j^{(1)}}} \\ \dots \\ \frac{z'_s(b-a)^{\mu'_s+\rho'_s}}{\prod_{j=1}^k (af_j+g_j)^{\lambda'_j^{(s)}}} \\ \tau_1(b-a)^{h_1} \\ \dots \\ \tau_l(b-a)^{h_l} \\ \frac{(b-a)f_1}{af_1+g_1} \\ \dots \\ \frac{(b-a)f_k}{af_k+g_k} \end{matrix} \right) \quad \left. \begin{matrix} A; K_1, K_2, K_P, K_j, K'_j : C \\ \\ \\ \\ \\ \\ \\ \\ \\ B; L_1, L_Q, L_j, L'_j : D \end{matrix} \right) \tag{4.1}$$

$$\text{b) If } B(L; R_1, \dots, R_r) = \frac{\prod_{j=1}^{\bar{A}} (a_j)_{R_1\theta'_j+\dots+R_r\theta_j^{(r)}} \prod_{j=1}^{B'} (b'_j)_{R_1\phi'_j} \dots \prod_{j=1}^{B^{(r)}} (b_j^{(r)})_{R_r\phi_j^{(r)}}}{\prod_{j=1}^{\bar{C}} (c_j)_{m_1\psi'_j+\dots+m_r\psi_j^{(r)}} \prod_{j=1}^{D'} (d'_j)_{R_1\delta'_j} \dots \prod_{j=1}^{D^{(r)}} (d_j^{(r)})_{R_r\delta_j^{(r)}}} \tag{4.2}$$

then the general class of multivariable polynomial $S_L^{h_1, \dots, h_u} [z_1, \dots, z_u]$ reduces to generalized Lauricella function defined by Srivastava et al [5]. We have the following integral

$$\int_a^b (t-a)^{\alpha-1} (b-t)^{\beta-1} \prod_{j=1}^l [1-\tau_j(t-a)^{h_j}]^{-\lambda_j} \prod_{j=1}^k (f_j t + g_j)^{\sigma_j}$$

$$F_{\bar{C}:D'; \dots; D^{(u)}}^{1+\bar{A}:B'; \dots; B^{(u)}} \left(\begin{matrix} z''_1 \theta''_1 (t-a)^{a_1} (b-t)^{b_1} \prod_{j=1}^k (f_j t + g_j)^{-\lambda''_j^{(1)}} \\ \vdots \\ z''_u \theta''_u (t-a)^{a_u} (b-t)^{b_u} \prod_{j=1}^k (f_j t + g_j)^{-\lambda''_j^{(u)}} \end{matrix} \right)$$

$$\left([(-L); R_1, \dots, R_u] [(a); \theta', \dots, \theta^{(u)}] : [(b'); \phi']; \dots; [(b^{(u)}); \phi^{(u)}] \right)$$

$$\left[[(c); \psi', \dots, \psi^{(u)}] : [(d'); \delta']; \dots; [(d^{(u)}); \delta^{(u)}] \right]$$

$$\bar{I} \begin{pmatrix} z_1''' \theta_1''' (t-a)^{a_1'} (b-t)^{b_1'} \prod_{j=1}^k (f_j t + g_j)^{-\lambda_j^{(1)}} \\ \vdots \\ z_v''' \theta_v''' (t-a)^{a_v'} (b-t)^{b_v'} \prod_{j=1}^k (f_j t + g_j)^{-\lambda_j^{(v)}} \end{pmatrix} \Re \begin{pmatrix} z_1 \theta_1 (t-a)^{\mu_1} (b-t)^{\rho_1} \prod_{j=1}^k (f_j t + g_j)^{-\lambda_j^{(1)}} \\ \vdots \\ z_r \theta_r (t-a)^{\mu_r} (b-t)^{\rho_r} \prod_{j=1}^k (f_j t + g_j)^{-\lambda_j^{(r)}} \end{pmatrix}$$

$${}^P F_Q \left[(A_P); (B_Q); - \sum_{i=1}^s z_i' \theta_i' (t-a)^{\mu_i'} (b-t)^{\rho_i'} \prod_{j=1}^k (f_j t + g_j)^{-\lambda_j^{(i)}} \right] dt = (b-a)^{\alpha+\beta-1} \frac{\prod_{j=1}^Q \Gamma(B_j)}{\prod_{j=1}^P \Gamma(A_j)} \prod_{j=1}^Q (af_j + g_j)^{\sigma_j}$$

$$\sum_{h_1=1}^{M_1} \dots \sum_{h_v=1}^{M_v} \sum_{k_1=0}^{\infty} \dots \sum_{k_v=0}^{\infty} \sum_{R_1, \dots, R_u=0}^{h_1 R_1 + \dots + h_u R_u \leq L} \prod_{i=1}^v \frac{(-)^{k_i}}{\delta h_i^{(i)} k_i!} z_i''' \eta_{h_i, k_i} \prod_{k=1}^u z_i'' R_k B_u B_{u,v} [\phi_1 (\eta_{h_1, k_1}, \dots, \eta_{h_r, k_r})]_{j \neq h_i}$$

$$\Re_{U_{P+l+k+2, Q+k+l+1}; V_1}^{0, n+P+k+k+2; W_1} \left(\begin{array}{c|c} \frac{z_1 (b-a)^{\mu_1 + \rho_1}}{\prod_{j=1}^k (af_j + g_j)^{\lambda_j^{(1)}}} & \vdots \\ \dots & \vdots \\ \frac{z_r (b-a)^{\mu_r + \rho_r}}{\prod_{j=1}^k (af_j + g_j)^{\lambda_j^{(r)}}} & \vdots \\ \frac{z_1' (b-a)^{\mu_1' + \rho_1'}}{\prod_{j=1}^k (af_j + g_j)^{\lambda_j'^{(1)}}} & \vdots \\ \dots & \vdots \\ \frac{z_s' (b-a)^{\mu_s' + \rho_s'}}{\prod_{j=1}^k (af_j + g_j)^{\lambda_j'^{(s)}}} & \vdots \\ \tau_1 (b-a)^{h_1} & \vdots \\ \dots & \vdots \\ \tau_l (b-a)^{h_l} & \vdots \\ \frac{(b-a)f_1}{af_1 + g_1} & \vdots \\ \dots & \vdots \\ \frac{(b-a)f_k}{af_k + g_k} & \vdots \end{array} \right) \begin{array}{l} A ; K_1, K_2, K_P, K_j, K_j', \mathfrak{A} : A' \\ \\ \\ \\ B ; L_1, L_Q, L_j, L_j', \mathfrak{B} : B' \end{array} \tag{4.3}$$

under the same notations and conditions that (3.16)

where $B'_u = \frac{(-L)^{h_1 R_1 + \dots + h_u R_u} B(E; R_1, \dots, R_u)}{R_1! \dots R_u!}$, $B[E; R_1, \dots, R_v]$ is defined by (4.2)

Remark:

By the following similar procedure, the results of this document can be extended to product of any finite number of multivariable Aleph-functions and a class of multivariable polynomials defined by Srivastava et al [6].

5. Conclusion

In this paper we have evaluated a generalized Eulerian integral involving the product of the multivariable Aleph-function, the multivariable \bar{I} -function defined by Nambisan et al [1], a class of multivariable polynomials and generalized hypergeometric function with general arguments. The formulae established in this paper is very general nature. Thus, the results established in this research work would serve as a key formula from which, upon specializing the parameters, as many as desired results involving the special functions of one and several variables can be obtained.

REFERENCES

- [1] Prathima J. Nambisan V. and Kurumujji S.K. A Study of I-function of Several Complex Variables, International Journal of Engineering Mathematics Vol(2014) , 2014 page 1-12
- [2] Saigo M. and Saxena R.K. Unified fractional integral formulas for the multivariable H-function I. J.Fractional Calculus 15 (1999), page 91-107.
- [3] Saigo M. and Saxena R.K. Unified fractional integral formulas for the multivariable H-function III. J.Fractional Calculus 20 (2001), page 45-68.
- [4] Sharma C.K.and Ahmad S.S.: On the multivariable I-function. Acta ciencia Indica Math , 1994 vol 20,no2, p 113-116.
- [5] Srivastava H.M. and Daoust M.C. Certain generalized Neumann expansions associated with Kampé de Fériet function. Nederl. Akad. Wetensch. Proc. Ser A72 = Indag Math 31(1969) page 449-457.
- [6] Srivastava H.M. And Garg M. Some integral involving a general class of polynomials and multivariable H-function. Rev. Roumaine Phys. 32(1987), page 685-692.
- [7] Srivastava H.M. and Karlsson P.W. Multiple Gaussian Hypergeometric series. Ellis.Horwood. Limited. New-York, Chichester. Brisbane. Toronto , 1985.
- [8] H.M. Srivastava And R.Panda. Some expansion theorems and generating relations for the H-function of several complex variables. Comment. Math. Univ. St. Paul. 24(1975), p.119-137.

Personal adress : 411 Avenue Joseph Raynaud
Le parc Fleuri , Bat B
83140 , Six-Fours les plages
Tel : 06-83-12-49-68
Department : VAR
Country : FRANCE