

Eulerian integral associated with product of two multivariable Aleph-functions and classes of polynomials

F.Y. AYANT¹

¹ Teacher in High School , France

ABSTRACT

The present paper is evaluated a new Eulerian integral associated with the product of two multivariable Aleph-functions, a generalized Lauricella function , a classes of multivariable polynomials with general arguments . We will study the case concerning the multivariable H-function defined by Srivastava et al [7] and Srivastava-Daoust polynomial [4].

Keywords: Eulerian integral, multivariable Aleph-function, generalized Lauricella function of several variables, multivariable H-function, generalized hypergeometric function, class of polynomials

2010 Mathematics Subject Classification. 33C99, 33C60, 44A20

1. Introduction

In this paper, we evaluate a new Eulerian integral of most general characters associated with the products of two multivariable Aleph-functions and classes of polynomials with general arguments.

The Aleph-function of several variables generalize the multivariable I-function defined by Sharma and Ahmad [2] , itself is an a generalisation of G and H-functions of several variables defined by Srivastava et al [7]. The multiple Mellin-Barnes integral occurring in this paper will be referred to as the multivariables Aleph-function throughout our present study and will be defined and represented as follows.

$$\begin{aligned}
 \text{We have : } \aleph(z_1, \dots, z_r) &= \aleph_{p_i, q_i, \tau_i; R: p_i(1), q_i(1), \tau_i(1); R^{(1)}; \dots; p_i(r), q_i(r), \tau_i(r); R^{(r)}}^{0, n: m_1, n_1, \dots, m_r, n_r} \left(\begin{matrix} z_1 \\ \cdot \\ \cdot \\ \cdot \\ z_r \end{matrix} \right) \\
 & [(a_j; \alpha_j^{(1)}, \dots, \alpha_j^{(r)})_{1, n}] , [\tau_i(a_{ji}; \alpha_{ji}^{(1)}, \dots, \alpha_{ji}^{(r)})_{n+1, p_i}] : \\
 & \dots \dots \dots [\tau_i(b_{ji}; \beta_{ji}^{(1)}, \dots, \beta_{ji}^{(r)})_{m+1, q_i}] : \\
 & [(c_j^{(1)}, \gamma_j^{(1)})_{1, n_1}], [\tau_{i(1)}(c_{ji(1)}, \gamma_{ji(1)})_{n_1+1, p_i^{(1)}}]; \dots ; [(c_j^{(r)}, \gamma_j^{(r)})_{1, n_r}], [\tau_{i(r)}(c_{ji(r)}, \gamma_{ji(r)})_{n_r+1, p_i^{(r)}}] \\
 & [(d_j^{(1)}, \delta_j^{(1)})_{1, m_1}], [\tau_{i(1)}(d_{ji(1)}, \delta_{ji(1)})_{m_1+1, q_i^{(1)}}]; \dots ; [(d_j^{(r)}, \delta_j^{(r)})_{1, m_r}], [\tau_{i(r)}(d_{ji(r)}, \delta_{ji(r)})_{m_r+1, q_i^{(r)}}] \\
 & = \frac{1}{(2\pi\omega)^r} \int_{L'_1} \dots \int_{L'_r} \psi(s_1, \dots, s_r) \prod_{k=1}^r \theta_k(s_k) z_k^{s_k} ds_1 \dots ds_r \tag{1.8}
 \end{aligned}$$

with $\omega = \sqrt{-1}$

$$\psi(s_1, \dots, s_r) = \frac{\prod_{j=1}^n \Gamma(1 - a_j + \sum_{k=1}^r \alpha_j^{(k)} s_k)}{\sum_{i=1}^R [\tau_i \prod_{j=n+1}^{p_i} \Gamma(a_{ji} - \sum_{k=1}^r \alpha_{ji}^{(k)} s_k) \prod_{j=1}^{q_i} \Gamma(1 - b_{ji} + \sum_{k=1}^r \beta_{ji}^{(k)} s_k)]} \tag{1.9}$$

$$\text{and } \theta_k(s_k) = \frac{\prod_{j=1}^{m_k} \Gamma(d_j^{(k)} - \delta_j^{(k)} s_k) \prod_{j=1}^{n_k} \Gamma(1 - c_j^{(k)} + \gamma_j^{(k)} s_k)}{\sum_{i^{(k)}=1}^{R^{(k)}} [\tau_{i^{(k)}} \prod_{j=m_k+1}^{q_{i^{(k)}}} \Gamma(1 - d_{ji^{(k)}}^{(k)} + \delta_{ji^{(k)}}^{(k)} s_k) \prod_{j=n_k+1}^{p_{i^{(k)}}} \Gamma(c_{ji^{(k)}}^{(k)} - \gamma_{ji^{(k)}}^{(k)} s_k)]} \quad (1.10)$$

Suppose, as usual, that the parameters

$$a_j, j = 1, \dots, p; b_j, j = 1, \dots, q;$$

$$c_j^{(k)}, j = 1, \dots, n_k; c_{ji^{(k)}}^{(k)}, j = n_k + 1, \dots, p_{i^{(k)}};$$

$$d_j^{(k)}, j = 1, \dots, m_k; d_{ji^{(k)}}^{(k)}, j = m_k + 1, \dots, q_{i^{(k)}};$$

$$\text{with } k = 1 \dots, r, i = 1, \dots, R, i^{(k)} = 1, \dots, R^{(k)}$$

are complex numbers, and the $\alpha' s, \beta' s, \gamma' s$ and $\delta' s$ are assumed to be positive real numbers for standardization purpose such that

$$U_i^{(k)} = \sum_{j=1}^n \alpha_j^{(k)} + \tau_i \sum_{j=n+1}^{p_i} \alpha_{ji}^{(k)} + \sum_{j=1}^{n_k} \gamma_j^{(k)} + \tau_{i^{(k)}} \sum_{j=n_k+1}^{p_{i^{(k)}}} \gamma_{ji^{(k)}}^{(k)} - \tau_i \sum_{j=1}^{q_i} \beta_{ji}^{(k)} - \sum_{j=1}^{m_k} \delta_j^{(k)} - \tau_{i^{(k)}} \sum_{j=m_k+1}^{q_{i^{(k)}}} \delta_{ji^{(k)}}^{(k)} \leq 0 \quad (1.11)$$

The real numbers τ_i are positives for $i = 1$ to R , $\tau_{i^{(k)}}$ are positives for $i^{(k)} = 1$ to $R^{(k)}$

The contour L_k is in the s_k -p lane and run from $\sigma - i\infty$ to $\sigma + i\infty$ where σ is a real number with loop, if necessary, ensure that the poles of $\Gamma(d_j^{(k)} - \delta_j^{(k)} s_k)$ with $j = 1$ to m_k are separated from those of $\Gamma(1 - a_j + \sum_{i=1}^r \alpha_j^{(k)} s_k)$ with $j = 1$ to n and $\Gamma(1 - c_j^{(k)} + \gamma_j^{(k)} s_k)$ with $j = 1$ to n_k to the left of the contour L_k . The condition for absolute convergence of multiple Mellin-Barnes type contour (1.9) can be obtained by extension of the corresponding conditions for multivariable H-function given by as :

$$|\arg z_k| < \frac{1}{2} A_i^{(k)} \pi, \text{ where}$$

$$A_i^{(k)} = \sum_{j=1}^n \alpha_j^{(k)} - \tau_i \sum_{j=n+1}^{p_i} \alpha_{ji}^{(k)} - \tau_i \sum_{j=1}^{q_i} \beta_{ji}^{(k)} + \sum_{j=1}^{n_k} \gamma_j^{(k)} - \tau_{i^{(k)}} \sum_{j=n_k+1}^{p_{i^{(k)}}} \gamma_{ji^{(k)}}^{(k)} + \sum_{j=1}^{m_k} \delta_j^{(k)} - \tau_{i^{(k)}} \sum_{j=m_k+1}^{q_{i^{(k)}}} \delta_{ji^{(k)}}^{(k)} > 0, \text{ with } k = 1 \dots, r, i = 1, \dots, R, i^{(k)} = 1, \dots, R^{(k)} \quad (1.12)$$

The complex numbers z_i are not zero. Throughout this document, we assume the existence and absolute convergence conditions of the multivariable Aleph-function.

We may establish the asymptotic expansion in the following convenient form :

$$\aleph(z_1, \dots, z_r) = O(|z_1|^{\alpha_1}, \dots, |z_r|^{\alpha_r}), \max(|z_1|, \dots, |z_r|) \rightarrow 0$$

$$\aleph(z_1, \dots, z_r) = O(|z_1|^{\beta_1}, \dots, |z_r|^{\beta_r}), \min(|z_1|, \dots, |z_r|) \rightarrow \infty$$

where $k = 1, \dots, r : \alpha_k = \min[\operatorname{Re}(d_j^{(k)} / \delta_j^{(k)})], j = 1, \dots, m_k$ and

$$\beta_k = \max[\operatorname{Re}((c_j^{(k)} - 1) / \gamma_j^{(k)})], j = 1, \dots, n_k$$

We will use these following notations in this paper

$$U = p_i, q_i, \tau_i; R; V = m_1, n_1; \dots; m_r, n_r \tag{1.13}$$

$$W = p_{i(1)}, q_{i(1)}, \tau_{i(1)}; R^{(1)}, \dots, p_{i(r)}, q_{i(r)}, \tau_{i(r)}; R^{(r)} \tag{1.14}$$

$$A = \{(a_j; \alpha_j^{(1)}, \dots, \alpha_j^{(r)})_{1,n}\}, \{\tau_i(a_{ji}; \alpha_{ji}^{(1)}, \dots, \alpha_{ji}^{(r)})_{n+1,p_i}\} \tag{1.15}$$

$$B = \{\tau_i(b_{ji}; \beta_{ji}^{(1)}, \dots, \beta_{ji}^{(r)})_{m+1,q_i}\} \tag{1.16}$$

$$C = \{(c_j^{(1)}; \gamma_j^{(1)})_{1,n_1}\}, \tau_{i(1)}(c_{ji(1)}; \gamma_{ji(1)})_{n_1+1,p_{i(1)}}, \dots, \{(c_j^{(r)}; \gamma_j^{(r)})_{1,n_r}\}, \tau_{i(r)}(c_{ji(r)}; \gamma_{ji(r)})_{n_r+1,p_{i(r)}} \tag{1.17}$$

$$D = \{(d_j^{(1)}; \delta_j^{(1)})_{1,m_1}\}, \tau_{i(1)}(d_{ji(1)}; \delta_{ji(1)})_{m_1+1,q_{i(1)}}, \dots, \{(d_j^{(r)}; \delta_j^{(r)})_{1,m_r}\}, \tau_{i(r)}(d_{ji(r)}; \delta_{ji(r)})_{m_r+1,q_{i(r)}} \tag{1.18}$$

The multivariable Aleph-function write :

$$\aleph(z_1, \dots, z_r) = \aleph_{U:W}^{0,n:V} \left(\begin{array}{c|c} z_1 & \text{A : C} \\ \cdot & \cdot \\ \cdot & \cdot \\ \cdot & \cdot \\ z_r & \text{B : D} \end{array} \right) \tag{1.19}$$

Consider the Aleph-function of s variables

$$\aleph(z_1, \dots, z_s) = \aleph_{p'_i, q'_i, \tau'_i; r^{(1)}; \dots; p'_i(s), q'_i(s), \tau'_i(s); r^{(s)}}^{0, n': m'_1, n'_1, \dots, m'_s, n'_s} \left(\begin{array}{c|c} z_1 & \\ \cdot & \\ \cdot & \\ \cdot & \\ z_s & \end{array} \right)$$

$$\begin{aligned} & [(u_j; \mu_j^{(1)}, \dots, \mu_j^{(r')})_{1, n'}] : [\tau_i(u_{ji}; \mu_{ji}^{(1)}, \dots, \mu_{ji}^{(r')})_{n'+1, p'_i}] : \\ & \dots \dots \dots [\tau_i(v_{ji}; \nu_{ji}^{(1)}, \dots, \nu_{ji}^{(r')})_{m'+1, q'_i}] : \end{aligned}$$

$$\left[(a_j^{(1)}; \alpha_j^{(1)})_{1, n'_1}, [\tau_{i(1)}(a_{ji(1)}; \alpha_{ji(1)})_{n'_1+1, p'_{i(1)}}]; \dots; [(a_j^{(s)}; \alpha_j^{(s)})_{1, n'_s}], [\tau_{i(s)}(a_{ji(s)}; \alpha_{ji(s)})_{n'_s+1, p'_{i(s)}}] \right] \\ \left[(b_j^{(1)}; \beta_j^{(1)})_{1, m'_1}, [\tau_{i(1)}(b_{ji(1)}; \beta_{ji(1)})_{m'_1+1, q'_{i(1)}}]; \dots; [(b_j^{(s)}; \beta_j^{(s)})_{1, m'_s}], [\tau_{i(s)}(b_{ji(s)}; \beta_{ji(s)})_{m'_s+1, q'_{i(s)}}] \right]$$

$$= \frac{1}{(2\pi\omega)^s} \int_{L''_1} \dots \int_{L''_s} \zeta(t_1, \dots, t_s) \prod_{k=1}^s \phi_k(t_k) z_k^{t_k} dt_1 \dots dt_s \tag{1.20}$$

with $\omega = \sqrt{-1}$

$$\zeta(t_1, \dots, t_s) = \frac{\prod_{j=1}^{n'} \Gamma(1 - u_j + \sum_{k=1}^s \mu_j^{(k)} t_k)}{\sum_{i=1}^{r'} [\tau_i \prod_{j=n'+1}^{p_i} \Gamma(u_{ji} - \sum_{k=1}^s \mu_{ji}^{(k)} t_k) \prod_{j=1}^{q'_i} \Gamma(1 - v_{ji} + \sum_{k=1}^s \nu_{ji}^{(k)} t_k)]} \tag{1.21}$$

$$\text{and } \phi_k(t_k) = \frac{\prod_{j=1}^{m'_k} \Gamma(b_j^{(k)} - \beta_j^{(k)} t_k) \prod_{j=1}^{n'_k} \Gamma(1 - a_j^{(k)} + \alpha_j^{(k)} s_k)}{\sum_{i^{(k)}=1}^{r^{(k)}} [\iota_{i^{(k)}} \prod_{j=m'_k+1}^{Q_{i^{(k)}}} \Gamma(1 - b_{ji^{(k)}}^{(k)} + \beta_{ji^{(k)}}^{(k)} t_k) \prod_{j=n'_k+1}^{P_{i^{(k)}}} \Gamma(a_{ji^{(k)}}^{(k)} - \alpha_{ji^{(k)}}^{(k)} s_k)]} \quad (1.22)$$

Suppose , as usual , that the parameters

$$u_j, j = 1, \dots, p'; v_j, j = 1, \dots, q';$$

$$a_j^{(k)}, j = 1, \dots, n'_k; a_{ji^{(k)}}^{(k)}, j = n_k + 1, \dots, p'_{i^{(k)}};$$

$$b_{ji^{(k)}}^{(k)}, j = m'_k + 1, \dots, q'_{i^{(k)}}; b_j^{(k)}, j = 1, \dots, m'_k;$$

$$\text{with } k = 1 \dots, s, i = 1, \dots, r', i^{(k)} = 1, \dots, r^{(k)}$$

are complex numbers , and the $\alpha' s, \beta' s, \gamma' s$ and $\delta' s$ are assumed to be positive real numbers for standardization purpose such that

$$U_i^{(k)} = \sum_{j=1}^{n'} \mu_j^{(k)} + \iota_i \sum_{j=n'+1}^{p'_i} \mu_{ji}^{(k)} + \sum_{j=1}^{n'_k} \alpha_j^{(k)} + \iota_{i^{(k)}} \sum_{j=n'_k+1}^{p'_{i^{(k)}}} \alpha_{ji^{(k)}}^{(k)} - \iota_i \sum_{j=1}^{q'_i} v_{ji}^{(k)} - \sum_{j=1}^{m'_k} \beta_j^{(k)} - \iota_{i^{(k)}} \sum_{j=m'_k+1}^{q'_{i^{(k)}}} \beta_{ji^{(k)}}^{(k)} \leq 0 \quad (1.23)$$

The reals numbers τ_i are positives for $i = 1, \dots, s$, $\iota_{i^{(k)}}$ are positives for $i^{(k)} = 1 \dots r^{(k)}$

The contour L_k is in the t_k -p lane and run from $\sigma - i\infty$ to $\sigma + i\infty$ where σ is a real number with loop , if necessary , ensure that the poles of $\Gamma(b_j^{(k)} - \beta_j^{(k)} t_k)$ with $j = 1$ to m'_k are separated from those of $\Gamma(1 - u_j + \sum_{i=1}^s \mu_j^{(k)} t_k)$ with $j = 1$ to N and $\Gamma(1 - a_j^{(k)} + \alpha_j^{(k)} t_k)$ with $j = 1$ to n'_k to the left of the contour L_k . The condition for absolute convergence of multiple Mellin-Barnes type contour (1.9) can be obtained by extension of the corresponding conditions for multivariable H-function given by as :

$$|arg z_k| < \frac{1}{2} B_i^{(k)} \pi, \text{ where}$$

$$B_i^{(k)} = \sum_{j=1}^{n'} \mu_j^{(k)} - \iota_i \sum_{j=n'+1}^{p'_i} \mu_{ji}^{(k)} - \iota_i \sum_{j=1}^{q'_i} v_{ji}^{(k)} + \sum_{j=1}^{n'_k} \alpha_j^{(k)} - \iota_{i^{(k)}} \sum_{j=n'_k+1}^{p'_{i^{(k)}}} \alpha_{ji^{(k)}}^{(k)} + \sum_{j=1}^{m'_k} \beta_j^{(k)} - \iota_{i^{(k)}} \sum_{j=m'_k+1}^{q'_{i^{(k)}}} \beta_{ji^{(k)}}^{(k)} > 0, \text{ with } k = 1, \dots, s, i = 1, \dots, r, i^{(k)} = 1, \dots, r^{(k)} \quad (1.24)$$

The complex numbers z_i are not zero. Throughout this document , we assume the existence and absolute convergence conditions of the multivariable Aleph-function.

We may establish the the asymptotic expansion in the following convenient form :

$$\aleph(z_1, \dots, z_s) = O(|z_1|^{\alpha'_1}, \dots, |z_s|^{\alpha'_s}), \max(|z_1|, \dots, |z_s|) \rightarrow 0$$

$$\aleph(z_1, \dots, z_s) = O(|z_1|^{\beta'_1}, \dots, |z_s|^{\beta'_s}), \min(|z_1|, \dots, |z_s|) \rightarrow \infty$$

where $k = 1, \dots, s, z : \alpha'_k = \min[Re(b_j^{(k)})/\beta_j^{(k)}], j = 1, \dots, m'_k$ and

$$\beta'_k = \max[\operatorname{Re}((a_j^{(k)} - 1)/\alpha_j^{(k)}), j = 1, \dots, n'_k]$$

We will use these following notations in this paper

$$U' = p'_i, q'_i, \iota_i; r'; V' = m'_1, n'_1; \dots; m'_s, n'_s \tag{1.25}$$

$$W' = p'_{i(1)}, q'_{i(1)}, \iota_{i(1)}; r^{(1)}, \dots, p'_{i(r)}, q'_{i(r)}, \iota_{i(s)}; r^{(s)} \tag{1.26}$$

$$A' = \{(u_j; \mu_j^{(1)}, \dots, \mu_j^{(s)})_{1, n'}\}, \{\iota_i(u_{ji}; \mu_{ji}^{(1)}, \dots, \mu_{ji}^{(s)})_{n'+1, p'_i}\} \tag{1.27}$$

$$B' = \{\iota_i(v_{ji}; v_{ji}^{(1)}, \dots, v_{ji}^{(s)})_{m'+1, q'_i}\} \tag{1.28}$$

$$C' = (a_j^{(1)}; \alpha_j^{(1)})_{1, n'_1}, \iota_{i(1)}(a_{ji(1)}^{(1)}; \alpha_{ji(1)}^{(1)})_{n'_1+1, p'_{i(1)}}, \dots, (a_j^{(s)}; \alpha_j^{(s)})_{1, n'_s}, \iota_{i(s)}(a_{ji(s)}^{(s)}; \alpha_{ji(s)}^{(s)})_{n'_s+1, p'_{i(s)}} \tag{1.29}$$

$$D' = (b_j^{(1)}; \beta_j^{(1)})_{1, m'_1}, \iota_{i(1)}(b_{ji(1)}^{(1)}; \beta_{ji(1)}^{(1)})_{m'_1+1, q'_{i(1)}}, \dots, (b_j^{(s)}; \beta_j^{(s)})_{1, m'_s}, \iota_{i(s)}(\beta_{ji(s)}^{(s)}; \beta_{ji(s)}^{(s)})_{m'_s+1, q'_{i(s)}} \tag{1.30}$$

The multivariable Aleph-function write :

$$\aleph(z_1, \dots, z_s) = \aleph_{U'; W'}^{0, n'; V'} \left(\begin{array}{c|c} z_1 & A' : C' \\ \cdot & \cdot \\ \cdot & \cdot \\ z_s & B' : D' \end{array} \right) \tag{1.31}$$

The generalized polynomials of multivariables defined by Srivastava [3], is given in the following manner :

$$S_{N_1, \dots, N_v}^{\mathfrak{M}_1, \dots, \mathfrak{M}_v} [y_1, \dots, y_v] = \sum_{K_1=0}^{[N_1/\mathfrak{M}_1]} \dots \sum_{K_v=0}^{[N_v/\mathfrak{M}_v]} \frac{(-N_1)_{\mathfrak{M}_1 K_1}}{K_1!} \dots \frac{(-N_v)_{\mathfrak{M}_v K_v}}{K_v!} A[N_1, K_1; \dots; N_v, K_v] y_1^{K_1} \dots y_v^{K_v} \tag{1.32}$$

where $\mathfrak{M}_1, \dots, \mathfrak{M}_v$ are arbitrary positive integers and the coefficients $A[N_1, K_1; \dots; N_v, K_v]$ are arbitrary constants, real or complex.

Srivastava and Garg [5] introduced and defined a general class of multivariable polynomials as follows

$$S_L^{h_1, \dots, h_u} [z_1, \dots, z_u] = \sum_{R_1, \dots, R_u=0}^{h_1 R_1 + \dots + h_u R_u \leq L} (-L)_{h_1 R_1 + \dots + h_u R_u} B(E; R_1, \dots, R_u) \frac{z_1^{R_1} \dots z_u^{R_u}}{R_1! \dots R_u!} \tag{1.33}$$

The coefficients are $B[E; R_1, \dots, R_u]$ arbitrary constants, real or complex.

We will note $a_v = \frac{(-N_1)_{\mathfrak{M}_1 K_1}}{K_1!} \dots \frac{(-N_v)_{\mathfrak{M}_v K_v}}{K_v!} A[N_1, K_1; \dots; N_v, K_v]$ and

$$b_u = \frac{(-E)_{F_1 L_1 + \dots + F_u L_u} B(E; L_1, \dots, L_u)}{L_1! \dots L_u!} \tag{1.9}$$

2. Integral representation of generalized Lauricella function of several variables

The following generalized hypergeometric function in terms of multiple contour integrals is also required [6 ,page 39

eq .30]

$$\frac{\prod_{j=1}^P \Gamma(A_j)}{\prod_{j=1}^Q \Gamma(B_j)} {}_P F_Q [(A_P); (B_Q); -(x_1 + \dots + x_r)]$$

$$= \frac{1}{(2\pi\omega)^r} \int_{L_1} \dots \int_{L_r} \frac{\prod_{j=1}^P \Gamma(A_j + s_1 + \dots + s_r)}{\prod_{j=1}^Q \Gamma(B_j + s_1 + \dots + s_r)} \Gamma(-s_1) \dots \Gamma(-s_r) x_1^{s_1} \dots x_r^{s_r} ds_1 \dots ds_r \tag{2.1}$$

where the contours are of Barnes type with indentations, if necessary, to ensure that the poles of $\Gamma(A_j + s_1 + \dots + s_r)$ are separated from those of $\Gamma(-s_j), j = 1, \dots, r$. The above result (1.23) can be easily established by an appeal to the calculus of residues by calculating the residues at the poles of $\Gamma(-s_j), j = 1, \dots, r$

In order to evaluate a number of integrals of multivariable I-function, we first establish the formula

$$\int_a^b (t-a)^{\alpha-1} (b-t)^{\beta-1} \prod_{j=1}^l [1 - \tau_j (t-a)^{h_j}]^{-\lambda_j} \prod_{j=1}^k (f_j t + g_j)^{\sigma_j} dt = (b-a)^{\alpha+\beta-1} B(\alpha, \beta) \prod_{j=1}^k (af_j + g_j)^{\sigma_j}$$

$$F_{1:0, \dots, 0; 0, \dots, 0}^{1:1, \dots, 1; 1, \dots, 1} \left(\begin{matrix} (\alpha : h_1, \dots, h_l, 1, \dots, 1) : (\lambda_1 : 1), \dots, (\lambda_l : 1); (-\sigma_1 : 1), \dots, (-\sigma_k : 1) \\ \dots \\ (\alpha + \beta : h_1, \dots, h_l, 1, \dots, 1) : -, \dots, -; -, \dots, - \end{matrix} \right)$$

$$; \tau_1 (b-a)^{h_1}, \dots, \tau_l (b-a)^{h_l}, -\frac{(b-a)f_1}{af_1 + g_1}, \dots, -\frac{(b-a)f_k}{af_k + g_k} \tag{2.2}$$

where $a, b \in \mathbb{R} (a < b), \alpha, \beta, f_i, g_i, \sigma_i, \tau_j, h_j \in \mathbb{C}, \lambda_j \in \mathbb{R}^+ (i = 1, \dots, k; j = 1, \dots, l)$

$$\min(Re(\alpha), Re(\beta)) > 0, \max_{1 \leq j \leq l} \{ |\tau_j (b-a)^{h_j}| \} < 1, \max_{1 \leq j \leq k} \left\{ \left| \frac{(b-a)f_j}{af_j + g_j} \right| \right\} < 1,$$

and $F_{1:0, \dots, 0; 0, \dots, 0}^{1:1, \dots, 1; 1, \dots, 1}$ is a particular case of the generalized Lauricella function introduced by Srivastava-Daoust[3,page 454] given by :

$$F_{1:0, \dots, 0; 0, \dots, 0}^{1:1, \dots, 1; 1, \dots, 1} \left(\begin{matrix} (\alpha : h_1, \dots, h_l, 1, \dots, 1) : (\lambda_1 : 1), \dots, (\lambda_l : 1); (-\sigma_1 : 1), \dots, (-\sigma_k : 1) \\ \dots \\ (\alpha + \beta : h_1, \dots, h_l, 1, \dots, 1) : -, \dots, -; -, \dots, - \end{matrix} \right)$$

$$; \tau_1 (b-a)^{h_1}, \dots, \tau_l (b-a)^{h_l}, -\frac{(b-a)f_1}{af_1 + g_1}, \dots, -\frac{(b-a)f_k}{af_k + g_k} \Bigg) = \frac{\Gamma(\alpha + \beta)}{\Gamma(\alpha) \prod_{j=1}^l \Gamma(\lambda_j) \prod_{j=1}^k \Gamma(-\sigma_j)}$$

$$\frac{1}{(2\pi\omega)^{l+k}} \int_{L_1} \dots \int_{L_{l+k}} \frac{\Gamma(\alpha + \sum_{j=1}^l h_j s_j + \sum_{j=1}^k s_{l+j})}{\Gamma(\alpha + \beta + \sum_{j=1}^l h_j s_j + \sum_{j=1}^k s_{l+j})} \prod_{j=1}^l \Gamma(\lambda_j + s_j) \prod_{j=1}^k \Gamma(-\sigma_j + s_{l+j})$$

$$\prod_{j=1}^{l+k} \Gamma(-s_j) z_1^{s_1} \dots z_l^{s_l} z_{l+1}^{s_{l+1}} \dots z_{l+k}^{s_{l+k}} ds_1 \dots ds_{l+k} \tag{2.3}$$

Here the contour $L'_j s$ are defined by $L_j = L_{\omega\zeta_j\infty}(Re(\zeta_j) = v''_j)$ starting at the point $v''_j - \omega\infty$ and terminating at the point $v''_j + \omega\infty$ with $v''_j \in \mathbb{R}(j = 1, \dots, l)$ and each of the remaining contour L_{l+1}, \dots, L_{l+k} run from $-\omega\infty$ to $\omega\infty$

(2.2) can be easily established by expanding $\prod_{j=1}^l [1 - \tau_j(t - a)^{h_i}]^{-\lambda_j}$ by means of the formula :

$$(1 - z)^{-\alpha} = \sum_{r=0}^{\infty} \frac{(\alpha)_r}{r!} z^r (|z| < 1) \tag{2.4}$$

integrating term by term with the help of the integral given by Saigo and Saxena [1, page 93, eq.(3.2)] and applying the definition of the generalized Lauricella function [6, page 454].

In this section , we note :

$$\theta_i = \prod_{j=1}^l [1 - \tau_j(t - a)^{h_i}]^{-\zeta_j^{(i)}}, \zeta_j^{(i)} > 0 (i = 1, \dots, r); \theta'_i = \prod_{j=1}^l [1 - \tau_j(t - a)^{h_i}]^{-\zeta_j'^{(i)}}, \zeta_j'^{(i)} > 0 (i = 1, \dots, s)$$

$$\theta''_i = \prod_{j=1}^l [1 - \tau_j(t - a)^{h_i}]^{-\zeta_j''^{(i)}}, \zeta_j''^{(i)} > 0 (i = 1, \dots, u)$$

$$\theta'''_i = \prod_{j=1}^l [1 - \tau_j(t - a)^{h_i}]^{-\zeta_j'''^{(i)}}, \zeta_j'''^{(i)} > 0 (i = 1, \dots, v) \tag{3.1}$$

$$K_1 = (1 - \alpha - \sum_{i=1}^u R_i a_i - \sum_{i=1}^v K_i a'_i; \mu_1, \dots, \mu_r, \mu'_1, \dots, \mu'_s, h_1, \dots, h_l, 1, \dots, 1) \tag{3.2}$$

$$K_2 = (1 - \beta - \sum_{i=1}^u R_i b_i - \sum_{i=1}^v K_i b'_i; \rho_1, \dots, \rho_r, \rho'_1, \dots, \rho'_s, 0, \dots, 0, 0 \dots, 0) \tag{3.3}$$

$$K_j = [1 - \lambda_j - \sum_{i=1}^u R_i \zeta_j''^{(i)} - \sum_{i=1}^v K_i \zeta_j'''^{(i)}; \zeta_j^{(1)}, \dots, \zeta_j^{(r)}, \zeta_j'^{(1)}, \dots, \zeta_j'^{(s)}, 0, \dots, 1, \dots, 0, 0 \dots, 0]_{1,l} \tag{3.4}$$

$$K'_j = [1 + \sigma_j - \sum_{i=1}^u R_i \lambda_j''^{(i)} - \sum_{i=1}^v K_i \lambda_j'''^{(i)}; \lambda_j^{(1)}, \dots, \lambda_j^{(r)}, \lambda_j'^{(1)}, \dots, \lambda_j'^{(s)}, 0, \dots, 0, 0 \dots, 1, \dots, 0]_{1,k} \tag{3.5}$$

$$L_1 = (1 - \alpha - \beta - \sum_{i=1}^u R_i (a_i + b_i) - \sum_{i=1}^v (a'_i + b'_i) K_i; \mu_1 + \rho_1, \dots, \mu_r + \rho_r, \mu'_1 + \rho'_1, \dots, \mu'_r + \rho'_r, h_1, \dots, h_l, 1, \dots, 1) \tag{3.6}$$

$$L_j = [1 - \lambda_j - \sum_{i=1}^u R_i \zeta_j^{''(i)} - \sum_{i=1}^v \zeta_j^{'''(i)} K_i; \zeta_j^{(1)}, \dots, \zeta_j^{(r)}, \zeta_j^{'(1)} \dots, \zeta_j^{'(s)}, 0, \dots, 0, 0 \dots, 0]_{1,l} \quad (3.7)$$

$$L'_j = [1 + \sigma_j - \sum_{i=1}^u R_i \lambda_j^{''(i)} - \sum_{i=1}^v \lambda_j^{'''(i)} K_i; \lambda_j^{(1)}, \dots, \lambda_j^{(r)}, \lambda_j^{'(1)} \dots, \lambda_j^{'(s)}, 0, \dots, 0, 0, \dots, 0]_{1,k} \quad (3.8)$$

$$P_1 = (b - a)^{\alpha+\beta-1} \left\{ \prod_{j=1}^h (af_j + g_j)^{\sigma_j} \right\} \quad (3.9)$$

$$B_{u,v} = (b - a)^{\sum_{i=1}^v K_i(a'_i+b'_i) + \sum_{i=1}^u (a_i+b_i)R_i} \left\{ \prod_{j=1}^h (af_j + g_j)^{-\sum_{i=1}^v K_i \lambda_i^{'''} - \sum_{i=1}^u \lambda_i^{''} R_i} \right\} \quad (3.10)$$

$$V_1 = V; V'; 1, 0; \dots; 1, 0; 1, 0; \dots; 1, 0; W_1 = W; W'; 0, 1; \dots; 0, 1; 0, 1; \dots; 0, 1 \quad (3.11)$$

$$C_1 = C; C'; (1, 0), \dots, (1, 0); (1, 0), \dots, (1, 0); D_1 = D; D'; (0, 1), \dots, (0, 1); (0, 1), \dots, (0, 1) \quad (3.12)$$

We have the following integral : $\int_a^b (t - a)^{\alpha-1} (b - t)^{\beta-1} \prod_{j=1}^l [1 - \tau_j (t - a)^{h_j}]^{-\lambda_j} \prod_{j=1}^k (f_j t + g_j)^{\sigma_j}$

$$S_L^{h_1, \dots, h_u} \left(\begin{matrix} z_1'' \theta_1'' (t - a)^{a_1} (b - t)^{b_1} \prod_{j=1}^k (f_j t + g_j)^{-\lambda_j^{''(1)}} \\ \vdots \\ z_u'' \theta_u'' (t - a)^{a_u} (b - t)^{b_u} \prod_{j=1}^k (f_j t + g_j)^{-\lambda_j^{''(u)}} \end{matrix} \right)$$

$$S_{N_1, \dots, N_v}^{\mathfrak{M}_1, \dots, \mathfrak{M}_v} \left(\begin{matrix} z_1''' \theta_1''' (t - a)^{a'_1} (b - t)^{b'_1} \prod_{j=1}^k (f_j t + g_j)^{-\lambda_j^{'''(1)}} \\ \vdots \\ z_v''' \theta_v''' (t - a)^{a'_v} (b - t)^{b'_v} \prod_{j=1}^k (f_j t + g_j)^{-\lambda_j^{'''(v)}} \end{matrix} \right)$$

$$\aleph \left(\begin{matrix} z_1 \theta_1 (t - a)^{\mu_1} (b - t)^{\rho_1} \prod_{j=1}^k (f_j t + g_j)^{-\lambda_j^{(1)}} \\ \vdots \\ z_r \theta_r (t - a)^{\mu_r} (b - t)^{\rho_r} \prod_{j=1}^k (f_j t + g_j)^{-\lambda_j^{(r)}} \end{matrix} \right)$$

$$\aleph \left(\begin{matrix} z'_1 \theta'_1 (t - a)^{\mu'_1} (b - t)^{\rho'_1} \prod_{j=1}^k (f_j t + g_j)^{-\lambda_j^{'(1)}} \\ \vdots \\ z'_s \theta'_s (t - a)^{\mu'_s} (b - t)^{\rho'_s} \prod_{j=1}^k (f_j t + g_j)^{-\lambda_j^{'(s)}} \end{matrix} \right) dt$$

$$(E) \operatorname{Re} \left(\alpha + \sum_{i=1}^v K_i a'_i + \sum_{i=1}^u R_i a_i + \sum_{i=1}^r \mu_i s_i + \sum_{i=1}^s t_i \mu'_i \right) > 0$$

$$\operatorname{Re} \left(\beta + \sum_{i=1}^v K_i b'_i + \sum_{i=1}^u R_i b_i + \sum_{i=1}^r v_i s_i + \sum_{i=1}^s t_i \rho'_i \right) > 0$$

$$\operatorname{Re} \left(\lambda_j + \sum_{i=1}^v K_i \lambda_j^{(i)} + \sum_{i=1}^u R_i \lambda_j^{(i)} + \sum_{i=1}^r s_i \zeta_j^{(i)} + \sum_{i=1}^s t_i \zeta_j^{(i)} \right) > 0 (j = 1, \dots, l);$$

$$\operatorname{Re} \left(-\sigma_j + \sum_{i=1}^v K_i \lambda_j^{(i)} + \sum_{i=1}^u R_i \lambda_j^{(i)} + \sum_{i=1}^r s_i \lambda_j^{(i)} + \sum_{i=1}^s t_i \lambda_j^{(i)} \right) > 0 (j = 1, \dots, k)$$

$$(F) U_i^{(k)} = \sum_{j=1}^n \alpha_j^{(k)} + \tau_i \sum_{j=n+1}^{p_i} \alpha_{ji}^{(k)} + \sum_{j=1}^{n_k} \gamma_j^{(k)} + \tau_{i(k)} \sum_{j=n_k+1}^{p_i(k)} \gamma_{ji}^{(k)} - \tau_i \sum_{j=1}^{q_i} \beta_{ji}^{(k)} - \sum_{j=1}^{m_k} \delta_j^{(k)} - \tau_{i(k)} \sum_{j=m_k+1}^{q_i(k)} \delta_{ji}^{(k)} \leq 0$$

$$U_i^{(k)} = \sum_{j=1}^{n'} \mu_j^{(k)} + \iota_i \sum_{j=n'+1}^{p'_i} \mu_{ji}^{(k)} + \sum_{j=1}^{n'_k} \alpha_j^{(k)} + \iota_{i(k)} \sum_{j=n'_k+1}^{p'_i(k)} \alpha_{ji}^{(k)} - \iota_i \sum_{j=1}^{q'_i} v_{ji}^{(k)} - \sum_{j=1}^{m'_k} \beta_j^{(k)} - \iota_{i(k)} \sum_{j=m'_k+1}^{q'_i(k)} \beta_{ji}^{(k)} \leq 0$$

$$(G) A_i^{(k)} = \sum_{j=1}^n \alpha_j^{(k)} - \tau_i \sum_{j=n+1}^{p_i} \alpha_{ji}^{(k)} - \tau_i \sum_{j=1}^{q_i} \beta_{ji}^{(k)} + \sum_{j=1}^{n_k} \gamma_j^{(k)} - \tau_{i(k)} \sum_{j=n_k+1}^{p_i(k)} \gamma_{ji}^{(k)} + \sum_{j=1}^{m_k} \delta_j^{(k)} - \tau_{i(k)} \sum_{j=m_k+1}^{q_i(k)} \delta_{ji}^{(k)} - \sum_{l=1}^k \lambda_j^{(i)} - \sum_{l=1}^l \zeta_j^{(i)} - \mu_k - \rho_k > 0, \text{ with } k = 1 \dots, r,$$

$$i = 1, \dots, R, i^{(k)} = 1, \dots, R^{(k)}$$

$$B_i^{(k)} = \sum_{j=1}^{n'} \mu_j^{(k)} - \iota_i \sum_{j=n'+1}^{p'_i} \mu_{ji}^{(k)} - \iota_i \sum_{j=1}^{q'_i} v_{ji}^{(k)} + \sum_{j=1}^{n'_k} \alpha_j^{(k)} - \iota_{i(k)} \sum_{j=n'_k+1}^{p'_i(k)} \alpha_{ji}^{(k)} + \sum_{j=1}^{m'_k} \beta_j^{(k)} - \iota_{i(k)} \sum_{j=m'_k+1}^{q'_i(k)} \beta_{ji}^{(k)} - \sum_{l=1}^k \lambda_j^{(i)} - \sum_{l=1}^l \zeta_j^{(i)} - \mu'_k - \rho'_k > 0, \text{ with } k = 1, \dots, s,$$

$$i = 1, \dots, r, i^{(k)} = 1, \dots, r^{(k)}$$

$$(H) \left| \arg \left(z_i \prod_{j=1}^l [1 - \tau_j(t-a)^{h_i}]^{-\zeta_j^{(i)}} \prod_{j=1}^k (f_j t + g_j)^{-\lambda_j^{(i)}} \right) \right| < \frac{1}{2} A_i^{(k)} \pi \quad (a \leq t \leq b; i = 1, \dots, r)$$

$$\left| \arg \left(z'_i \prod_{j=1}^l [1 - \tau'_j(t-a)^{h'_i}]^{-\zeta'_j{}^{(i)}} \prod_{j=1}^k (f_j t + g_j)^{-\lambda'_j{}^{(i)}} \right) \right| < \frac{1}{2} B_i^{(k)} \pi \quad (a \leq t \leq b; i = 1, \dots, s)$$

Proof

To prove (3.13), first expressing a class of multivariable polynomials $S_{N_1, \dots, N_v}^{\mathfrak{M}_1, \dots, \mathfrak{M}_v} [.]$ defined by Srivastava [3] in serie with the help of (1.32), a class of multivariable polynomials $S_L^{h_1, \dots, h_u} [.]$ defined by Srivastava et al [5] in serie with the help of (1.33) and we interchange the order of summations and t-integral (which is permissible under the conditions stated). Expressing the Aleph-functions of r-variables and s-variables in terms of Mellin-Barnes type contour integral with the help of (1.8) and (1.20) respectively and interchange the order of integrations which is justifiable due to absolute convergence of the integral involved in the process. Now collect the power of $[1 - \tau_j(t-a)^{h_i}]$ with $(i = 1, \dots, r; j = 1, \dots, l)$ and collect the power of $(f_j t + g_j)$ with $j = 1, \dots, k$. Use the equations (2.2) and (2.3) and express the result in Mellin-Barnes contour integral. Interpreting the $(r + s + k + l)$ dimensional Mellin-Barnes integral in multivariable Aleph-function, we obtain the equation (3.13).

Remarks

If a) $\rho_1 = \dots, \rho_r = \rho'_1 = \dots, \rho'_s = 0$; b) $\mu_1 = \dots, \mu_r = \mu'_1 = \dots, \mu'_s = 0$, we obtain the similar formulas that (3.13) with the corresponding simplifications.

4. Particular cases

a) If $\tau_i, \tau_{i(1)}, \dots, \tau_{i(r)}, l_i, l_{i(1)}, \dots, l_{i(s)} \rightarrow 1$, the multivariable Aleph-functions of r and s-variables reduces to multivariable I-functions of r and s-variables defined by Sharma and al [2] respectively and we have

$$\int_a^b (t-a)^{\alpha-1} (b-t)^{\beta-1} \prod_{j=1}^l [1 - \tau_j(t-a)^{h_i}]^{-\lambda_j} \prod_{j=1}^k (f_j t + g_j)^{\sigma_j}$$

$$S_L^{h_1, \dots, h_u} \left(\begin{matrix} z_1'' \theta_1'' (t-a)^{a_1} (b-t)^{b_1} \prod_{j=1}^k (f_j t + g_j)^{-\lambda_j''^{(1)}} \\ \vdots \\ z_u'' \theta_u'' (t-a)^{a_u} (b-t)^{b_u} \prod_{j=1}^k (f_j t + g_j)^{-\lambda_j''^{(u)}} \end{matrix} \right)$$

$$S_{N_1, \dots, N_v}^{\mathfrak{M}_1, \dots, \mathfrak{M}_v} \left(\begin{matrix} z_1''' \theta_1''' (t-a)^{a'_1} (b-t)^{b'_1} \prod_{j=1}^k (f_j t + g_j)^{-\lambda_j'''^{(1)}} \\ \vdots \\ z_v''' \theta_v''' (t-a)^{a'_v} (b-t)^{b'_v} \prod_{j=1}^k (f_j t + g_j)^{-\lambda_j'''^{(v)}} \end{matrix} \right)$$

under the same conditions and notations that (3.13) with $\tau_i, \tau_{i(1)}, \dots, \tau_{i(r)}, l_i, l_{i(1)}, \dots, l_{i(s)} \rightarrow 1$

$$b) \text{ If } B(L; R_1, \dots, R_u) = \frac{\prod_{j=1}^{\bar{A}} (a_j)_{R_1 \theta_j' + \dots + R_u \theta_j^{(u)}} \prod_{j=1}^{B'} (b_j')_{R_1 \phi_j'} \dots \prod_{j=1}^{B^{(u)}} (b_j^{(u)})_{R_u \phi_j^{(u)}}}{\prod_{j=1}^{\bar{C}} (c_j)_{R_1 \psi_j' + \dots + R_u \psi_j^{(u)}} \prod_{j=1}^{D'} (d_j')_{R_1 \delta_j'} \dots \prod_{j=1}^{D^{(u)}} (d_j^{(u)})_{R_u \delta_j^{(u)}}} \quad (4.2)$$

then the general class of multivariable polynomial $S_L^{h_1, \dots, h_u} [z_1, \dots, z_u]$ reduces to generalized Lauricella function defined by Srivastava et al [4]. We have

$$\int_a^b (t-a)^{\alpha-1} (b-t)^{\beta-1} \prod_{j=1}^l [1 - \tau_j (t-a)^{h_i}]^{-\lambda_j} \prod_{j=1}^k (f_j t + g_j)^{\sigma_j}$$

$$F_{\bar{C}; D'; \dots; D^{(u)}}^{1+\bar{A}; B'; \dots; B^{(u)}} \left(\begin{matrix} z_1'' \theta_1'' (t-a)^{a_1} (b-t)^{b_1} \prod_{j=1}^k (f_j t + g_j)^{-\lambda_j''(1)} \\ \vdots \\ z_u'' \theta_u'' (t-a)^{a_u} (b-t)^{b_u} \prod_{j=1}^k (f_j t + g_j)^{-\lambda_j''(u)} \end{matrix} \right)$$

$$\left[(-L); R_1, \dots, R_u \right] [(a); \theta', \dots, \theta^{(u)}] : [(b'); \phi']; \dots; [(b^{(u)}); \phi^{(u)}] \\ [(c); \psi', \dots, \psi^{(u)}] : [(d'); \delta']; \dots; [(d^{(u)}); \delta^{(u)}]$$

$$S_{N_1, \dots, N_v}^{\mathfrak{M}_1, \dots, \mathfrak{M}_v} \left(\begin{matrix} z_1''' \theta_1''' (t-a)^{a_1'} (b-t)^{b_1'} \prod_{j=1}^k (f_j t + g_j)^{-\lambda_j'''(1)} \\ \vdots \\ z_v''' \theta_v''' (t-a)^{a_v'} (b-t)^{b_v'} \prod_{j=1}^k (f_j t + g_j)^{-\lambda_j'''(v)} \end{matrix} \right)$$

$$\mathfrak{N} \left(\begin{matrix} z_1 \theta_1 (t-a)^{\mu_1} (b-t)^{\rho_1} \prod_{j=1}^k (f_j t + g_j)^{-\lambda_j^{(1)}} \\ \vdots \\ z_r \theta_r (t-a)^{\mu_r} (b-t)^{\rho_r} \prod_{j=1}^k (f_j t + g_j)^{-\lambda_j^{(r)}} \end{matrix} \right)$$

$$\mathfrak{N} \left(\begin{matrix} z_1' \theta_1' (t-a)^{\mu_1'} (b-t)^{\rho_1'} \prod_{j=1}^k (f_j t + g_j)^{-\lambda_j'^{(1)}} \\ \vdots \\ z_s' \theta_s' (t-a)^{\mu_s'} (b-t)^{\rho_s'} \prod_{j=1}^k (f_j t + g_j)^{-\lambda_j'^{(s)}} \end{matrix} \right) dt$$

REFERENCES

- [1] Saigo M. and Saxena R.K. Unified fractional integral formulas for the multivariable H-function I. J.Fractional Calculus 15 (1999), page 91-107.
- [2] Sharma C.K.and Ahmad S.S.: On the multivariable I-function. Acta ciencia Indica Math , 1994 vol 20,no2, p 113-116.
- [3] Srivastava H.M. A multilinear generating function for the Konhauser set of biorthogonal polynomials suggested by Laguerre polynomial, Pacific. J. Math. 177(1985), page183-191.
- [4] Srivastava H.M. and Daoust M.C. Certain generalized Neumann expansions associated with Kampé de Fériet function. Nederl. Akad. Wetensch. Proc. Ser A72 = Indag Math 31(1969) page 449-457.
- [5] Srivastava H.M. And Garg M. Some integral involving a general class of polynomials and multivariable H-function. Rev. Roumaine Phys. 32(1987), page 685-692.
- [6] Srivastava H.M. and Karlsson P.W. Multiple Gaussian Hypergeometric series. Ellis.Horwood. Limited. New-York, Chichester. Brisbane. Toronto , 1985.
- [7] H.M. Srivastava and R.Panda. Some expansion theorems and generating relations for the H-function of several complex variables. Comment. Math. Univ. St. Paul. 24(1975), p.119-137.

Personal adress : 411 Avenue Joseph Raynaud
Le parc Fleuri , Bat B
83140 , Six-Fours les plages
Tel : 06-83-12-49-68
Department : VAR
Country : FRANCE