

Finite integral involving the spheroidal function, a class of polynomials multivariable Aleph-functions and Fresnel integral

F.Y. AYANT¹

¹ Teacher in High School , France

ABSTRACT

In the present paper we evaluate a generalized finite integral involving the product of the spheroidal function, the Fresnel integral, the multivariable Aleph-functions and general class of polynomials of several variables with general arguments. The importance of the result established in this paper lies in the fact they involve the Aleph-function of several variables which is sufficiently general in nature and capable to yielding a large of results merely by specializing the parameters their in.

Keywords: Multivariable Aleph-function, general class of polynomials, spheroidal function, multivariable I-function, Aleph-function of two variable, I-function of two variables, Fresnel integral.

2010 Mathematics Subject Classification. 33C99, 33C60, 44A20

1. Introduction and preliminaries.

The function Aleph of several variables generalize the multivariable I-function recently study by C.K. Sharma and Ahmad [4] , itself is an a generalisation of G and H-functions of multiple variables. The multiple Mellin-Barnes integral occuring in this paper will be referred to as the multivariables Aleph-function throughout our present study and will be defined and represented as follows.

$$\text{We define : } \mathfrak{N}(z_1, \dots, z_r) = \mathfrak{N}_{p_i, q_i, \tau_i; R; p_i(1), q_i(1), \tau_i(1); R^{(1)}; \dots; p_i(r), q_i(r), \tau_i(r); R^{(r)}}^{0, n; m_1, n_1, \dots, m_r, n_r} \left(\begin{matrix} z_1 \\ \cdot \\ \cdot \\ \cdot \\ z_r \end{matrix} \right)$$

$$[(a_j; \alpha_j^{(1)}, \dots, \alpha_j^{(r)})_{1, n}] , [\tau_i(a_{ji}; \alpha_{ji}^{(1)}, \dots, \alpha_{ji}^{(r)})_{n+1, p_i}] :$$

$$\dots, [\tau_i(b_{ji}; \beta_{ji}^{(1)}, \dots, \beta_{ji}^{(r)})_{m+1, q_i}] :$$

$$\left[[(c_j^{(1)}); \gamma_j^{(1)})_{1, n_1}], [\tau_{i(1)}(c_{ji(1)}^{(1)}; \gamma_{ji(1)}^{(1)})_{n_1+1, p_i^{(1)}}]; \dots ; [(c_j^{(r)}); \gamma_j^{(r)})_{1, n_r}], [\tau_{i(r)}(c_{ji(r)}^{(r)}; \gamma_{ji(r)}^{(r)})_{n_r+1, p_i^{(r)}}] \right]$$

$$\left[[(d_j^{(1)}); \delta_j^{(1)})_{1, m_1}], [\tau_{i(1)}(d_{ji(1)}^{(1)}; \delta_{ji(1)}^{(1)})_{m_1+1, q_i^{(1)}}]; \dots ; [(d_j^{(r)}); \delta_j^{(r)})_{1, m_r}], [\tau_{i(r)}(d_{ji(r)}^{(r)}; \delta_{ji(r)}^{(r)})_{m_r+1, q_i^{(r)}}] \right]$$

$$= \frac{1}{(2\pi\omega)^r} \int_{L_1} \dots \int_{L_r} \psi(s_1, \dots, s_r) \prod_{k=1}^r \theta_k(s_k) y_k^{s_k} ds_1 \dots ds_r \tag{1.1}$$

with $\omega = \sqrt{-1}$

$$\psi(s_1, \dots, s_r) = \frac{\prod_{j=1}^n \Gamma(1 - a_j + \sum_{k=1}^r \alpha_j^{(k)} s_k)}{\sum_{i=1}^R [\tau_i \prod_{j=n+1}^{p_i} \Gamma(a_{ji} - \sum_{k=1}^r \alpha_{ji}^{(k)} s_k) \prod_{j=1}^{q_i} \Gamma(1 - b_{ji} + \sum_{k=1}^r \beta_{ji}^{(k)} s_k)]} \tag{1.2}$$

$$\text{and } \theta_k(s_k) = \frac{\prod_{j=1}^{m_k} \Gamma(d_j^{(k)} - \delta_j^{(k)} s_k) \prod_{j=1}^{n_k} \Gamma(1 - c_j^{(k)} + \gamma_j^{(k)} s_k)}{\sum_{i^{(k)}=1}^{R^{(k)}} [\tau_{i^{(k)}} \prod_{j=m_k+1}^{q_i^{(k)}} \Gamma(1 - d_{ji^{(k)}}^{(k)} + \delta_{ji^{(k)}}^{(k)} s_k) \prod_{j=n_k+1}^{p_i^{(k)}} \Gamma(c_{ji^{(k)}}^{(k)} - \gamma_{ji^{(k)}}^{(k)} s_k)]} \quad (1.3)$$

Suppose, as usual, that the parameters

$$a_j, j = 1, \dots, p; b_j, j = 1, \dots, q;$$

$$c_j^{(k)}, j = 1, \dots, n_k; c_{ji^{(k)}}^{(k)}, j = n_k + 1, \dots, p_i^{(k)};$$

$$d_j^{(k)}, j = 1, \dots, m_k; d_{ji^{(k)}}^{(k)}, j = m_k + 1, \dots, q_i^{(k)};$$

$$\text{with } k = 1, \dots, r, i = 1, \dots, R, i^{(k)} = 1, \dots, R^{(k)}$$

are complex numbers, and the $\alpha's, \beta's, \gamma's$ and $\delta's$ are assumed to be positive real numbers for standardization purpose such that

$$U_i^{(k)} = \sum_{j=1}^n \alpha_j^{(k)} + \tau_i \sum_{j=n+1}^{p_i} \alpha_{ji}^{(k)} + \sum_{j=1}^{n_k} \gamma_j^{(k)} + \tau_{i^{(k)}} \sum_{j=n_k+1}^{p_i^{(k)}} \gamma_{ji^{(k)}}^{(k)} - \tau_i \sum_{j=1}^{q_i} \beta_{ji}^{(k)} - \sum_{j=1}^{m_k} \delta_j^{(k)} - \tau_{i^{(k)}} \sum_{j=m_k+1}^{q_i^{(k)}} \delta_{ji^{(k)}}^{(k)} \leq 0 \quad (1.4)$$

The real numbers τ_i are positives for $i = 1$ to R , $\tau_{i^{(k)}}$ are positives for $i^{(k)} = 1$ to $R^{(k)}$

The contour L_k is in the s_k -p lane and run from $\sigma - i\infty$ to $\sigma + i\infty$ where σ is a real number with loop, if necessary, ensure that the poles of $\Gamma(d_j^{(k)} - \delta_j^{(k)} s_k)$ with $j = 1$ to m_k are separated from those of $\Gamma(1 - a_j + \sum_{i=1}^r \alpha_j^{(k)} s_k)$ with $j = 1$ to n and $\Gamma(1 - c_j^{(k)} + \gamma_j^{(k)} s_k)$ with $j = 1$ to n_k to the left of the

contour L_k . The condition for absolute convergence of multiple Mellin-Barnes type contour (1.9) can be obtained by extension of the corresponding conditions for multivariable H-function given by as :

$$|\arg z_k| < \frac{1}{2} A_i^{(k)} \pi, \text{ where}$$

$$A_i^{(k)} = \sum_{j=1}^n \alpha_j^{(k)} - \tau_i \sum_{j=n+1}^{p_i} \alpha_{ji}^{(k)} - \tau_i \sum_{j=1}^{q_i} \beta_{ji}^{(k)} + \sum_{j=1}^{n_k} \gamma_j^{(k)} - \tau_{i^{(k)}} \sum_{j=n_k+1}^{p_i^{(k)}} \gamma_{ji^{(k)}}^{(k)} + \sum_{j=1}^{m_k} \delta_j^{(k)} - \tau_{i^{(k)}} \sum_{j=m_k+1}^{q_i^{(k)}} \delta_{ji^{(k)}}^{(k)} > 0, \text{ with } k = 1, \dots, r, i = 1, \dots, R, i^{(k)} = 1, \dots, R^{(k)} \quad (1.5)$$

The complex numbers z_i are not zero. Throughout this document, we assume the existence and absolute convergence conditions of the multivariable Aleph-function.

We may establish the asymptotic expansion in the following convenient form :

$$\aleph(z_1, \dots, z_r) = O(|z_1|^{\alpha_1}, \dots, |z_r|^{\alpha_r}), \max(|z_1|, \dots, |z_r|) \rightarrow 0$$

$$\aleph(z_1, \dots, z_r) = O(|z_1|^{\beta_1}, \dots, |z_r|^{\beta_r}), \min(|z_1|, \dots, |z_r|) \rightarrow \infty$$

where, with $k = 1, \dots, r : \alpha_k = \min[Re(d_j^{(k)} / \delta_j^{(k)})], j = 1, \dots, m_k$ and

$$\beta_k = \max[\operatorname{Re}((c_j^{(k)} - 1)/\gamma_j^{(k)})], j = 1, \dots, n_k$$

Series representation of Aleph-function of several variables is given by

$$\aleph(y_1, \dots, y_r) = \sum_{G_1, \dots, G_r=0}^{\infty} \sum_{g_1=0}^{m_1} \dots \sum_{g_r=0}^{m_r} \frac{(-)^{G_1+\dots+G_r}}{\delta_{g_1}^{G_1} \dots \delta_{g_r}^{G_r}} \psi(\eta_{G_1, g_1}, \dots, \eta_{G_r, g_r}) \times \theta_1(\eta_{G_1, g_1}) \dots \theta_r(\eta_{G_r, g_r}) y_1^{-\eta_{G_1, g_1}} \dots y_r^{-\eta_{G_r, g_r}} \tag{1.6}$$

Where $\psi(\dots), \theta_i(\dots), i = 1, \dots, r$ are given respectively in (1.2), (1.3) and

$$\eta_{G_1, g_1} = \frac{d_{g_1}^{(1)} + G_1}{\delta_{g_1}^{(1)}}, \dots, \eta_{G_r, g_r} = \frac{d_{g_r}^{(r)} + G_r}{\delta_{g_r}^{(r)}}$$

which is valid under the conditions $\delta_{g_i}^{(i)} [d_j^i + p_i] \neq \delta_{j_i}^{(i)} [d_{g_i}^i + G_i]$ (1.7)

for $j \neq m_i, m_i = 1, \dots, \eta_{G_i, g_i}; p_i, n_i = 0, 1, 2, \dots, ; y_i \neq 0, i = 1, \dots, r$ (1.8)

Consider the Aleph-function of s variables

$$\aleph(z_1, \dots, z_s) = \aleph_{P_i, Q_i, \mu_i; r': P_i^{(1)}, Q_i^{(1)}, \mu_i^{(1)}; \dots; P_i^{(s)}, Q_i^{(s)}, \mu_i^{(s)}; r^{(s)}}^{0, N; M_1, N_1, \dots, M_s, N_s} \left(\begin{matrix} z_1 \\ \cdot \\ \cdot \\ \cdot \\ z_s \end{matrix} \right)$$

$$[(u_j; \mu_j^{(1)}, \dots, \mu_j^{(r')})_{1, N_1}, \dots, [l_i(u_{ji}; \mu_{ji}^{(1)}, \dots, \mu_{ji}^{(r')})_{N_1+1, P_i}], \dots, [l_i(v_{ji}; \nu_{ji}^{(1)}, \dots, \nu_{ji}^{(r')})_{M_1+1, Q_i}], \dots, [(a_j^{(1)}; \alpha_j^{(1)})_{1, N_1}], [l_{i(1)}(a_{ji(1)}^{(1)}; \alpha_{ji(1)}^{(1)})_{N_1+1, P_i^{(1)}}]; \dots; [(a_j^{(s)}; \alpha_j^{(s)})_{1, N_s}], [l_{i(s)}(a_{ji(s)}^{(s)}; \alpha_{ji(s)}^{(s)})_{N_s+1, P_i^{(s)}}] \dots [(b_j^{(1)}; \beta_j^{(1)})_{1, M_1}], [l_{i(1)}(b_{ji(1)}^{(1)}; \beta_{ji(1)}^{(1)})_{M_1+1, Q_i^{(1)}}]; \dots; [(b_j^{(s)}; \beta_j^{(s)})_{1, M_s}], [l_{i(s)}(b_{ji(s)}^{(s)}; \beta_{ji(s)}^{(s)})_{M_s+1, Q_i^{(s)}}]$$

$$= \frac{1}{(2\pi\omega)^s} \int_{L_1} \dots \int_{L_s} \zeta(t_1, \dots, t_s) \prod_{k=1}^s \phi_k(t_k) z_k^{t_k} dt_1 \dots dt_s \tag{1.9}$$

with $\omega = \sqrt{-1}$

$$\zeta(t_1, \dots, t_s) = \frac{\prod_{j=1}^N \Gamma(1 - u_j + \sum_{k=1}^s \mu_j^{(k)} t_k)}{\sum_{i=1}^{r'} [l_i \prod_{j=N+1}^{P_i} \Gamma(u_{ji} - \sum_{k=1}^s \mu_{ji}^{(k)} t_k) \prod_{j=1}^{Q_i} \Gamma(1 - v_{ji} + \sum_{k=1}^s \nu_{ji}^{(k)} t_k)]} \tag{1.10}$$

and $\phi_k(t_k) = \frac{\prod_{j=1}^{M_k} \Gamma(b_j^{(k)} - \beta_j^{(k)} t_k) \prod_{j=1}^{N_k} \Gamma(1 - a_j^{(k)} + \alpha_j^{(k)} t_k)}{\sum_{i^{(k)}=1}^{r^{(k)}} [l_{i^{(k)}} \prod_{j=M_k+1}^{Q_{i^{(k)}}} \Gamma(1 - b_{ji^{(k)}}^{(k)} + \beta_{ji^{(k)}}^{(k)} t_k) \prod_{j=N_k+1}^{P_{i^{(k)}}} \Gamma(a_{ji^{(k)}}^{(k)} - \alpha_{ji^{(k)}}^{(k)} t_k)]} \tag{1.11}$

Suppose , as usual , that the parameters

$$u_j, j = 1, \dots, P; v_j, j = 1, \dots, Q;$$

$$a_j^{(k)}, j = 1, \dots, N_k; a_{j i^{(k)}}^{(k)}, j = n_k + 1, \dots, P_{i^{(k)}};$$

$$b_{j i^{(k)}}^{(k)}, j = m_k + 1, \dots, Q_{i^{(k)}}; b_j^{(k)}, j = 1, \dots, M_k;$$

$$\text{with } k = 1 \dots, s, i = 1, \dots, r', i^{(k)} = 1, \dots, r^{(k)}$$

are complex numbers , and the $\alpha' s, \beta' s, \gamma' s$ and $\delta' s$ are assumed to be positive real numbers for standardization purpose such that

$$U_i^{(k)} = \sum_{j=1}^N \mu_j^{(k)} + \iota_i \sum_{j=N+1}^{P_i} \mu_{j i}^{(k)} + \sum_{j=1}^{N_k} \alpha_j^{(k)} + \iota_{i^{(k)}} \sum_{j=N_k+1}^{P_{i^{(k)}}} \alpha_{j i^{(k)}}^{(k)} - \iota_i \sum_{j=1}^{Q_i} v_{j i}^{(k)} - \sum_{j=1}^{M_k} \beta_j^{(k)} - \iota_{i^{(k)}} \sum_{j=M_k+1}^{Q_{i^{(k)}}} \beta_{j i^{(k)}}^{(k)} \leq 0 \tag{1.12}$$

The reals numbers τ_i are positives for $i = 1, \dots, r$, $\iota_{i^{(k)}}$ are positives for $i^{(k)} = 1 \dots r^{(k)}$

The contour L_k is in the t_k -p lane and run from $\sigma - i\infty$ to $\sigma + i\infty$ where σ is a real number with loop , if necessary ,ensure that the poles of $\Gamma(b_j^{(k)} - \beta_j^{(k)} t_k)$ with $j = 1$ to M_k are separated from those of $\Gamma(1 - u_j + \sum_{i=1}^s \mu_j^{(k)} t_k)$ with $j = 1$ to N and $\Gamma(1 - a_j^{(k)} + \alpha_j^{(k)} t_k)$ with $j = 1$ to N_k to the left of the contour L_k . The condition for absolute convergence of multiple Mellin-Barnes type contour (1.9) can be obtained by extension of the corresponding conditions for multivariable H-function given by as :

$$|arg z_k| < \frac{1}{2} B_i^{(k)} \pi, \text{ where}$$

$$B_i^{(k)} = \sum_{j=1}^N \mu_j^{(k)} - \iota_i \sum_{j=N+1}^{P_i} \mu_{j i}^{(k)} - \iota_i \sum_{j=1}^{Q_i} v_{j i}^{(k)} + \sum_{j=1}^{N_k} \alpha_j^{(k)} - \iota_{i^{(k)}} \sum_{j=N_k+1}^{P_{i^{(k)}}} \alpha_{j i^{(k)}}^{(k)} + \sum_{j=1}^{M_k} \beta_j^{(k)} - \iota_{i^{(k)}} \sum_{j=M_k+1}^{Q_{i^{(k)}}} \beta_{j i^{(k)}}^{(k)} > 0, \text{ with } k = 1, \dots, s, i = 1, \dots, r, i^{(k)} = 1, \dots, r^{(k)} \tag{1.13}$$

The complex numbers z_i are not zero. Throughout this document , we assume the existence and absolute convergence conditions of the multivariable Aleph-function.

We may establish the the asymptotic expansion in the following convenient form :

$$\aleph(z_1, \dots, z_s) = O(|z_1|^{\alpha'_1}, \dots, |z_s|^{\alpha'_s}), \max(|z_1|, \dots, |z_s|) \rightarrow 0$$

$$\aleph(z_1, \dots, z_s) = O(|z_1|^{\beta'_1}, \dots, |z_s|^{\beta'_s}), \min(|z_1|, \dots, |z_s|) \rightarrow \infty$$

where, with $k = 1, \dots, s, z : \alpha'_k = \min[Re(b_j^{(k)} / \beta_j^{(k)})], j = 1, \dots, M_k$ and

$$\beta'_k = \max[Re((a_j^{(k)} - 1) / \alpha_j^{(k)})], j = 1, \dots, N_k$$

We will use these following notations in this paper

$$U = P_i, Q_i, l_i; r'; V = M_1, N_1; \dots; M_s, N_s \tag{1.15}$$

$$W = P_{i(1)}, Q_{i(1)}, l_{i(1)}; r^{(1)}, \dots, P_{i(r)}, Q_{i(r)}, l_{i(r)}; r^{(s)} \tag{1.16}$$

$$A = \{(u_j; \mu_j^{(1)}, \dots, \mu_j^{(s)})_{1,N}\}, \{l_i(u_{ji}; \mu_{ji}^{(1)}, \dots, \mu_{ji}^{(s)})_{N+1, P_i}\} \tag{1.17}$$

$$B = \{l_i(v_{ji}; v_{ji}^{(1)}, \dots, v_{ji}^{(s)})_{M+1, Q_i}\} \tag{1.18}$$

$$C = (a_j^{(1)}; \alpha_j^{(1)})_{1, N_1}, l_{i(1)}(a_{ji(1)}^{(1)}; \alpha_{ji(1)}^{(1)})_{N_1+1, P_{i(1)}}, \dots, (a_j^{(s)}; \alpha_j^{(s)})_{1, N_s}, l_{i(s)}(a_{ji(s)}^{(s)}; \alpha_{ji(s)}^{(s)})_{N_s+1, P_{i(s)}} \tag{1.19}$$

$$D = (b_j^{(1)}; \beta_j^{(1)})_{1, M_1}, l_{i(1)}(b_{ji(1)}^{(1)}; \beta_{ji(1)}^{(1)})_{M_1+1, Q_{i(1)}}, \dots, (b_j^{(s)}; \beta_j^{(s)})_{1, M_s}, l_{i(s)}(\beta_{ji(s)}^{(s)}; \beta_{ji(s)}^{(s)})_{M_s+1, Q_{i(s)}} \tag{1.20}$$

The multivariable Aleph-function write :

$$\aleph(z_1, \dots, z_s) = \aleph_{U:W}^{0, N; V} \left(\begin{array}{c|c} z_1 & \text{A : C} \\ \cdot & \cdot \cdot \cdot \\ \cdot & \text{B : D} \\ z_s & \end{array} \right) \tag{1.21}$$

The generalized polynomials defined by Srivastava [7], is given in the following manner :

$$S_{N_1, \dots, N_t}^{M_1, \dots, M_t} [y_1, \dots, y_t] = \sum_{K_1=0}^{[N_1/M_1]} \dots \sum_{K_t=0}^{[N_t/M_t]} \frac{(-N_1)_{M_1 K_1}}{K_1!} \dots \frac{(-N_t)_{M_t K_t}}{K_t!} A[N_1, K_1; \dots; N_t, K_t] y_1^{K_1} \dots y_t^{K_t} \tag{1.22}$$

Where M_1, \dots, M_s are arbitrary positive integers and the coefficients $A[N_1, K_1; \dots; N_t, K_t]$ are arbitrary constants, real or complex. In the present paper, we use the following notation

$$a_1 = \frac{(-N_1)_{M_1 K_1}}{K_1!} \dots \frac{(-N_t)_{M_t K_t}}{K_t!} A[N_1, K_1; \dots; N_t, K_t] \tag{1.23}$$

In the document , we note :

$$G(\eta_{G_1, g_1}, \dots, \eta_{G_r, g_r}) = \phi(\eta_{G_1, g_1}, \dots, \eta_{G_r, g_r}) \theta_1(\eta_{G_1, g_1}) \dots \theta_r(\eta_{G_r, g_r}) \tag{1.24}$$

where $\phi(\eta_{G_1, g_1}, \dots, \eta_{G_r, g_r}), \theta_1(\eta_{G_1, g_1}), \dots, \theta_r(\eta_{G_r, g_r})$ are given respectively in (1.2) and (1.3)

The spheroidal function $\psi_{\alpha n}(c, \eta)$ of general order $\alpha > -1$ can be expanded as ([3] an [8]).

$$\psi_{\alpha n}(c, \eta) = \frac{i^n \sqrt{2\pi}}{V_{\alpha n}(c)} \sum_{k=0, \text{ or } 1}^{\infty} a_k(c|\alpha n) (c\eta)^{-\alpha - \frac{1}{2}} J_{k+\alpha + \frac{1}{2}}(c\eta) \tag{1.25}$$

which represents the function uniformly on (∞, ∞) , where the coefficients $a_k(c|\alpha n)$ satisfy the recursion formula [14 ,eq.67]and the asterisk over the summation sign indicates that the sum is taken over only even or odd values of k according as n is even or odd. As $c \rightarrow 0, a_k(c|\alpha n) \rightarrow 0, k \neq n$

The Fresnel integral denoted $S(z)$ is defined by : see Abramowitz M and Stegun I.A. ([1],page 89(300))

$$S(z) = \int_0^z \sin\left(\frac{\pi}{2}t^2\right) dt \tag{1.25}$$

2. Required integral

We have the following integral, see Brychkow ([2], 4.5.1, 1page 189).

$$\int_0^a x^{s-1}(a-x)^{t-1} S(b\sqrt{x(a-x)}) dx = \frac{a^{s+t+\frac{1}{2}}}{3} \sqrt{\frac{2b^3}{\pi}} B\left(s + \frac{3}{4}, t + \frac{3}{4}\right) \times {}_3F_4\left(\begin{matrix} \frac{3}{4}, s + \frac{3}{4}, t + \frac{3}{4} \\ \frac{3}{2}, \frac{7}{4}, \frac{s+t}{2} + \frac{3}{4}, \frac{s+t}{2} + \frac{5}{4} \end{matrix}; -\frac{(ab)^2}{16}\right) \tag{2.1}$$

where $a > 0, Re(s) > -\frac{3}{4}, Re(t) > -\frac{3}{4}$

3. Main integral

Let $X_{s,t} = x^s(a-x)^t$, We have the following generalized finite integral :

$$\int_0^a x^{s'-1}(a-x)^{t'-1} S(b\sqrt{x(a-x)}) \psi_{\alpha n}(c^\sigma, X_{\alpha,\beta}) S_{N_1,\dots,N_t}^{M_1,\dots,M_t} \begin{pmatrix} y_1 X_{\gamma_1,\mu_1} \\ \dots \\ y_t X_{\gamma_t,\mu_t} \end{pmatrix} \mathbb{N}_{u:v}^{0,n:v} \begin{pmatrix} z_1 X_{\alpha_1,\beta_1} \\ \dots \\ z_r X_{\alpha_r,\beta_r} \end{pmatrix} \mathbb{N}_{U:W}^{0,N:V} \begin{pmatrix} Z_1 X_{\eta_1,\epsilon_1} \\ \dots \\ Z_s X_{\eta_s,\epsilon_s} \end{pmatrix} dx = \frac{a^{s'+t'+\frac{1}{2}}}{3} \sqrt{\frac{2b^3}{\pi}} \sum_{n'=0}^{\infty} \sum_{k=0,or 1}^{\infty} \sum_{m=0}^{\infty} \sum_{G_1,\dots,G_r=0}^{\infty} \sum_{g_1=0}^{m_1} \dots \sum_{g_r=0}^{m_r} \sum_{K_1=0}^{[N_1/M_1]} \dots \sum_{K_t=0}^{[N_t/M_t]} a_1 \frac{(-)^{G_1+\dots+G_r}}{\delta_{g_1} G_1! \dots \delta_{g_r} G_r!} G(\eta_{G_1,g_1}, \dots, \eta_{G_r,g_r}) \frac{(-)^m a_k (c^\sigma | \alpha n)}{m! \Gamma(m+k+\alpha+\frac{3}{2})} \frac{(\frac{3}{4})_{n'} (-ab)^{2n'}}{16^{n'} (\frac{3}{2})_{n'} (\frac{7}{4})_{n'} n'!} x_1^{p_1} \dots x_s^{p_s} z_1^{\eta_{G_1,g_1}} \dots z_r^{\eta_{G_r,g_r}} y_1^{K_1} \dots y_t^{K_t} c^{\sigma(2m+k)}$$

$$a^{(2m+k)(\alpha+\beta)+\sum_{i=1}^t K_i(\gamma_i+\mu_i)+\sum_{i=1}^r \eta_{G_i,g_i}(\alpha_i+\beta_i)} \mathbb{N}_{U_{43}:W}^{0,N+3:V} \begin{pmatrix} Z_1 a^{\eta_1+\epsilon_1} \\ \dots \\ Z_s a^{\eta_s+\epsilon_s} \end{pmatrix}$$

the Mellin-barnes contour integral, we arrive at the desired result.

4. Multivariable I-function

If $l_i, l_{i(1)}, \dots, l_{i(s)} \rightarrow 1$, the Aleph-function of several variables degenerate to the I-function of several variables. The simple integral have been derived in this section for multivariable I-functions defined by Sharma et al [4].

Corollary 1

$$\int_0^a x^{s'-1} (a-x)^{t'-1} S(b\sqrt{x(a-x)}) \psi_{\alpha n}(c^\sigma, X_{\alpha, \beta}) S_{N_1, \dots, N_t}^{M_1, \dots, M_t} \begin{pmatrix} y_1 X_{\gamma_1, \mu_1} \\ \dots \\ y_t X_{\gamma_t, \mu_t} \end{pmatrix} \\ N_{u:w}^{0, n:v} \begin{pmatrix} z_1 X_{\alpha_1, \beta_1} \\ \dots \\ z_r X_{\alpha_r, \beta_r} \end{pmatrix} I_{U:W}^{0, N:V} \begin{pmatrix} Z_1 X_{\eta_1, \epsilon_1} \\ \dots \\ Z_s X_{\eta_s, \epsilon_s} \end{pmatrix} dx = \frac{a^{s'+t'+\frac{1}{2}}}{3} \sqrt{\frac{2b^3}{\pi}} \sum_{n'=0}^{\infty} \sum_{k=0, or 1}^{\infty} \sum_{m=0}^{\infty} \\ \sum_{G_1, \dots, G_r=0}^{\infty} \sum_{g_1=0}^{m_1} \dots \sum_{g_r=0}^{m_r} \sum_{K_1=0}^{[N_1/M_1]} \dots \sum_{K_t=0}^{[N_t/M_t]} a_1 \frac{(-)^{G_1+\dots+G_r}}{\delta_{g_1} G_1! \dots \delta_{g_r} G_r!} G(\eta_{G_1, g_1}, \dots, \eta_{G_r, g_r}) \\ \frac{(-)^m a_k (c^\sigma | \alpha n)}{m! \Gamma(m+k+\alpha+\frac{3}{2})} \frac{(\frac{3}{4})_{n'} (-ab)^{2n'}}{16^{n'} (\frac{3}{2})_{n'} (\frac{7}{4})_{n'} n'!} x_1^{p_1} \dots x_s^{p_s} z_1^{\eta_{G_1, g_1}} \dots z_r^{\eta_{G_r, g_r}} y_1^{K_1} \dots y_t^{K_t} c^{\sigma(2m+k)} \\ a^{(2m+k)(\alpha+\beta)+\sum_{i=1}^t K_i(\gamma_i+\mu_i)+\sum_{i=1}^r \eta_{G_i, g_i}(\alpha_i+\beta_i)} I_{U_{43}:W}^{0, N+4:V} \begin{pmatrix} Z_1 a^{\eta_1+\epsilon_1} \\ \dots \\ Z_s a^{\eta_s+\epsilon_s} \end{pmatrix} \\ (\frac{1}{4}-n'-(s'+(2m+k)\alpha+\sum_{i=1}^t K_i \gamma_i+\sum_{i=1}^r \eta_{G_i, g_i} \alpha_i); \eta_1, \dots, \eta_s), \\ (-\frac{1}{2}-(s'+t'+(2m+k)(\alpha+\beta)+\sum_{i=1}^t K_i(\gamma_i+\mu_i)+\sum_{i=1}^r \eta_{G_i, g_i}(\alpha_i+\beta_i)); \epsilon_1+\eta_1, \dots, \epsilon_s+\eta_s), \\ (\frac{1}{4}-\frac{1}{2}(s'+t'+(2m+k)(\alpha+\beta)+\sum_{i=1}^t K_i(\gamma_i+\mu_i)+\sum_{i=1}^r \eta_{G_i, g_i}(\alpha_i+\beta_i)); \frac{\epsilon_1+\eta_1}{2}, \dots, \frac{\epsilon_s+\eta_s}{2}), \\ (\frac{1}{4}-n'-\frac{1}{2}(s'+t'+(2m+k)(\alpha+\beta)+\sum_{i=1}^t K_i(\gamma_i+\mu_i)+\sum_{i=1}^r \eta_{G_i, g_i}(\alpha_i+\beta_i)); \frac{\epsilon_1+\eta_1}{2}, \dots, \frac{\epsilon_s+\eta_s}{2}), \\ (-\frac{1}{4}-\frac{1}{2}(s'+t'+(2m+k)(\alpha+\beta)+\sum_{i=1}^t K_i(\gamma_i+\mu_i)+\sum_{i=1}^r \eta_{G_i, g_i}(\alpha_i+\beta_i)); \frac{\epsilon_1+\eta_1}{2}, \dots, \frac{\epsilon_s+\eta_s}{2}), \\ (-\frac{1}{4}-n'-\frac{1}{2}(s'+t'+(2m+k)(\alpha+\beta)+\sum_{i=1}^t K_i(\gamma_i+\mu_i)+\sum_{i=1}^r \eta_{G_i, g_i}(\alpha_i+\beta_i)); \frac{\epsilon_1+\eta_1}{2}, \dots, \frac{\epsilon_s+\eta_s}{2}),$$

$$\left(\begin{array}{c} (\frac{1}{4}-n'-(t'+(2m+k)\beta + \sum_{i=1}^t K_i\mu_i + \sum_{i=1}^r \eta_{G_i,g_i}\beta_i); \epsilon_1, \dots, \epsilon_s), A : C \\ \vdots \\ B : D \end{array} \right) \tag{4.1}$$

under the same notations and conditions that (3.1) with $l_i, l_{i(1)}, \dots, l_{i(s)} \rightarrow 1$

5. Aleph-function of two variables

If $s = 2$, we obtain the Aleph-function of two variables defined by K.Sharma [6], and we have the following simple integrals.

Corollary 2

$$\int_0^a x^{s'-1}(a-x)^{t'-1} S(b\sqrt{x(a-x)}) \psi_{\alpha n}(c^\sigma, X_{\alpha,\beta}) S_{N_1,\dots,N_t}^{M_1,\dots,M_t} \left(\begin{array}{c} y_1 X_{\gamma_1,\mu_1} \\ \dots \\ y_t X_{\gamma_t,\mu_t} \end{array} \right)$$

$$\mathfrak{N}_{u:w}^{0,n:v} \left(\begin{array}{c} z_1 X_{\alpha_1,\beta_1} \\ \dots \\ z_r X_{\alpha_r,\beta_r} \end{array} \right) \mathfrak{N}_{U:W}^{0,N:V} \left(\begin{array}{c} Z_1 X_{\eta_1,\epsilon_1} \\ \dots \\ Z_2 X_{\eta_2,\epsilon_2} \end{array} \right) dx = \frac{a^{s'+t'+\frac{1}{2}}}{3} \sqrt{\frac{2b^3}{\pi}} \sum_{n'=0}^{\infty} \sum_{k=0,or1}^{\infty} \sum_{m=0}^{\infty}$$

$$\sum_{G_1,\dots,G_r=0}^{\infty} \sum_{g_1=0}^{m_1} \dots \sum_{g_r=0}^{m_r} \sum_{K_1=0}^{[N_1/M_1]} \dots \sum_{K_t=0}^{[N_t/M_t]} a_1 \frac{(-)^{G_1+\dots+G_r}}{\delta_{g_1} G_1! \dots \delta_{g_r} G_r!} G(\eta_{G_1,g_1}, \dots, \eta_{G_r,g_r})$$

$$\frac{(-)^m a_k (c^\sigma | \alpha n)}{m! \Gamma(m+k+\alpha+\frac{3}{2})} \frac{(\frac{3}{4})_{n'} (-ab)^{2n'}}{16^{n'} (\frac{3}{2})_{n'} (\frac{7}{4})_{n'} n'!} x_1^{p_1} \dots x_s^{p_s} z_1^{\eta_{G_1,g_1}} \dots z_r^{\eta_{G_r,g_r}} y_1^{K_1} \dots y_t^{K_t} c^{\sigma(2m+k)}$$

$$a^{(2m+k)(\alpha+\beta)+\sum_{i=1}^t K_i(\gamma_i+\mu_i)+\sum_{i=1}^r \eta_{G_i,g_i}(\alpha_i+\beta_i)} \mathfrak{N}_{U_{43}:W}^{0,N+4:V} \left(\begin{array}{c} Z_1 a^{\eta_1+\epsilon_1} \\ \dots \\ Z_2 a^{\eta_2+\epsilon_2} \end{array} \right)$$

$$\left(\begin{array}{c} (\frac{1}{4}-n'-(s'+(2m+k)\alpha + \sum_{i=1}^t K_i\gamma_i + \sum_{i=1}^r \eta_{G_i,g_i}\alpha_i); \eta_1, \eta_2), \\ \vdots \\ (-\frac{1}{2}-(s'+t'+(2m+k)(\alpha + \beta) + \sum_{i=1}^t K_i(\gamma_i + \mu_i) + \sum_{i=1}^r \eta_{G_i,g_i}(\alpha_i + \beta_i)); \epsilon_1 + \eta_1, \epsilon_2 + \eta_2), \\ \vdots \\ (\frac{1}{4}-\frac{1}{2}(s'+t'+(2m+k)(\alpha + \beta) + \sum_{i=1}^t K_i(\gamma_i + \mu_i) + \sum_{i=1}^r \eta_{G_i,g_i}(\alpha_i + \beta_i)); \frac{\epsilon_1+\eta_1}{2}, \frac{\epsilon_2+\eta_2}{2}), \\ \vdots \\ (\frac{1}{4}-n'-\frac{1}{2}(s'+t'+(2m+k)(\alpha + \beta) + \sum_{i=1}^t K_i(\gamma_i + \mu_i) + \sum_{i=1}^r \eta_{G_i,g_i}(\alpha_i + \beta_i)); \frac{\epsilon_1+\eta_1}{2}, \frac{\epsilon_2+\eta_2}{2}), \end{array} \right)$$

$$\begin{aligned}
 & \left(\frac{1}{4} - \frac{1}{2}(s'+t'+(2m+k)(\alpha + \beta) + \sum_{i=1}^t K_i(\gamma_i + \mu_i) + \sum_{i=1}^r \eta_{G_i, g_i}(\alpha_i + \beta_i)); \frac{\epsilon_1 + \eta_1}{2}, \frac{\epsilon_2 + \eta_2}{2} \right), \\
 & \left(\frac{1}{4} - n' - \frac{1}{2}(s'+t'+(2m+k)(\alpha + \beta) + \sum_{i=1}^t K_i(\gamma_i + \mu_i) + \sum_{i=1}^r \eta_{G_i, g_i}(\alpha_i + \beta_i)); \frac{\epsilon_1 + \eta_1}{2}, \frac{\epsilon_2 + \eta_2}{2} \right), \\
 & \left(-\frac{1}{4} - \frac{1}{2}(s'+t'+(2m+k)(\alpha + \beta) + \sum_{i=1}^t K_i(\gamma_i + \mu_i) + \sum_{i=1}^r \eta_{G_i, g_i}(\alpha_i + \beta_i)); \frac{\epsilon_1 + \eta_1}{2}, \frac{\epsilon_2 + \eta_2}{2} \right), \\
 & \left(-\frac{1}{4} - n' - \frac{1}{2}(s'+t'+(2m+k)(\alpha + \beta) + \sum_{i=1}^t K_i(\gamma_i + \mu_i) + \sum_{i=1}^r \eta_{G_i, g_i}(\alpha_i + \beta_i)); \frac{\epsilon_1 + \eta_1}{2}, \frac{\epsilon_2 + \eta_2}{2} \right), \\
 & \left. \begin{aligned}
 & \left(\frac{1}{4} - n' - (t'+(2m+k)\beta + \sum_{i=1}^t K_i\mu_i + \sum_{i=1}^r \eta_{G_i, g_i}\beta_i); \epsilon_1, \epsilon_2 \right), A : C \\
 & \quad \quad \quad \cdot \quad \cdot \quad \cdot \\
 & \quad \quad \quad B : D
 \end{aligned} \right) \tag{5.1}
 \end{aligned}$$

under the same notation and conditions that (3.1) with $s = 2$ and $l_i, l'_i, l''_i \rightarrow 1$

7. Conclusion

In this paper we have evaluated a generalized finite integral involving the multivariable Aleph-functions, the Fresnel integral function, a class of polynomials of several variables and the spheroidal function. The integral established in this paper is of very general nature as it contains Multivariable Aleph-function, which is a general function of several variables studied so far. Thus, the integral established in this research work would serve as a key formula from which, upon specializing the parameters, as many as desired results involving the special functions of one and several variables can be obtained.

REFERENCES

- [1] Abramowitz M and Stegun I.A. Handbook of Mathematical Functions ,1964,
- [2] Brychkow Y.A. Handbook of Special Functions. Derivatives. Integrals, Series and Other Formulas. CRC. Press. Taylor and Francis Group. Boca. Raton. London. New York. 2008.
- [3] Rhodes D.R. On the spheroidal functions. J. Res. Nat. Bur. Standards. Sect. B 74(1970), page187-209.
- [4] Sharma C.K. and Ahmad S.S.: On the multivariable I-function. Acta ciencia Indica Math , 1994 vol 20, no2, p 113-116.
- [5] C.K. Sharma and P.L. Mishra : On the I-function of two variables and its properties. Acta Ciencia Indica Math , 1991 Vol 17 page 667-672.
- [6] Sharma K. On the integral representation and applications of the generalized function of two variables , International Journal of Mathematical Engineering and Sciences , Vol 3 , issue1 (2014) , page1-13.
- [7] Srivastava H.M. A multilinear generating function for the Konhauser set of biorthogonal polynomials suggested by Laguerre polynomial, Pacific. J. Math. 177(1985), page183-191.
- [8] Stratton J.A. And Chu L.J. Elliptic and spheroidal wave function J. Math. And Phys. 20 (1941), page 259-309.

Personal adress : 411 Avenue Joseph Raynaud
Le parc Fleuri , Bat B
83140 , Six-Fours les plages
Tel : 06-83-12-49-68
Department : VAR
Country : FRANCE