

Generalized finite integral involving the extension of Zeta-function, a general class of polynomials, the multivariable Aleph-function and the multivariable I-function I

F.Y. AYANT¹

¹ Teacher in High School , France

ABSTRACT

In the present paper we evaluate a generalized finite integral involving the product of a extension of the Hurwitz-lerch Zeta-function, the arctangent function, the multivariable Aleph-function, the multivariable I-function defined by Prasad and general class of polynomials of several variables. The importance of the result established in this paper lies in the fact they involve the Aleph-function of several variables which is sufficiently general in nature and capable to yielding a large of results merely by specializing the parameters their in.

Keywords:Multivariable Aleph-function, general class of polynomial, extension of the Hurwitz-lerch Zeta-function, multivariable H-function, multivariable I-function.

2010 Mathematics Subject Classification. 33C99, 33C60, 44A20

1.Introduction and preliminaries.

The function Aleph of several variables generalize the multivariable I-function recently study by C.K. Sharma and Ahmad [3] , itself is an a generalisation of G and H-functions of multiple variables. The multiple Mellin-Barnes integral occurring in this paper will be referred to as the multivariables Aleph-function throughout our present study and will be defined and represented as follows.

$$\text{We define : } \aleph(z_1, \dots, z_r) = \aleph_{P_i, Q_i, \tau_i; R: P_i^{(1)}, Q_i^{(1)}, \tau_i^{(1)}; R^{(1)}; \dots; P_i^{(r)}, Q_i^{(r)}, \tau_i^{(r)}; R^{(r)}}^{0, N: M_1, N_1, \dots, M_r, N_r} \left(\begin{matrix} z_1 \\ \cdot \\ \cdot \\ \cdot \\ z_r \end{matrix} \right)$$

$$[(a_j; \alpha_j^{(1)}, \dots, \alpha_j^{(r)})_{1, N_1}], [\tau_i(a_{ji}; \alpha_{ji}^{(1)}, \dots, \alpha_{ji}^{(r)})_{N_1+1, P_i^{(1)}}] ;$$

$$\dots, [\tau_i(b_{ji}; \beta_{ji}^{(1)}, \dots, \beta_{ji}^{(r)})_{M_1+1, Q_i^{(1)}}] ;$$

$$[(c_j^{(1)}; \gamma_j^{(1)})_{1, N_1}], [\tau_i^{(1)}(c_{ji}^{(1)}; \gamma_{ji}^{(1)})_{N_1+1, P_i^{(1)}}]; \dots ; [(c_j^{(r)}; \gamma_j^{(r)})_{1, N_r}], [\tau_i^{(r)}(c_{ji}^{(r)}; \gamma_{ji}^{(r)})_{N_r+1, P_i^{(r)}}]$$

$$[(d_j^{(1)}; \delta_j^{(1)})_{1, M_1}], [\tau_i^{(1)}(d_{ji}^{(1)}; \delta_{ji}^{(1)})_{M_1+1, Q_i^{(1)}}]; \dots ; [(d_j^{(r)}; \delta_j^{(r)})_{1, M_r}], [\tau_i^{(r)}(d_{ji}^{(r)}; \delta_{ji}^{(r)})_{M_r+1, Q_i^{(r)}}]$$

$$= \frac{1}{(2\pi\omega)^r} \int_{L_1} \dots \int_{L_r} \psi(s_1, \dots, s_r) \prod_{k=1}^r \theta_k(s_k) y_k^{s_k} ds_1 \dots ds_r \tag{1.1}$$

with $\omega = \sqrt{-1}$

$$\psi(s_1, \dots, s_r) = \frac{\prod_{j=1}^N \Gamma(1 - a_j + \sum_{k=1}^r \alpha_j^{(k)} s_k)}{\sum_{i=1}^R [\tau_i \prod_{j=N+1}^{P_i} \Gamma(a_{ji} - \sum_{k=1}^r \alpha_{ji}^{(k)} s_k) \prod_{j=1}^{Q_i} \Gamma(1 - b_{ji} + \sum_{k=1}^r \beta_{ji}^{(k)} s_k)]} \tag{1.2}$$

$$\text{and } \theta_k(s_k) = \frac{\prod_{j=1}^{M_k} \Gamma(d_j^{(k)} - \delta_j^{(k)} s_k) \prod_{j=1}^{N_k} \Gamma(1 - c_j^{(k)} + \gamma_j^{(k)} s_k)}{\sum_{i^{(k)}=1}^{R^{(k)}} [\tau_{i^{(k)}} \prod_{j=M_k+1}^{Q_{i^{(k)}}} \Gamma(1 - d_{ji^{(k)}}^{(k)} + \delta_{ji^{(k)}}^{(k)} s_k) \prod_{j=N_k+1}^{P_{i^{(k)}}} \Gamma(c_{ji^{(k)}}^{(k)} - \gamma_{ji^{(k)}}^{(k)} s_k)]} \quad (1.3)$$

Suppose, as usual, that the parameters

$$b_j, j = 1, \dots, Q; a_j, j = 1, \dots, P;$$

$$c_{ji^{(k)}}^{(k)}, j = n_k + 1, \dots, P_{i^{(k)}}; c_j^{(k)}, j = 1, \dots, N_k;$$

$$d_{ji^{(k)}}^{(k)}, j = M_k + 1, \dots, Q_{i^{(k)}}; d_j^{(k)}, j = 1, \dots, M_k;$$

with $k = 1 \dots, r, i = 1, \dots, R, i^{(k)} = 1, \dots, R^{(k)}$

are complex numbers, and the α' s, β' s, γ' s and δ' s are assumed to be positive real numbers for standardization purpose such that

$$U_i^{(k)} = \sum_{j=1}^N \alpha_j^{(k)} + \tau_i \sum_{j=N+1}^{P_i} \alpha_{ji}^{(k)} + \sum_{j=1}^{N_k} \gamma_j^{(k)} + \tau_{i^{(k)}} \sum_{j=n_k+1}^{P_{i^{(k)}}} \gamma_{ji^{(k)}}^{(k)} - \tau_i \sum_{j=1}^{Q_i} \beta_{ji}^{(k)} - \sum_{j=1}^{M_k} \delta_j^{(k)} - \tau_{i^{(k)}} \sum_{j=M_k+1}^{Q_{i^{(k)}}} \delta_{ji^{(k)}}^{(k)} \leq 0 \quad (1.4)$$

The real numbers τ_i are positives for $i = 1$ to R , $\tau_{i^{(k)}}$ are positives for $i^{(k)} = 1$ to $R^{(k)}$

The contour L_k is in the s_k -p lane and run from $\sigma - i\infty$ to $\sigma + i\infty$ where σ is a real number with loop, if necessary, ensure that the poles of $\Gamma(d_j^{(k)} - \delta_j^{(k)} s_k)$ with $j = 1$ to m_k are separated from those of $\Gamma(1 - a_j + \sum_{i=1}^r \alpha_j^{(k)} s_k)$ with $j = 1$ to N and $\Gamma(1 - c_j^{(k)} + \gamma_j^{(k)} s_k)$ with $j = 1$ to N_k to the left of the contour L_k . The condition for absolute convergence of multiple Mellin-Barnes type contour (1.9) can be obtained by extension of the corresponding conditions for multivariable H-function given by as :

$$|arg z_k| < \frac{1}{2} A_i^{(k)} \pi, \text{ where}$$

$$A_i^{(k)} = \sum_{j=1}^N \alpha_j^{(k)} - \tau_i \sum_{j=N+1}^{P_i} \alpha_{ji}^{(k)} - \tau_i \sum_{j=1}^{Q_i} \beta_{ji}^{(k)} + \sum_{j=1}^{N_k} \gamma_j^{(k)} - \tau_{i^{(k)}} \sum_{j=N_k+1}^{P_{i^{(k)}}} \gamma_{ji^{(k)}}^{(k)} + \sum_{j=1}^{M_k} \delta_j^{(k)} - \tau_{i^{(k)}} \sum_{j=M_k+1}^{Q_{i^{(k)}}} \delta_{ji^{(k)}}^{(k)} > 0, \text{ with } k = 1, \dots, r, i = 1, \dots, R, i^{(k)} = 1, \dots, R^{(k)} \quad (1.5)$$

The complex numbers z_i are not zero. Throughout this document, we assume the existence and absolute convergence conditions of the multivariable Aleph-function.

We may establish the asymptotic expansion in the following convenient form :

$$\aleph(z_1, \dots, z_r) = O(|z_1|^{\alpha_1}, \dots, |z_r|^{\alpha_r}), \max(|z_1|, \dots, |z_r|) \rightarrow 0$$

$$\aleph(z_1, \dots, z_r) = O(|z_1|^{\beta_1}, \dots, |z_r|^{\beta_r}), \min(|z_1|, \dots, |z_r|) \rightarrow \infty$$

where, with $k = 1, \dots, r : \alpha_k = \min[Re(d_j^{(k)} / \delta_j^{(k)})], j = 1, \dots, M_k$ and

$$\beta_k = \max[Re((c_j^{(k)} - 1) / \gamma_j^{(k)})], j = 1, \dots, N_k$$

Serie representation of Aleph-function of several variables is given by

$$\begin{aligned} \aleph(y_1, \dots, y_r) = & \sum_{G_1, \dots, G_r=0}^{\infty} \sum_{g_1=0}^{M_1} \dots \sum_{g_r=0}^{M_r} \frac{(-)^{G_1+\dots+G_r}}{\delta_{g_1}^{G_1} \dots \delta_{g_r}^{G_r}} \psi(\eta_{G_1, g_1}, \dots, \eta_{G_r, g_r}) \\ & \times \theta_1(\eta_{G_1, g_1}) \dots \theta_r(\eta_{G_r, g_r}) y_1^{-\eta_{G_1, g_1}} \dots y_r^{-\eta_{G_r, g_r}} \end{aligned} \tag{1.6}$$

Where $\psi(\dots), \theta_i(\dots), i = 1, \dots, r$ are given respectively in (1.2), (1.3) and

$$\eta_{G_1, g_1} = \frac{d_{g_1}^{(1)} + G_1}{\delta_{g_1}^{(1)}}, \dots, \eta_{G_r, g_r} = \frac{d_{g_r}^{(r)} + G_r}{\delta_{g_r}^{(r)}}$$

which is valid under the conditions $\delta_{g_i}^{(i)} [d_j^i + p_i] \neq \delta_j^{(i)} [d_{g_i}^i + G_i]$ (1.7)

for $j \neq M_i, M_i = 1, \dots, \eta_{G_i, g_i}; P_i, N_i = 0, 1, 2, \dots, ; y_i \neq 0, i = 1, \dots, r$ (1.8)

In the document , we will note :

$$G(\eta_{G_1, g_1}, \dots, \eta_{G_r, g_r}) = \phi(\eta_{G_1, g_1}, \dots, \eta_{G_r, g_r}) \theta_1(\eta_{G_1, g_1}) \dots \theta_r(\eta_{G_r, g_r}) \tag{1.9}$$

where $\phi(\eta_{G_1, g_1}, \dots, \eta_{G_r, g_r}), \theta_1(\eta_{G_1, g_1}), \dots, \theta_r(\eta_{G_r, g_r})$ are given respectively in (1.2) and (1.3)

We will note the Aleph-function of r variables $\aleph_{u:w}^{0,N:v} \left(\begin{matrix} z_1 \\ \dots \\ z_r \end{matrix} \right)$ (1.10)

The multivariable I-function is defined in term of multiple Mellin-Barnes type integral :

$$I(z_1, z_2, \dots, z_s) = I_{p_2, q_2, p_3, q_3; \dots; p_s, q_s; p', q'; \dots; p^{(s)}, q^{(s)}}^{0, n_2; 0, n_3; \dots; 0, n_r; m', n'; \dots; m^{(s)}, n^{(s)}} \left(\begin{matrix} z_1 \\ \cdot \\ \cdot \\ \cdot \\ z_s \end{matrix} \middle| \begin{matrix} (a_{2j}; \alpha'_{2j}, \alpha''_{2j})_{1, p_2}; \dots; \\ (b_{2j}; \beta'_{2j}, \beta''_{2j})_{1, q_2}; \dots; \end{matrix} \right)$$

$$\left(\begin{matrix} (a_{sj}; \alpha'_{sj}, \dots, \alpha^{(s)}_{sj})_{1, p_s} : (a'_j, \alpha'_j)_{1, p'}; \dots; (a_j^{(s)}, \alpha_j^{(s)})_{1, p^{(s)}} \\ (b_{sj}; \beta'_{sj}, \dots, \beta^{(s)}_{sj})_{1, q_s} : (b'_j, \beta'_j)_{1, q'}; \dots; (b_j^{(s)}, \beta_j^{(s)})_{1, q^{(s)}} \end{matrix} \right) \tag{1.11}$$

$$= \frac{1}{(2\pi\omega)^s} \int_{L_1} \cdots \int_{L_s} \xi(t_1, \dots, t_s) \prod_{i=1}^s \phi_i(t_i) z_i^{t_i} dt_1 \cdots dt_s \tag{1.12}$$

The defined integral of the above function, the existence and convergence conditions, see Y,N Prasad [2]. Throughout the present document, we assume that the existence and convergence conditions of the multivariable I-function.

The condition for absolute convergence of multiple Mellin-Barnes type contour (1.9) can be obtained by extension of the corresponding conditions for multivariable H-function given by as :

$$|argz_k| < \frac{1}{2} \Omega_i^{(k)} \pi, \text{ where}$$

$$\Omega_i^{(k)} = \sum_{k=1}^{n^{(i)}} \alpha_k^{(i)} - \sum_{k=n^{(i)}+1}^{p^{(i)}} \alpha_k^{(i)} + \sum_{k=1}^{m^{(i)}} \beta_k^{(i)} - \sum_{k=m^{(i)}+1}^{q^{(i)}} \beta_k^{(i)} + \left(\sum_{k=1}^{n_2} \alpha_{2k}^{(i)} - \sum_{k=n_2+1}^{p_2} \alpha_{2k}^{(i)} \right) + \cdots +$$

$$\left(\sum_{k=1}^{n_s} \alpha_{sk}^{(i)} - \sum_{k=n_s+1}^{p_s} \alpha_{sk}^{(i)} \right) - \left(\sum_{k=1}^{q_2} \beta_{2k}^{(i)} + \sum_{k=1}^{q_3} \beta_{3k}^{(i)} + \cdots + \sum_{k=1}^{q_s} \beta_{sk}^{(i)} \right) \tag{1.13}$$

where $i = 1, \dots, s$

The complex numbers z_i are not zero. Throughout this document, we assume the existence and absolute convergence conditions of the multivariable I-function.

We may establish the the asymptotic expansion in the following convenient form :

$$I(z_1, \dots, z_s) = O(|z_1|^{\alpha'_1}, \dots, |z_s|^{\alpha'_s}), \max(|z_1|, \dots, |z_s|) \rightarrow 0$$

$$I(z_1, \dots, z_s) = O(|z_1|^{\beta'_1}, \dots, |z_s|^{\beta'_s}), \min(|z_1|, \dots, |z_s|) \rightarrow \infty$$

where, with $k = 1, \dots, s : \alpha'_k = \min[Re(b_j^{(k)})/\beta_j^{(k)}], j = 1, \dots, m_k$ and

$$\beta'_k = \max[Re((a_j^{(k)} - 1)/\alpha_j^{(k)}), j = 1, \dots, n_k$$

We will use these following notations in this paper :

$$U = p_2, q_2; p_3, q_3; \cdots; p_{s-1}, q_{s-1}; V = 0, n_2; 0, n_3; \cdots; 0, n_{s-1} \tag{1.14}$$

$$W = (p', q'); \cdots; (p^{(s)}, q^{(s)}); X = (m', n'); \cdots; (m^{(s)}, n^{(s)}) \tag{1.15}$$

$$A = (a_{2k}, \alpha'_{2k}, \alpha''_{2k}); \cdots; (a_{(s-1)k}, \alpha'_{(s-1)k}, \alpha''_{(s-1)k}, \cdots, \alpha^{(s-1)}_{(s-1)k}) \tag{1.16}$$

$$B = (b_{2k}, \beta'_{2k}, \beta''_{2k}); \cdots; (b_{(s-1)k}, \beta'_{(s-1)k}, \beta''_{(s-1)k}, \cdots, \beta^{(s-1)}_{(s-1)k}) \tag{1.17}$$

$$\mathfrak{A} = (a_{sk}; \alpha'_{sk}, \alpha''_{sk}, \cdots, \alpha^s_{sk}) : \mathfrak{B} = (b_{sk}; \beta'_{sk}, \beta''_{sk}, \cdots, \beta^s_{sk}) \tag{1.18}$$

$$A' = (a'_k, \alpha'_k)_{1,p'}; \dots; (a_k^{(s)}, \alpha_k^{(s)})_{1,p^{(s)}}; B' = (b'_k, \beta'_k)_{1,q'}; \dots; (b_k^{(s)}, \beta_k^{(s)})_{1,q^{(s)}} \tag{1.19}$$

The multivariable I-function write :

$$I(z_1, \dots, z_s) = I_{U:p_s, q_s; W}^{V; 0, n_s; X} \left(\begin{matrix} z_1 \\ \cdot \\ \cdot \\ z_s \end{matrix} \middle| \begin{matrix} A; \mathfrak{A}; A' \\ B; \mathfrak{B}; B' \end{matrix} \right) \tag{1.20}$$

The generalized polynomials defined by Srivastava [4], is given in the following manner :

$$S_{N'_1, \dots, N'_t}^{M'_1, \dots, M'_t} [y_1, \dots, y_t] = \sum_{K_1=0}^{[N'_1/M'_1]} \dots \sum_{K_t=0}^{[N'_t/M'_t]} \frac{(-N'_1)_{M'_1 K_1}}{K_1!} \dots \frac{(-N'_t)_{M'_t K_t}}{K_t!} A[N'_1, K_1; \dots; N'_t, K_t] y_1^{K_1} \dots y_t^{K_t} \tag{1.21}$$

Where M'_1, \dots, M'_s are arbitrary positive integers and the coefficients $A[N'_1, K_1; \dots; N'_t, K_t]$ are arbitrary constants, real or complex. In the present paper, we use the following notation

$$a_1 = \frac{(-N'_1)_{M'_1 K_1}}{K_1!} \dots \frac{(-N'_t)_{M'_t K_t}}{K_t!} A[N'_1, K_1; \dots; N'_t, K_t] \tag{1.22}$$

2. Extension of the Hurwitz-Lerch Zeta function

The extension of the Hurwitz-Lerch Zeta function $\phi(z, S, a)$ is introduced by Srivastava et al ([6], eq.(6.2), page 503) as follows :

$$\phi_{(\lambda_1, \dots, \lambda_p, \mu_1, \dots, \mu_q)}^{(\rho_1, \dots, \rho_p, \sigma_1, \dots, \sigma_q)}(z; S, a) = \sum_{n=0}^{\infty} \frac{\prod_{j=1}^p (\lambda_j)_{n\rho_j}}{(a+n)^S \prod_{j=1}^q (\mu_j)_{n\sigma_j}} \times \frac{z^n}{n!} \tag{2.1}$$

with : $p, q \in \mathbb{N}_0, \lambda_j \in \mathbb{C} (j = 1, \dots, p), a, \mu_j \in \mathbb{C} \setminus \mathbb{Z}_0^* (j = 1, \dots, q), \rho_j, \sigma_k \in \mathbb{R}^+$

$$(j = 1, \dots, p; k = 1, \dots, q)$$

where $\Delta > -1$ when $S, z \in \mathbb{C}; \Delta = -1$ and $s \in \mathbb{C}$, when $|z| < \nabla^*$, $\Delta = -1$ and $Re(\chi) > \frac{1}{2}$ when $|z| = \nabla^*$

$$\nabla^* = \prod_{j=1}^p \rho_j^{\rho_j} \prod_{j=1}^q \sigma_j^{\sigma_j}; \Delta = \sum_{j=1}^q \sigma_j - \sum_{j=1}^p \rho_j; \chi = S + \sum_{j=1}^q \mu_j - \sum_{j=1}^p \lambda_j + \frac{p-q}{2}$$

We denote these conditions the conditions (f).

3. Required integral

We have the following integral, see Brychkow ([1], 4.1.5, 33 page 136).

$$\int_0^a x^{s-1}(a-x)^{t-1} \arctan(b\sqrt{x(a-x)}) dx = a^{s+t} b B\left(s + \frac{1}{2}, t + \frac{1}{2}\right) \times {}_4F_3\left(\begin{matrix} \frac{1}{2}, 1, s + \frac{1}{2}, t + \frac{1}{2} \\ \frac{3}{2}, \frac{s+t}{2} + 1, \frac{s+t+1}{2} \end{matrix}; -\frac{(ab)^2}{4}\right) \tag{3.1}$$

where $a > 0, Re(s) > -\frac{1}{2}, Re(t) > -\frac{1}{2}, |arg(4 + a^2b^2)| < \pi$

4. Main integral

Let $b_n = \frac{\prod_{j=1}^p (\lambda_j)_{n\rho_j}}{(a+n)^S \prod_{j=1}^q (\mu_j)_{n\sigma_j}}$ and $X_{c,d} = x^c(a-x)^d$, we have the following integral,

$$\int_0^a x^{s'-1}(a-x)^{t'-1} \arctan(b\sqrt{x(a-x)}) \phi_{(\lambda_1, \dots, \lambda_p, \sigma_1, \dots, \sigma_q)}^{(\rho_1, \dots, \rho_p, \sigma_1, \dots, \sigma_q)}(zX_{\alpha, \beta}; S, a) S_{N_1, \dots, N_t}^{M'_1, \dots, M'_t} \begin{pmatrix} y_1 X_{\gamma_1, \mu_1} \\ \dots \\ y_t X_{\gamma_t, \mu_t} \end{pmatrix}$$

$$N_{u:w}^{0, N:v} \begin{pmatrix} z_1 X_{\alpha_1, \beta_1} \\ \dots \\ z_r X_{\alpha_r, \beta_r} \end{pmatrix} I_{U:p_s, q_s; W}^{V; 0, n_s; X} \begin{pmatrix} Z_1 X_{\eta_1, \epsilon_1} \\ \dots \\ Z_s X_{\eta_s, \epsilon_s} \end{pmatrix} dx = a^{s'+t'} b \sum_{K_1=0}^{[N'_1/M'_1]} \dots \sum_{K_t=0}^{[N'_t/M'_t]} \sum_{n=0}^{\infty} \sum_{n'=0}^{\infty}$$

$$\sum_{G_1, \dots, G_r=0}^{\infty} \sum_{g_1=0}^{M_1} \dots \sum_{g_r=0}^{M_r} \frac{(-)^{G_1 + \dots + G_r}}{\delta_{g_1} G_1! \dots \delta_{g_r} G_r!} G(\eta_{G_1, g_1}, \dots, \eta_{G_r, g_r}) a_1 \frac{b_n z^n (\frac{1}{2})_{n'}}{n!} \frac{(-a^2 b^2)^{n'}}{4^{n'} (\frac{3}{2})_{n'}} z_1^{\eta_{G_1, g_1}} \dots z_r^{\eta_{G_r, g_r}}$$

$$y_1^{K_1} \dots y_t^{K_t} a^{n(\alpha + \beta) + \sum_{i=1}^t K_i(\gamma_i + \mu_i) + \sum_{i=1}^r \eta_{G_i, g_i}(\alpha_i + \beta_i)} I_{U:p_s+4, q_s+3; W}^{V; 0, n_s+4; X} \begin{pmatrix} Z_1 a^{\eta_1 + \epsilon_1} & | & \mathbf{A}; \mathfrak{A}, \\ \dots & & \\ \dots & & \\ Z_s a^{\eta_s + \epsilon_s} & | & \mathbf{B}; \mathfrak{B}, \end{pmatrix}$$

$$(\frac{1}{2} - n' - (s' + n\alpha + \sum_{i=1}^t K_i \gamma_i + \sum_{i=1}^r \eta_{G_i, g_i} \alpha_i); \eta_1, \dots, \eta_s),$$

$$(-(s'+t'+n(\alpha + \beta) + \sum_{i=1}^t K_i(\gamma_i + \mu_i) + \sum_{i=1}^r \eta_{G_i, g_i}(\alpha_i + \beta_i)); \epsilon_1 + \eta_1, \dots, \epsilon_s + \eta_s),$$

$$(-\frac{1}{2}(s'+t'+n(\alpha + \beta) + \sum_{i=1}^t K_i(\gamma_i + \mu_i) + \sum_{i=1}^r \eta_{G_i, g_i}(\alpha_i + \beta_i)); \frac{\epsilon_1 + \eta_1}{2}, \dots, \frac{\epsilon_s + \eta_s}{2}),$$

$$(-n' - \frac{1}{2}(s'+t'+n(\alpha + \beta) + \sum_{i=1}^t K_i(\gamma_i + \mu_i) + \sum_{i=1}^r \eta_{G_i, g_i}(\alpha_i + \beta_i)); \frac{\epsilon_1 + \eta_1}{2}, \dots, \frac{\epsilon_s + \eta_s}{2}),$$

$$\int_0^a x^{s'-1} (a-x)^{t'-1} \arctan(b\sqrt{x(a-x)}) \phi_{(\lambda_1, \dots, \lambda_p, \sigma_1, \dots, \sigma_q)}^{(\rho_1, \dots, \rho_p, \sigma_1, \dots, \sigma_q)}(zX_{\alpha, \beta}; S, a) S_{N_1, \dots, N_t}^{M'_1, \dots, M'_t} \left(\begin{matrix} y_1 X_{\gamma_1, \mu_1} \\ \dots \\ y_t X_{\gamma_t, \mu_t} \end{matrix} \right)$$

$$\delta_{u:w}^{0, N:v} \left(\begin{matrix} z_1 X_{\alpha_1, \beta_1} \\ \dots \\ z_r X_{\alpha_r, \beta_r} \end{matrix} \right) H_{p_s, q_s; W}^{0, n_s; X} \left(\begin{matrix} Z_1 X_{\eta_1, \epsilon_1} \\ \dots \\ Z_s X_{\eta_s, \epsilon_s} \end{matrix} \right) dx = a^{s'+t'} b \sum_{K_1=0}^{[N'_1/M'_1]} \dots \sum_{K_t=0}^{[N'_t/M'_t]} \sum_{n=0}^{\infty} \sum_{n'=0}^{\infty}$$

$$\sum_{G_1, \dots, G_r=0}^{\infty} \sum_{g_1=0}^{M_1} \dots \sum_{g_r=0}^{M_r} \frac{(-)^{G_1+\dots+G_r}}{\delta_{g_1} G_1! \dots \delta_{g_r} G_r!} G(\eta_{G_1, g_1}, \dots, \eta_{G_r, g_r}) a_1 \frac{b_n z^n (\frac{1}{2})_{n'}}{n!} \frac{(-a^2 b^2)^{n'}}{4^{n'} (\frac{3}{2})_{n'}} z_1^{\eta_{G_1, g_1}} \dots z_r^{\eta_{G_r, g_r}}$$

$$y_1^{K_1} \dots y_t^{K_t} a^{n(\alpha+\beta)+\sum_{i=1}^t K_i(\gamma_i+\mu_i)+\sum_{i=1}^r \eta_{G_i, g_i}(\alpha_i+\beta_i)} H_{p_s+4, q_s+3; W}^{0, n_s+4; X} \left(\begin{matrix} Z_1 a^{\eta_1+\epsilon_1} \\ \dots \\ Z_s a^{\eta_s+\epsilon_s} \end{matrix} \middle| \begin{matrix} \mathfrak{A}, \\ \mathfrak{B}, \end{matrix} \right)$$

$$\left(\frac{1}{2} - n' - (s' + n\alpha + \sum_{i=1}^t K_i \gamma_i + \sum_{i=1}^r \eta_{G_i, g_i} \alpha_i); \eta_1, \dots, \eta_s \right),$$

$$(- (s'+t'+n(\alpha + \beta) + \sum_{i=1}^t K_i(\gamma_i + \mu_i) + \sum_{i=1}^r \eta_{G_i, g_i}(\alpha_i + \beta_i)); \epsilon_1 + \eta_1, \dots, \epsilon_s + \eta_s),$$

$$(-\frac{1}{2}(s'+t'+n(\alpha + \beta) + \sum_{i=1}^t K_i(\gamma_i + \mu_i) + \sum_{i=1}^r \eta_{G_i, g_i}(\alpha_i + \beta_i)); \frac{\epsilon_1+\eta_1}{2}, \dots, \frac{\epsilon_s+\eta_s}{2}),$$

$$(-n'-\frac{1}{2}(s'+t'+n(\alpha + \beta) + \sum_{i=1}^t K_i(\gamma_i + \mu_i) + \sum_{i=1}^r \eta_{G_i, g_i}(\alpha_i + \beta_i)); \frac{\epsilon_1+\eta_1}{2}, \dots, \frac{\epsilon_s+\eta_s}{2}),$$

$$(\frac{1}{2}-\frac{1}{2}(s'+t'+n(\alpha + \beta) + \sum_{i=1}^t K_i(\gamma_i + \mu_i) + \sum_{i=1}^r \eta_{G_i, g_i}(\alpha_i + \beta_i)); \frac{\epsilon_1+\eta_1}{2}, \dots, \frac{\epsilon_s+\eta_s}{2}),$$

$$(\frac{1}{2}-n'-\frac{1}{2}(s'+t'+n(\alpha + \beta) + \sum_{i=1}^t K_i(\gamma_i + \mu_i) + \sum_{i=1}^r \eta_{G_i, g_i}(\alpha_i + \beta_i)); \frac{\epsilon_1+\eta_1}{2}, \dots, \frac{\epsilon_s+\eta_s}{2}),$$

$$\left(\frac{1}{2}-n'-(t'+n\beta + \sum_{i=1}^t K_i \mu_i + \sum_{i=1}^r \eta_{G_i, g_i} \beta_i); \epsilon_1, \dots, \epsilon_s, A' \right) \left(\begin{matrix} \dots \\ B' \end{matrix} \right) \tag{5.1}$$

under the same notations and conditions that (4.1) with $U = V = A = B = 0$

6. Conclusion

In this paper we have evaluated a generalized finite integral involving the multivariable Aleph-function, a class of polynomials of several variables a extension of the Hurwitz-Lerch Zeta-function and the multivariable I-function

defined by Prasad. The integral established in this paper is of very general nature as it contains Multivariable Aleph-function, which is a general function of several variables studied so far. Thus, the integral established in this research work would serve as a key formula from which, upon specializing the parameters, as many as desired results involving the special functions of one and several variables can be obtained.

REFERENCES

- [1] Brychkow Y.A. **Handbook of Special Functions. Derivatives, Integrals, Series and Other Formulas.** CRC. Press. Taykor and Francis Group. Boca. Raton. London. New York. 2008.
- [2] Y.N. Prasad , Multivariable I-function , Vijnana Parishad Anusandhan Patrika 29 (1986) , page 231-237.
- [3] Sharma C.K.and Ahmad S.S.: On the multivariable I-function. Acta ciencia Indica Math , 1994 vol 20,no2, p 113-116.
- [4] Srivastava H.M. A multilinear generating function for the Konhauser set of biorthogonal polynomials suggested by Laguerre polynomial, Pacific. J. Math. 177(1985), page183-191.
- [5] H.M. Srivastava And R.Panda. Some expansion theorems and generating relations for the H-function of several complex variables. Comment. Math. Univ. St. Paul. 24(1975), p.119-137.
- [6] H.M. Srivastava, R.K. Saxena, T.K. Pogány, R. Saxena, Integral and computational representations of the extended Hurwitz–Lerch zeta function, Integr.Transf. Spec. Funct. 22 (2011) 487–506
- [7] Whittaker E.T and Watson G.N. A course of modern analysis. Naw York. Mac Millan 1943.

Personal adress : 411 Avenue Joseph Raynaud

Le parc Fleuri , Bat B

83140 , Six-Fours les plages

Tel : 06-83-12-49-68

Department : VAR

Country : FRANCE