

Finite integral involving the generalized multiple Zeta-function, a general class of polynomials and multivariable Aleph-functions dilogarithm function and the multivariable I-function

F.Y. AYANT¹

¹ Teacher in High School , France

ABSTRACT

In the present paper we evaluate a generalized finite integral involving the product of generalized multiple zeta-function, the dilogarithm function, the multivariable Aleph-function, the multivariable I-function defined by Prasad and general class of polynomials of several variables. The importance of the result established in this paper lies in the fact they involve the Aleph-function of several variables which is sufficiently general in nature and capable to yielding a large of results merely by specializing the parameters their in.

Keywords: Multivariable Aleph-function, general class of polynomial, multiple Zeta-function, dilogarithm function, multivariable I-function, multivariable H-function

2010 Mathematics Subject Classification. 33C99, 33C60, 44A20

1.Introduction and preliminaries.

The function Aleph of several variables generalize the multivariable I-function recently study by C.K. Sharma and Ahmad [5], itself is an a generalisation of G and H-functions of multiple variables. The multiple Mellin-Barnes integral occuring in this paper will be referred to as the multivariables Aleph-function throughout our present study and will be defined and represented as follows.

$$\text{We define : } \aleph(z_1, \dots, z_r) = \aleph_{P_i, Q_i, \tau_i; R: P_i(1), Q_i(1), \tau_i(1); R^{(1)}; \dots; P_i(r), Q_i(r), \tau_i(r); R^{(r)}}^{0, N: M_1, N_1, \dots, M_r, N_r} \left(\begin{matrix} Z_1 \\ \cdot \\ \cdot \\ \cdot \\ Z_r \end{matrix} \right)$$

$$[(a_j; \alpha_j^{(1)}, \dots, \alpha_j^{(r)})_{1, N}] , [\tau_i(a_{ji}; \alpha_{ji}^{(1)}, \dots, \alpha_{ji}^{(r)})_{N+1, P_i}] :$$

$$\dots\dots\dots, [\tau_i(b_{ji}; \beta_{ji}^{(1)}, \dots, \beta_{ji}^{(r)})_{M+1, Q_i}] :$$

$$[(c_j^{(1)}; \gamma_j^{(1)})_{1, N_1}], [\tau_{i(1)}(c_{ji(1)}^{(1)}; \gamma_{ji(1)}^{(1)})_{N_1+1, P_i^{(1)}}]; \dots; [(c_j^{(r)}; \gamma_j^{(r)})_{1, N_r}], [\tau_{i(r)}(c_{ji(r)}^{(r)}; \gamma_{ji(r)}^{(r)})_{N_r+1, P_i^{(r)}}]$$

$$[(d_j^{(1)}; \delta_j^{(1)})_{1, M_1}], [\tau_{i(1)}(d_{ji(1)}^{(1)}; \delta_{ji(1)}^{(1)})_{M_1+1, Q_i^{(1)}}]; \dots; [(d_j^{(r)}; \delta_j^{(r)})_{1, M_r}], [\tau_{i(r)}(d_{ji(r)}^{(r)}; \delta_{ji(r)}^{(r)})_{M_r+1, Q_i^{(r)}}]$$

$$= \frac{1}{(2\pi\omega)^r} \int_{L_1} \dots \int_{L_r} \psi(s_1, \dots, s_r) \prod_{k=1}^r \theta_k(s_k) y_k^{s_k} ds_1 \dots ds_r \quad (1.1)$$

with $\omega = \sqrt{-1}$

$$\psi(s_1, \dots, s_r) = \frac{\prod_{j=1}^N \Gamma(1 - a_j + \sum_{k=1}^r \alpha_j^{(k)} s_k)}{\sum_{i=1}^R [\tau_i \prod_{j=N+1}^{P_i} \Gamma(a_{ji} - \sum_{k=1}^r \alpha_{ji}^{(k)} s_k) \prod_{j=1}^{Q_i} \Gamma(1 - b_{ji} + \sum_{k=1}^r \beta_{ji}^{(k)} s_k)]} \quad (1.2)$$

$$\text{and } \theta_k(s_k) = \frac{\prod_{j=1}^{M_k} \Gamma(d_j^{(k)} - \delta_j^{(k)} s_k) \prod_{j=1}^{N_k} \Gamma(1 - c_j^{(k)} + \gamma_j^{(k)} s_k)}{\sum_{i^{(k)}=1}^{R^{(k)}} [\tau_{i^{(k)}} \prod_{j=M_k+1}^{Q_{i^{(k)}}} \Gamma(1 - d_{ji^{(k)}}^{(k)} + \delta_{ji^{(k)}}^{(k)} s_k) \prod_{j=N_k+1}^{P_{i^{(k)}}} \Gamma(c_{ji^{(k)}}^{(k)} - \gamma_{ji^{(k)}}^{(k)} s_k)]} \quad (1.3)$$

Suppose, as usual, that the parameters

$$b_j, j = 1, \dots, Q; a_j, j = 1, \dots, P;$$

$$c_{ji^{(k)}}^{(k)}, j = n_k + 1, \dots, P_{i^{(k)}}; c_j^{(k)}, j = 1, \dots, N_k;$$

$$d_{ji^{(k)}}^{(k)}, j = M_k + 1, \dots, Q_{i^{(k)}}; d_j^{(k)}, j = 1, \dots, M_k;$$

with $k = 1, \dots, r, i = 1, \dots, R, i^{(k)} = 1, \dots, R^{(k)}$

are complex numbers, and the $\alpha's, \beta's, \gamma's$ and $\delta's$ are assumed to be positive real numbers for standardization purpose such that

$$U_i^{(k)} = \sum_{j=1}^N \alpha_j^{(k)} + \tau_i \sum_{j=N+1}^{P_i} \alpha_{ji}^{(k)} + \sum_{j=1}^{N_k} \gamma_j^{(k)} + \tau_{i^{(k)}} \sum_{j=n_k+1}^{P_{i^{(k)}}} \gamma_{ji^{(k)}}^{(k)} - \tau_i \sum_{j=1}^{Q_i} \beta_{ji}^{(k)} - \sum_{j=1}^{M_k} \delta_j^{(k)} - \tau_{i^{(k)}} \sum_{j=M_k+1}^{Q_{i^{(k)}}} \delta_{ji^{(k)}}^{(k)} \leq 0 \quad (1.4)$$

The real numbers τ_i are positives for $i = 1$ to R , $\tau_{i^{(k)}}$ are positives for $i^{(k)} = 1$ to $R^{(k)}$

The contour L_k is in the s_k -p lane and run from $\sigma - i\infty$ to $\sigma + i\infty$ where σ is a real number with loop, if necessary, ensure that the poles of $\Gamma(d_j^{(k)} - \delta_j^{(k)} s_k)$ with $j = 1$ to m_k are separated from those of $\Gamma(1 - a_j + \sum_{i=1}^r \alpha_j^{(k)} s_k)$ with $j = 1$ to N and $\Gamma(1 - c_j^{(k)} + \gamma_j^{(k)} s_k)$ with $j = 1$ to N_k to the left of the contour L_k . The condition for absolute convergence of multiple Mellin-Barnes type contour (1.9) can be obtained by extension of the corresponding conditions for multivariable H-function given by as :

$$|arg z_k| < \frac{1}{2} A_i^{(k)} \pi, \text{ where}$$

$$A_i^{(k)} = \sum_{j=1}^N \alpha_j^{(k)} - \tau_i \sum_{j=N+1}^{P_i} \alpha_{ji}^{(k)} - \tau_i \sum_{j=1}^{Q_i} \beta_{ji}^{(k)} + \sum_{j=1}^{N_k} \gamma_j^{(k)} - \tau_{i^{(k)}} \sum_{j=N_k+1}^{P_{i^{(k)}}} \gamma_{ji^{(k)}}^{(k)} + \sum_{j=1}^{M_k} \delta_j^{(k)} - \tau_{i^{(k)}} \sum_{j=M_k+1}^{Q_{i^{(k)}}} \delta_{ji^{(k)}}^{(k)} > 0, \text{ with } k = 1, \dots, r, i = 1, \dots, R, i^{(k)} = 1, \dots, R^{(k)} \quad (1.5)$$

The complex numbers z_i are not zero. Throughout this document, we assume the existence and absolute convergence conditions of the multivariable Aleph-function.

We may establish the asymptotic expansion in the following convenient form :

$$\aleph(z_1, \dots, z_r) = O(|z_1|^{\alpha_1}, \dots, |z_r|^{\alpha_r}), \max(|z_1|, \dots, |z_r|) \rightarrow 0$$

$$\aleph(z_1, \dots, z_r) = O(|z_1|^{\beta_1}, \dots, |z_r|^{\beta_r}), \min(|z_1|, \dots, |z_r|) \rightarrow \infty$$

where, with $k = 1, \dots, r : \alpha_k = \min[Re(d_j^{(k)} / \delta_j^{(k)})], j = 1, \dots, M_k$ and

$$\beta_k = \max[Re((c_j^{(k)} - 1) / \gamma_j^{(k)})], j = 1, \dots, N_k$$

Serie representation of Aleph-function of several variables is given by

$$\aleph(y_1, \dots, y_r) = \sum_{G_1, \dots, G_r=0}^{\infty} \sum_{g_1=0}^{M_1} \dots \sum_{g_r=0}^{M_r} \frac{(-)^{G_1+\dots+G_r}}{\delta_{g_1} G_1! \dots \delta_{g_r} G_r!} \psi(\eta_{G_1, g_1}, \dots, \eta_{G_r, g_r}) \\ \times \theta_1(\eta_{G_1, g_1}) \dots \theta_r(\eta_{G_r, g_r}) y_1^{-\eta_{G_1, g_1}} \dots y_r^{-\eta_{G_r, g_r}} \quad (1.6)$$

Where $\psi(\cdot, \dots, \cdot), \theta_i(\cdot), i = 1, \dots, r$ are given respectively in (1.2), (1.3) and

$$\eta_{G_1, g_1} = \frac{d_{g_1}^{(1)} + G_1}{\delta_{g_1}^{(1)}}, \dots, \eta_{G_r, g_r} = \frac{d_{g_r}^{(r)} + G_r}{\delta_{g_r}^{(r)}}$$

$$\text{which is valid under the conditions } \delta_{g_i}^{(i)}[d_j^i + p_i] \neq \delta_j^{(i)}[d_{g_i}^i + G_i] \quad (1.7)$$

$$\text{for } j \neq M_i, M_i = 1, \dots, \eta_{G_i, g_i}; P_i, N_i = 0, 1, 2, \dots; y_i \neq 0, i = 1, \dots, r \quad (1.8)$$

In the document, we will note :

$$G(\eta_{G_1, g_1}, \dots, \eta_{G_r, g_r}) = \phi(\eta_{G_1, g_1}, \dots, \eta_{G_r, g_r}) \theta_1(\eta_{G_1, g_1}) \dots \theta_r(\eta_{G_r, g_r}) \quad (1.9)$$

where $\phi(\eta_{G_1, g_1}, \dots, \eta_{G_r, g_r}), \theta_1(\eta_{G_1, g_1}), \dots, \theta_r(\eta_{G_r, g_r})$ are given respectively in (1.2) and (1.3)

$$\text{We will note the Aleph-function of r variables } \aleph_{u:w}^{0,N:v} \left(\begin{matrix} z_1 \\ \cdot \\ \cdot \\ z_r \end{matrix} \right) \quad (1.10)$$

The multivariable I-function is defined in term of multiple Mellin-Barnes type integral :

$$I(z_1, z_2, \dots, z_s) = I_{p_2, q_2, p_3, q_3; \dots; p_s, q_s: p', q'; \dots; p^{(s)}, q^{(s)}}^{0, n_2; 0, n_3; \dots; 0, n_r: m', n'; \dots; m^{(s)}, n^{(s)}} \left(\begin{matrix} z_1 \\ \cdot \\ \cdot \\ z_s \end{matrix} \middle| \begin{matrix} (a_{2j}; \alpha'_{2j}, \alpha''_{2j})_{1, p_2}; \dots; \\ (b_{2j}; \beta'_{2j}, \beta''_{2j})_{1, q_2}; \dots; \end{matrix} \right) \\ \left(\begin{matrix} (a_{sj}; \alpha'_{sj}, \dots, \alpha_{sj}^{(s)})_{1, p_s}; (a'_j, \alpha'_j)_{1, p'}; \dots; (a_j^{(s)}, \alpha_j^{(s)})_{1, p^{(s)}} \\ (b_{sj}; \beta'_{sj}, \dots, \beta_{sj}^{(s)})_{1, q_s}; (b'_j, \beta'_j)_{1, q'}; \dots; (b_j^{(s)}, \beta_j^{(s)})_{1, q^{(s)}} \end{matrix} \right) \quad (1.11)$$

$$= \frac{1}{(2\pi\omega)^s} \int_{L_1} \cdots \int_{L_s} \xi(t_1, \dots, t_s) \prod_{i=1}^s \phi_i(t_i) z_i^{t_i} dt_1 \cdots dt_s \quad (1.12)$$

The defined integral of the above function, the existence and convergence conditions, see Y,N Prasad [4]. Throughout the present document, we assume that the existence and convergence conditions of the multivariable I-function.

The condition for absolute convergence of multiple Mellin-Barnes type contour (1.9) can be obtained by extension of the corresponding conditions for multivariable H-function given by as :

$$|arg z_k| < \frac{1}{2} \Omega_i^{(k)} \pi, \text{ where}$$

$$\Omega_i^{(k)} = \sum_{k=1}^{n^{(i)}} \alpha_k^{(i)} - \sum_{k=n^{(i)}+1}^{p^{(i)}} \alpha_k^{(i)} + \sum_{k=1}^{m^{(i)}} \beta_k^{(i)} - \sum_{k=m^{(i)}+1}^{q^{(i)}} \beta_k^{(i)} + \left(\sum_{k=1}^{n_2} \alpha_{2k}^{(i)} - \sum_{k=n_2+1}^{p_2} \alpha_{2k}^{(i)} \right) + \cdots +$$

$$\left(\sum_{k=1}^{n_s} \alpha_{sk}^{(i)} - \sum_{k=n_s+1}^{p_s} \alpha_{sk}^{(i)} \right) - \left(\sum_{k=1}^{q_2} \beta_{2k}^{(i)} + \sum_{k=1}^{q_3} \beta_{3k}^{(i)} + \cdots + \sum_{k=1}^{q_s} \beta_{sk}^{(i)} \right) \quad (1.13)$$

where $i = 1, \dots, s$

The complex numbers z_i are not zero. Throughout this document, we assume the existence and absolute convergence conditions of the multivariable I-function.

We may establish the asymptotic expansion in the following convenient form :

$$I(z_1, \dots, z_s) = O(|z_1|^{\alpha'_1}, \dots, |z_s|^{\alpha'_s}), \max(|z_1|, \dots, |z_s|) \rightarrow 0$$

$$I(z_1, \dots, z_s) = O(|z_1|^{\beta'_1}, \dots, |z_s|^{\beta'_s}), \min(|z_1|, \dots, |z_s|) \rightarrow \infty$$

where, with $k = 1, \dots, s : \alpha'_k = \min[Re(b_j^{(k)})/\beta_j^{(k)}], j = 1, \dots, m_k$ and

$$\beta'_k = \max[Re((a_j^{(k)} - 1)/\alpha_j^{(k)}), j = 1, \dots, n_k]$$

We will use these following notations in this paper :

$$U = p_2, q_2; p_3, q_3; \dots; p_{s-1}, q_{s-1}; V = 0, n_2; 0, n_3; \dots; 0, n_{s-1} \quad (1.14)$$

$$W = (p', q'); \dots; (p^{(s)}, q^{(s)}); X = (m', n'); \dots; (m^{(s)}, n^{(s)}) \quad (1.15)$$

$$A = (a_{2k}, \alpha'_{2k}, \alpha''_{2k}); \dots; (a_{(s-1)k}, \alpha'_{(s-1)k}, \alpha''_{(s-1)k}, \dots, \alpha_{(s-1)k}^{(s-1)}) \quad (1.16)$$

$$B = (b_{2k}, \beta'_{2k}, \beta''_{2k}); \dots; (b_{(s-1)k}, \beta'_{(s-1)k}, \beta''_{(s-1)k}, \dots, \beta_{(s-1)k}^{(s-1)}) \quad (1.17)$$

$$\mathfrak{A} = (a_{sk}; \alpha'_{sk}, \alpha''_{sk}, \dots, \alpha_{sk}^s) : \mathfrak{B} = (b_{sk}; \beta'_{sk}, \beta''_{sk}, \dots, \beta_{sk}^s) \quad (1.18)$$

$$A' = (a'_k, \alpha'_k)_{1,p'}; \dots; (a_k^{(s)}, \alpha_k^{(s)})_{1,p^{(s)}}; B' = (b'_k, \beta'_k)_{1,q'}; \dots; (b_k^{(s)}, \beta_k^{(s)})_{1,q^{(s)}} \quad (1.19)$$

The multivariable I-function write :

$$I(z_1, \dots, z_s) = I_{U; p_s, q_s; W}^{V; 0, n_s; X} \left(\begin{matrix} z_1 \\ \cdot \\ \cdot \\ \cdot \\ z_s \end{matrix} \middle| \begin{matrix} A; \mathfrak{A}; A' \\ \\ B; \mathfrak{B}; B' \end{matrix} \right) \quad (1.20)$$

The generalized polynomials defined by Srivastava [6], is given in the following manner :

$$S_{N'_1, \dots, N'_t}^{M'_1, \dots, M'_t} [y_1, \dots, y_t] = \sum_{K_1=0}^{[N'_1/M'_1]} \dots \sum_{K_t=0}^{[N'_t/M'_t]} \frac{(-N'_1)_{M'_1 K_1}}{K_1!} \dots \frac{(-N'_t)_{M'_t K_t}}{K_t!} A[N'_1, K_1; \dots; N'_t, K_t] y_1^{K_1} \dots y_t^{K_t} \quad (1.21)$$

Where M'_1, \dots, M'_s are arbitrary positive integers and the coefficients $A[N'_1, K_1; \dots; N'_t, K_t]$ are arbitrary constants, real or complex. In the present paper, we use the following notation

$$a_1 = \frac{(-N'_1)_{M'_1 K_1}}{K_1!} \dots \frac{(-N'_t)_{M'_t K_t}}{K_t!} A[N'_1, K_1; \dots; N'_t, K_t] \quad (1.22)$$

2. Generalized multiple Zeta-function and dilogarithm function

Bin Saad et al [2] have defined the generalized multiple Zeta-function $\phi_{\mu, \lambda_1, \dots, \lambda_s}^{(s)}(x_1, \dots, x_s, z, a)$ by

$$\phi_{\mu, \lambda_1, \dots, \lambda_s}^{(s)}(x_1, \dots, x_s, z, a) = \sum_{p_1, \dots, p_s=0}^{\infty} \frac{(\mu)_{p_1+\dots+p_s} x_1^{p_1} \dots x_s^{p_s}}{p_1! \dots p_s! (a + \lambda_1 p_1 + \dots + \lambda_s p_s)^z} \quad (2.1)$$

where $|x_i| < 1, i = 1 \dots, s, s \in \mathbb{Z}^+, \mu \in \mathbb{C} \setminus \{0, -1, -2, \dots\}$ and

$a \in \mathbb{C} \setminus \{-(\lambda_i, p_i), p_i \in \mathbb{N} \cup \{0\}\}, \operatorname{Re}(z) > 0$

The dilogarithm function denoted $Li_2(z)$ is defined by : serie expansion and integral.

$$Li_2(z) = \sum_{k=0}^{\infty} \frac{z^k}{k^2}, \text{ where } |z| < 1 \quad (2.2)$$

$$Li_2(z) = \int_z^0 \frac{(1-t)}{t} dt \quad (2.3)$$

For more details, see Berndt ([1], page 323-326)

3. Required integral

we have the following integral, see Brychkow ([3], 4.2.3, 1 page 180).

$$\int_0^a x^{\mu-1} \ln \left(\frac{\sqrt{a} + \sqrt{a-x}}{\sqrt{x}} \right) Li_2(bx) dx = \frac{\sqrt{\pi} a^{\mu+1} b \Gamma(\mu)}{2\mu(\mu+1)\Gamma(\mu + \frac{3}{2})} \times \left[{}_3F_2 \left(\begin{matrix} 1, \mu+1, \mu+1 \\ \mu + \frac{3}{2}, \mu+2 \end{matrix}; ab \right) - (\mu+1) {}_3F_2 \left(\begin{matrix} 1, 1, \mu+1 \\ \mu + \frac{3}{2}, 2 \end{matrix}; ab \right) - \mu(\mu+1) {}_4F_3 \left(\begin{matrix} 1, 1, 1, \mu+1 \\ \mu + \frac{3}{2}, 2, 2 \end{matrix}; ab \right) \right] \quad (3.1)$$

with $Re(\mu) > -2, |arg(1-ab)| < \pi, a > 0$

4. Main integral

We have the following generalized finite integral :

$$\int_0^1 x^{\mu-1} \ln \left(\frac{\sqrt{a} + \sqrt{a-x}}{\sqrt{x}} \right) Li_2(bx) \phi_{\mu, \lambda_1, \dots, \lambda_s}^{(s)}(t_1 x^{\rho_1}, \dots, t_s x^{\rho_s}, z, a) S_{N_1, \dots, N_t}^{M'_1, \dots, M'_t} \left(\begin{matrix} y_1 x^{\gamma_1} \\ \dots \\ y_t x^{\gamma_t} \end{matrix} \right)$$

$$\mathfrak{N}_{u:w}^{0,N:v} \left(\begin{matrix} z_1 x^{\alpha_1} \\ \dots \\ z_r x^{\alpha_r} \end{matrix} \right) I_{U:p_R, q_R; W}^{V; 0, n_R; X} \left(\begin{matrix} Z_1 x^{\eta_1} \\ \dots \\ Z_R x^{\eta_R} \end{matrix} \right) dx = \frac{\sqrt{\pi} a^{\mu+1} b}{2} \left[\sum_{p_1, \dots, p_s=0}^{\infty} \right]$$

$$\sum_{K_1=0}^{[N_1/M'_1]} \dots \sum_{K_t=0}^{[N_t/M'_t]} \sum_{n'=0}^{\infty} \sum_{G_1, \dots, G_r=0}^{\infty} \sum_{g_1=0}^{M_1} \dots \sum_{g_r=0}^{M_r} \frac{(-)^{G_1+\dots+G_r}}{\delta_{g_1} G_1! \dots \delta_{g_r} G_r!} G(\eta_{G_1, g_1}, \dots, \eta_{G_r, g_r}) a_1$$

$$\frac{(\mu)_{p_1+\dots+p_s}}{p_1! \dots p_s! (a + \lambda_1 p_1 + \dots + \lambda_s p_s) z} t_1^{p_1} \dots t_s^{p_s} z_1^{\eta_{G_1, g_1}} \dots z_r^{\eta_{G_r, g_r}} y_1^{K_1} \dots y_t^{K_t} (ab)^{n'}$$

$$a^{\sum_{i=1}^s \rho_i p_i + \sum_{i=1}^t K_i \gamma_i + \sum_{i=1}^r \eta_{G_i, g_i} \alpha_i} I_{U:p_R+5, q_R+5; W}^{V; 0, n_R+5; X} \left(\begin{matrix} a^{\eta_1} Z_1 \\ \dots \\ a^{\eta_R} Z_R \end{matrix} \right)$$

$$A : (1 - (\mu + \sum_{i=1}^s \rho_i p_i + \sum_{i=1}^t K_i \gamma_i + \sum_{i=1}^r \eta_{G_i, g_i} \alpha_i); \eta_1, \dots, \eta_s),$$

$$\dots$$

$$B : (-1 - (\mu + \sum_{i=1}^s \rho_i p_i + \sum_{i=1}^t K_i \gamma_i + \sum_{i=1}^r \eta_{G_i, g_i} \alpha_i); \eta_1, \dots, \eta_s),$$

$$\begin{aligned}
 & (1-(\mu + \sum_{i=1}^s \rho_i p_i + \sum_{i=1}^t K_i \gamma_i + \sum_{i=1}^\eta G_{i,g_i} \alpha_i); \eta_1, \dots, \eta_s), \\
 & \quad \vdots \\
 & \quad \vdots \\
 & (-\frac{1}{2}\text{-n}'-(\mu + \sum_{i=1}^s \rho_i p_i + \sum_{i=1}^t K_i \gamma_i + \sum_{i=1}^\eta G_{i,g_i} \alpha_i); \eta_1, \dots, \eta_s), \\
 & \quad \vdots \\
 & (-\text{n}'-(\mu + \sum_{i=1}^s \rho_i p_i + \sum_{i=1}^t K_i \gamma_i + \sum_{i=1}^\eta G_{i,g_i} \alpha_i); \eta_1, \dots, \eta_s), \\
 & \quad \vdots \\
 & \quad \vdots \\
 & (-(\mu + \sum_{i=1}^s \rho_i p_i + \sum_{i=1}^t K_i \gamma_i + \sum_{i=1}^\eta G_{i,g_i} \alpha_i); \eta_1, \dots, \eta_s), \\
 & \quad \vdots \\
 & (-\text{n}'-(\mu + \sum_{i=1}^s \rho_i p_i + \sum_{i=1}^t K_i \gamma_i + \sum_{i=1}^\eta G_{i,g_i} \alpha_i); \eta_1, \dots, \eta_s), \\
 & \quad \vdots \\
 & \quad \vdots \\
 & (-(\mu + \sum_{i=1}^s \rho_i p_i + \sum_{i=1}^t K_i \gamma_i + \sum_{i=1}^\eta G_{i,g_i} \alpha_i); \eta_1, \dots, \eta_s), \\
 & \quad \vdots \\
 & \quad \vdots \\
 & (-1-(\mu + \sum_{i=1}^s \rho_i p_i + \sum_{i=1}^t K_i \gamma_i + \sum_{i=1}^\eta G_{i,g_i} \alpha_i); \eta_1, \dots, \eta_s); \mathfrak{A}; A' \\
 & \quad \vdots \\
 & \quad \vdots \\
 & (-1-\text{n}'-(\mu + \sum_{i=1}^s \rho_i p_i + \sum_{i=1}^t K_i \gamma_i + \sum_{i=1}^\eta G_{i,g_i} \alpha_i); \eta_1, \dots, \eta_s), \mathfrak{B}; B' \Bigg) - \sum_{p_1, \dots, p_s=0}^{\infty} \\
 & \sum_{K_1=0}^{[N_1/M'_1]} \dots \sum_{K_t=0}^{[N_t/M'_t]} \sum_{n'=0}^{\infty} \sum_{G_1, \dots, G_r=0}^{\infty} \sum_{g_1=0}^{M_1} \dots \sum_{g_r=0}^{M_r} \frac{(-)^{G_1+\dots+G_r}}{\delta_{g_1} G_1! \dots \delta_{g_r} G_r!} G(\eta_{G_1, g_1}, \dots, \eta_{G_r, g_r}) a_1 \\
 & \frac{(\mu)_{p_1+\dots+p_s}}{p_1! \dots p_s! (a + \lambda_1 p_1 + \dots + \lambda_s p_s)^z} t_1^{p_1} \dots t_s^{p_s} z_1^{\eta_{G_1, g_1}} \dots z_r^{\eta_{G_r, g_r}} y_1^{K_1} \dots y_t^{K_t} \frac{(ab)^{n'}}{n' + 1} \\
 & a^{\sum_{i=1}^s \rho_i p_i + \sum_{i=1}^t K_i \gamma_i + \sum_{i=1}^r \eta_{G_i, g_i} \alpha_i} I_{U: p_R+3, q_R+3; W}^{V; 0, n_R+3; X} \left(\begin{array}{c} a^{\eta_1} Z_1 \\ \vdots \\ \dots \\ a^{\eta_R} Z_R \end{array} \right) \\
 & A : (1-(\mu + \sum_{i=1}^s \rho_i p_i + \sum_{i=1}^t K_i \gamma_i + \sum_{i=1}^\eta G_{i,g_i} \alpha_i); \eta_1, \dots, \eta_s), \\
 & \quad \vdots \\
 & \quad \vdots \\
 & B : (-(\mu + \sum_{i=1}^s \rho_i p_i + \sum_{i=1}^t K_i \gamma_i + \sum_{i=1}^\eta G_{i,g_i} \alpha_i); \eta_1, \dots, \eta_s), \\
 & \quad \vdots \\
 & \quad \vdots \\
 & (1-(\mu + \sum_{i=1}^s \rho_i p_i + \sum_{i=1}^t K_i \gamma_i + \sum_{i=1}^\eta G_{i,g_i} \alpha_i); \eta_1, \dots, \eta_s), \\
 & \quad \vdots \\
 & \quad \vdots \\
 & (-\frac{1}{2}\text{-n}'-(\mu + \sum_{i=1}^s \rho_i p_i + \sum_{i=1}^t K_i \gamma_i + \sum_{i=1}^\eta G_{i,g_i} \alpha_i); \eta_1, \dots, \eta_s),
 \end{aligned}$$

$$\left(\begin{matrix} (-n'-(\mu + \sum_{i=1}^s \rho_i p_i + \sum_{i=1}^t K_i \gamma_i + \sum_{i=1}^{\eta} G_{i,g_i} \alpha_i); \eta_1, \dots, \eta_s), \mathfrak{A}; A' \\ \vdots \\ (-(\mu + \sum_{i=1}^s \rho_i p_i + \sum_{i=1}^t K_i \gamma_i + \sum_{i=1}^{\eta} G_{i,g_i} \alpha_i); \eta_1, \dots, \eta_s), \mathfrak{B}; B' \end{matrix} \right) - \sum_{p_1, \dots, p_s=0}^{\infty}$$

$$\sum_{K_1=0}^{[N_1/M'_1]} \cdots \sum_{K_t=0}^{[N_t/M'_t]} \sum_{n'=0}^{\infty} \sum_{G_1, \dots, G_r=0}^{\infty} \sum_{g_1=0}^{M_1} \cdots \sum_{g_r=0}^{M_r} \frac{(-)^{G_1+\dots+G_r}}{\delta_{g_1} G_1! \cdots \delta_{g_r} G_r!} G(\eta_{G_1, g_1}, \dots, \eta_{G_r, g_r}) a_1$$

$$\frac{(\mu)_{p_1+\dots+p_s}}{p_1! \cdots p_s! (a + \lambda_1 p_1 + \dots + \lambda_s p_s) z} t_1^{p_1} \cdots t_s^{p_s} z_1^{\eta_{G_1, g_1}} \cdots z_r^{\eta_{G_r, g_r}} y_1^{K_1} \cdots y_t^{K_t} \frac{(ab)^{n'}}{(n' + 1)^2}$$

$$a^{\sum_{i=1}^s \rho_i p_i + \sum_{i=1}^t K_i \gamma_i + \sum_{i=1}^r \eta_{G_i, g_i} \alpha_i} I_{U: p_R+2, q_R+2; W}^{V; 0, n_R+2; X} \left(\begin{matrix} a^{\eta_1} Z_1 \\ \vdots \\ \dots \\ a^{\eta_R} Z_R \end{matrix} \right)$$

$$A ; (1-(\mu + \sum_{i=1}^s \rho_i p_i + \sum_{i=1}^t K_i \gamma_i + \sum_{i=1}^{\eta} G_{i,g_i} \alpha_i); \eta_1, \dots, \eta_s),$$

$$\vdots$$

$$B ; (-\frac{1}{2}n'-(\mu + \sum_{i=1}^s \rho_i p_i + \sum_{i=1}^t K_i \gamma_i + \sum_{i=1}^{\eta} G_{i,g_i} \alpha_i); \eta_1, \dots, \eta_s),$$

$$\left(\begin{matrix} (-n'-(\mu + \sum_{i=1}^s \rho_i p_i + \sum_{i=1}^t K_i \gamma_i + \sum_{i=1}^{\eta} G_{i,g_i} \alpha_i); \eta_1, \dots, \eta_s), \mathfrak{A}; A' \\ \vdots \\ (-(\mu + \sum_{i=1}^s \rho_i p_i + \sum_{i=1}^t K_i \gamma_i + \sum_{i=1}^{\eta} G_{i,g_i} \alpha_i); \eta_1, \dots, \eta_s), \mathfrak{B}; B' \end{matrix} \right) \Bigg] \quad (4.1)$$

Provided that

$$a) \min\{\rho_i \gamma_j, \alpha_k, \eta_l\} > 0, i = 1, \dots, s, j = 1, \dots, t, k = 1, \dots, r, l = 1, \dots, R, Re(z) > 0$$

$$b) Re[s' + \sum_{i=1}^s \rho_i p_i + \sum_{i=1}^r \alpha_i \min_{1 \leq j \leq M_i} \frac{d_j^{(i)}}{\delta_j^{(i)}} + \sum_{i=1}^R \eta_i \min_{1 \leq j \leq m^{(i)}} \frac{b_j^{(i)}}{\beta_j^{(i)}}] > -2$$

$$c) |arg z_k| < \frac{1}{2} A_i^{(k)} \pi, \text{ where } A_i^{(k)} \text{ is defined by (1.5); } i = 1, \dots, r$$

$$d) |arg Z_k| < \frac{1}{2} \Omega_i^{(k)} \pi, \text{ where } \Omega_i^{(k)} \text{ is defined by (1.11); } i = 1, \dots, s$$

e) The multiple serie occuring on the right-hand side of (3.1) is absolutely and uniformly convergent.

$$f) |arg(1 - ab)| < \pi, a > 0$$

$$g) f) |x_i| < 1, i = 1 \cdots, s, s \in \mathbb{Z}^+, \mu \in \mathbb{C} \setminus \{0, -1, -2, \dots\} \text{ and } a \in \mathbb{C} \setminus \{-(\lambda_i, p_i), p_i \in N \cup \{0\}\}$$

Proof

First, expressing the multiple generalized the Zeta-function $\phi_{\mu, \lambda_1, \dots, \lambda_s}^{(s)}(x_1 X_1^{\rho_1}, \dots, x_s X_s^{\rho_s}, z, a)$ in multiple serie with the help of equation (2.1), the Aleph-function of r variables in series with the help of equation (1.6), the general class of polynomial of several variables $S_{N_1, \dots, N_t}^{M_1, \dots, M_t}$ with the help of equation (1.22) and and the Prasad's multivariable I-function of s variables in Mellin-Barnes contour integral with the help of equation (1.9), changing the order of integration ans summation (which is easily seen to be justified due to the absolute convergence of the integral and the summations involved in the process) and then evaluating the resulting integral with the help of equation (3.1) and expressing the generalized hypergeometric functions ${}_3F_2$ and ${}_4F_3$ in series. Use several times the following relations $\Gamma(a)(a)_n = \Gamma(a+n)$ and $a = \frac{\Gamma(a+1)}{\Gamma(a)}$ with $Re(a) > 0$. Finally interpreting the result thus obtained with the Mellin-barnes contour integral, we arrive at the desired result.

The quantities $U, V, W, X, A, B, \mathfrak{A}, \mathfrak{B}, A'$ and B' are defined by the equations (1.14) to (1;19)

5. Particular case

If $U = V = A = B = 0$, the multivariable I-function defined by Prasad degenerere in multivariable H-function defined by Srivastava et al [7]. We have the following result.

$$\int_0^1 x^{\mu-1} \ln \left(\frac{\sqrt{a} + \sqrt{a-x}}{\sqrt{x}} \right) Li_2(bx) \phi_{\mu, \lambda_1, \dots, \lambda_s}^{(s)}(t_1 x_1^{\rho_1}, \dots, t_s x_s^{\rho_s}, z, a) S_{N_1, \dots, N_t}^{M'_1, \dots, M'_t} \left(\begin{matrix} y_1 x^{\gamma_1} \\ \vdots \\ y_t x^{\gamma_t} \end{matrix} \right)$$

$$\mathfrak{N}_{u:w}^{0,N:v} \left(\begin{matrix} z_1 x^{\alpha_1} \\ \vdots \\ z_r x^{\alpha_r} \end{matrix} \right) H_{p_R, q_R; W}^{0, n_R; X} \left(\begin{matrix} Z_1 x^{\eta_1} \\ \vdots \\ Z_R x^{\eta_R} \end{matrix} \right) dx = \frac{\sqrt{\pi} a^{\mu+1} b}{2} \left[\sum_{p_1, \dots, p_s=0}^{\infty} \right.$$

$$\sum_{K_1=0}^{[N_1/M'_1]} \cdots \sum_{K_t=0}^{[N_t/M'_t]} \sum_{n'=0}^{\infty} \sum_{G_1, \dots, G_r=0}^{\infty} \sum_{g_1=0}^{M_1} \cdots \sum_{g_r=0}^{M_r} \frac{(-)^{G_1+\dots+G_r}}{\delta_{g_1} G_1! \cdots \delta_{g_r} G_r!} G(\eta_{G_1, g_1}, \dots, \eta_{G_r, g_r}) a_1$$

$$\frac{(\mu)_{p_1+\dots+p_s}}{p_1! \cdots p_s! (a + \lambda_1 p_1 + \dots + \lambda_s p_s)^z} t_1^{p_1} \cdots t_s^{p_s} z_1^{\eta_{G_1, g_1}} \cdots z_r^{\eta_{G_r, g_r}} y_1^{K_1} \cdots y_t^{K_t} (ab)^{n'}$$

$$a^{\sum_{i=1}^s \rho_i p_i + \sum_{i=1}^t K_i \gamma_i + \sum_{i=1}^r \eta_{G_i, g_i} \alpha_i} H_{p_R+5, q_R+5; W}^{0, n_R+5; X} \left(\begin{matrix} a^{\eta_1} Z_1 \\ \vdots \\ \dots \\ a^{\eta_R} Z_R \end{matrix} \right)$$

$$\begin{aligned} & (1-(\mu + \sum_{i=1}^s \rho_i p_i + \sum_{i=1}^t K_i \gamma_i + \sum_{i=1}^r \eta_{G_i, g_i} \alpha_i); \eta_1, \dots, \eta_s), \\ & \quad \vdots \\ & \quad \vdots \\ & (-1-(\mu + \sum_{i=1}^s \rho_i p_i + \sum_{i=1}^t K_i \gamma_i + \sum_{i=1}^r \eta_{G_i, g_i} \alpha_i); \eta_1, \dots, \eta_s), \end{aligned}$$

$$\begin{aligned}
 & (1-(\mu + \sum_{i=1}^s \rho_i p_i + \sum_{i=1}^t K_i \gamma_i + \sum_{i=1}^\eta G_{i,g_i} \alpha_i); \eta_1, \dots, \eta_s), \\
 & \quad \vdots \\
 & \quad \vdots \\
 & (-\frac{1}{2}\text{-n}'-(\mu + \sum_{i=1}^s \rho_i p_i + \sum_{i=1}^t K_i \gamma_i + \sum_{i=1}^\eta G_{i,g_i} \alpha_i); \eta_1, \dots, \eta_s), \\
 & \quad \vdots \\
 & (-\text{n}'-(\mu + \sum_{i=1}^s \rho_i p_i + \sum_{i=1}^t K_i \gamma_i + \sum_{i=1}^\eta G_{i,g_i} \alpha_i); \eta_1, \dots, \eta_s), \\
 & \quad \vdots \\
 & \quad \vdots \\
 & (-(\mu + \sum_{i=1}^s \rho_i p_i + \sum_{i=1}^t K_i \gamma_i + \sum_{i=1}^\eta G_{i,g_i} \alpha_i); \eta_1, \dots, \eta_s), \\
 & \quad \vdots \\
 & (-\text{n}'-(\mu + \sum_{i=1}^s \rho_i p_i + \sum_{i=1}^t K_i \gamma_i + \sum_{i=1}^\eta G_{i,g_i} \alpha_i); \eta_1, \dots, \eta_s), \\
 & \quad \vdots \\
 & \quad \vdots \\
 & (-(\mu + \sum_{i=1}^s \rho_i p_i + \sum_{i=1}^t K_i \gamma_i + \sum_{i=1}^\eta G_{i,g_i} \alpha_i); \eta_1, \dots, \eta_s), \\
 & \quad \vdots \\
 & \quad \vdots \\
 & (-1-(\mu + \sum_{i=1}^s \rho_i p_i + \sum_{i=1}^t K_i \gamma_i + \sum_{i=1}^\eta G_{i,g_i} \alpha_i); \eta_1, \dots, \eta_s); \mathfrak{A}; A' \\
 & \quad \vdots \\
 & \quad \vdots \\
 & (-1-\text{n}'-(\mu + \sum_{i=1}^s \rho_i p_i + \sum_{i=1}^t K_i \gamma_i + \sum_{i=1}^\eta G_{i,g_i} \alpha_i); \eta_1, \dots, \eta_s); \mathfrak{B}; B' \Bigg) - \sum_{p_1, \dots, p_s=0}^{\infty}
 \end{aligned}$$

$$\sum_{K_1=0}^{[N_1/M'_1]} \cdots \sum_{K_t=0}^{[N_t/M'_t]} \sum_{n'=0}^{\infty} \sum_{G_1, \dots, G_r=0}^{\infty} \sum_{g_1=0}^{M_1} \cdots \sum_{g_r=0}^{M_r} \frac{(-)^{G_1+\dots+G_r}}{\delta_{g_1} G_1! \cdots \delta_{g_r} G_r!} G(\eta_{G_1, g_1}, \dots, \eta_{G_r, g_r}) a_1$$

$$\frac{(\mu)_{p_1+\dots+p_s}}{p_1! \cdots p_s! (a + \lambda_1 p_1 + \dots + \lambda_s p_s)^z} t_1^{p_1} \cdots t_s^{p_s} z_1^{\eta_{G_1, g_1}} \cdots z_r^{\eta_{G_r, g_r}} y_1^{K_1} \cdots y_t^{K_t} \frac{(ab)^{n'}}{n' + 1}$$

$$a^{\sum_{i=1}^s \rho_i p_i + \sum_{i=1}^t K_i \gamma_i + \sum_{i=1}^r \eta_{G_{i,g_i}} \alpha_i} H_{p_R+3, q_R+3; W}^{0, n_R+3; X} \left(\begin{array}{c} a^{\eta_1} Z_1 \\ \vdots \\ \dots \\ a^{\eta_R} Z_R \end{array} \right)$$

$$\begin{aligned}
 & (1-(\mu + \sum_{i=1}^s \rho_i p_i + \sum_{i=1}^t K_i \gamma_i + \sum_{i=1}^\eta G_{i,g_i} \alpha_i); \eta_1, \dots, \eta_s), \\
 & \quad \vdots \\
 & \quad \vdots \\
 & (-(\mu + \sum_{i=1}^s \rho_i p_i + \sum_{i=1}^t K_i \gamma_i + \sum_{i=1}^\eta G_{i,g_i} \alpha_i); \eta_1, \dots, \eta_s),
 \end{aligned}$$

$$\begin{aligned}
 & (1-(\mu + \sum_{i=1}^s \rho_i p_i + \sum_{i=1}^t K_i \gamma_i + \sum_{i=1}^\eta G_{i,g_i} \alpha_i); \eta_1, \dots, \eta_s), \\
 & \quad \vdots \\
 & \quad \vdots \\
 & (-\frac{1}{2}\text{-n}'-(\mu + \sum_{i=1}^s \rho_i p_i + \sum_{i=1}^t K_i \gamma_i + \sum_{i=1}^\eta G_{i,g_i} \alpha_i); \eta_1, \dots, \eta_s),
 \end{aligned}$$

$$\left(\begin{array}{c} (-n'-(\mu + \sum_{i=1}^s \rho_i p_i + \sum_{i=1}^t K_i \gamma_i + \sum_{i=1}^{\eta} G_{i,g_i} \alpha_i); \eta_1, \dots, \eta_s), \mathfrak{A}; A' \\ \vdots \\ (-(\mu + \sum_{i=1}^s \rho_i p_i + \sum_{i=1}^t K_i \gamma_i + \sum_{i=1}^{\eta} G_{i,g_i} \alpha_i); \eta_1, \dots, \eta_s), \mathfrak{B}; B' \end{array} \right) - \sum_{p_1, \dots, p_s=0}^{\infty}$$

$$\sum_{K_1=0}^{[N_1/M'_1]} \cdots \sum_{K_t=0}^{[N_t/M'_t]} \sum_{n'=0}^{\infty} \sum_{G_1, \dots, G_r=0}^{\infty} \sum_{g_1=0}^{M_1} \cdots \sum_{g_r=0}^{M_r} \frac{(-)^{G_1+\dots+G_r}}{\delta_{g_1} G_1! \cdots \delta_{g_r} G_r!} G(\eta_{G_1, g_1}, \dots, \eta_{G_r, g_r}) a_1$$

$$\frac{(\mu)_{p_1+\dots+p_s}}{p_1! \cdots p_s! (a + \lambda_1 p_1 + \dots + \lambda_s p_s)^z} t_1^{p_1} \cdots t_s^{p_s} z_1^{\eta_{G_1, g_1}} \cdots z_r^{\eta_{G_r, g_r}} y_1^{K_1} \cdots y_t^{K_t} \frac{(ab)^{n'}}{(n' + 1)^2}$$

$$a^{\sum_{i=1}^s \rho_i p_i + \sum_{i=1}^t K_i \gamma_i + \sum_{i=1}^r \eta_{G_i, g_i} \alpha_i} H_{p_R+2, q_R+2; W}^{0, n_R+2; X} \left(\begin{array}{c} a^{\eta_1} Z_1 \\ \vdots \\ \dots \\ a^{\eta_R} Z_R \end{array} \right)$$

$$(1-(\mu + \sum_{i=1}^s \rho_i p_i + \sum_{i=1}^t K_i \gamma_i + \sum_{i=1}^{\eta} G_{i,g_i} \alpha_i); \eta_1, \dots, \eta_s),$$

$$\vdots$$

$$(-\frac{1}{2}n'-(\mu + \sum_{i=1}^s \rho_i p_i + \sum_{i=1}^t K_i \gamma_i + \sum_{i=1}^{\eta} G_{i,g_i} \alpha_i); \eta_1, \dots, \eta_s),$$

$$\left(\begin{array}{c} (-n'-(\mu + \sum_{i=1}^s \rho_i p_i + \sum_{i=1}^t K_i \gamma_i + \sum_{i=1}^{\eta} G_{i,g_i} \alpha_i); \eta_1, \dots, \eta_s), \mathfrak{A}; A' \\ \vdots \\ (-(\mu + \sum_{i=1}^s \rho_i p_i + \sum_{i=1}^t K_i \gamma_i + \sum_{i=1}^{\eta} G_{i,g_i} \alpha_i); \eta_1, \dots, \eta_s), \mathfrak{B}; B' \end{array} \right) \Bigg] \quad (4.1)$$

7. Conclusion

In this paper we have evaluated a generalized finite integral involving the multivariable Aleph-function, a class of polynomials of several variables a sequence of functions and the multivariable I-function defined by Prasad. The integral established in this paper is of very general nature as it contains Multivariable Aleph-function, which is a general function of several variables studied so far. Thus, the integral established in this research work would serve as a key formula from which, upon specializing the parameters, as many as desired results involving the special functions of one and several variables can be obtained.

REFERENCES

- [1] Berndt B.C. Ramanujan's notebooks. Part iv. New York. Springer-Verlag 1994. page 323-326.
- [2] Bin-Saad M.G. Pathan M.A. And Hanballa A.M. On power series associated with generalized multiple Zeta-functions. Math.Sci.Res.J. 17(10) 2013, page 279-291.
- [3] Brychkow Y.A. Handbook of Special Functions. Derivatives. Integrals, Series and Other Formulas. CRC. Press. Taykor and Francis Group. Boca. Raton. London. New York. 2008.

- [4] Y.N. Prasad , Multivariable I-function , Vijnana Parishad Anusandhan Patrika 29 (1986) , page 231-237.
- [5] Sharma C.K.and Ahmad S.S.: On the multivariable I-function. Acta ciencia Indica Math , 1994 vol 20,no2, p 113-116.
- [6] Srivastava H.M. A multilinear generating function for the Konhauser set of biorthogonal polynomials suggested by Laguerre polynomial, Pacific. J. Math. 177(1985), page183-191.
- [7] H.M. Srivastava And R.Panda. Some expansion theorems and generating relations for the H-function of several complex variables. Comment. Math. Univ. St. Paul. 24(1975), p.119-137.

Personal adress : 411 Avenue Joseph Raynaud
Le parc Fleuri , Bat B
83140 , Six-Fours les plages
Tel : 06-83-12-49-68
Department : VAR
Country : **FRANCE**