

EVEN VERTEX MAGIC TOTAL LABELING OF SOME 2-REGULAR GRAPHS

CT. NAGARAJ^{#1}, C.Y. PONNAPPAN^{*2}, G. PRABAKARAN^{#3}

¹ Research Scholar, Department of Mathematics

Research and Development centre

Bharathiar University, Coimbatore-641046, Tamilnadu, India.

e-mail : *ct_nagaraj@yahoo.co.in*

² Department of Mathematics

Government Arts College, Melur - 625 106, Tamilnadu, India.

e-mail : *pons_mdu1969@yahoo.com*

³ Department of Mathematics

Thiagarajar College, Madurai - 625009, Tamilnadu, India.

e-mail : *tcmaths@yahoo.in*

ABSTRACT. A vertex magic total labeling of a graph $G(V, E)$ is defined as one - to - one map f from $V \cup E$ onto the integers $\{1, 2, 3, \dots, |V| + |E|\}$ such that $f(u) + \sum f(uv)$ where the sum is over all vertices v adjacent to u , is a constant, independent of the choice of vertex u . Such a labeling is even if $f(V(G)) = \{2, 4, 6, \dots, 2n\}$. In this paper, we present an even vertex magic total labeling of some 2 - regular graphs.

Key words: Even vertex magic total labeling, Even vertex magic graph.

2010 Mathematics Subject Classification: 05C78.

1. INTRODUCTION

All graphs considered in this paper are simple, finite and undirected. The graph G has vertex set $V = V(G)$ and edge set $E = E(G)$ and we let $n = |V|$ and $m = |E|$. The degree of a vertex v is the number of edges that have v as an end point and the set of neighbours of v is denoted by $N(v)$.

A labeling of a graph G is a mapping that carries a set of graph elements usually the vertices and / or edges, into a set of numbers, usually integers, called labels. Many kinds of labelings have been studied and an excellent survey of graph labelings can be found in [1]

Macdougall et al. [2] introduced the concept of vertex magic total labeling. If G is a finite simple undirected graph with n vertices and e edges, then a vertex

magic total labeling (*VMTL*) is a bijection f from $V(G) \cup E(G)$ to the integers $1, 2, \dots, m + n$ with the property that for every u in $V(G)$, $f(u) + \sum_{v \in N(u)} f(uv) = k$, for some constant k (which we call the magic constant for f).

MacDougall et al. [4] introduced the concept of super vertex magic total labeling. They call a vertex magic total labeling is super if $f(V(G)) = \{1, 2, 3, \dots, n\}$. In this labeling, the smallest labels are assigned to the vertices.

Nagaraj et al. [3] introduced the concept of an Even vertex magic total labeling. They call vertex magic total labeling is even if $f(V(G)) = \{2, 4, 6, \dots, 2n\}$. A graph is called an even vertex magic if the graph has an even vertex magic total labeling.

2. MAIN RESULTS

In this section, we will use the notation $[x_1, x_2, x_3, \dots, x_k]$ to denote labels of a cycle of length k in the sequence vertex - edge - vertex - edge and so on. Thus, x_1 is the label of a vertex with weight $x_1 + x_2 + x_{2k}$.

Theorem 2.1. *The graph $C_3 \cup C_{4t}, t > 1$ is even vertex magic graph.*

Proof. we label the vertex and edges of C_3 with $[8t+2, 2t+5, 8t+4, 2t+1, 8t+6, 2t+3]$ and then edge label C_{4t} as follows.

$$f(e_i) = \begin{cases} i + 2 & \text{if } i \text{ odd, } 1 \leq i \leq 2t - 3 \\ i + 8 & \text{if } i \text{ odd, } 2t - 1 \leq i \leq 4t - 3 \\ 1 & \text{if } i = 4t - 1 \\ 4t + 5 + i & \text{if } i \text{ even} \end{cases}$$

The vertex label C_{4t} as follows

$$f(v_i) = \begin{cases} 4t + 2 & i = 1 \\ 8t + 4 - 2i & i = 2, 3, \dots, 2t - 2 \\ 8t - 2 - 2i & 2t - 1 \leq i \leq 4t - 2 \\ 4t + 4 & i = 4t \\ 4t + 6 & i = 4t - 1. \end{cases}$$


We show firstly that each edge label appears only once.

Edge Number	Range of labels
$4t - 1$	1
i odd, $1 \leq i \leq 2t - 3$	3, 5, 7, ..., $2t - 1$
C_3 edges	$2t + 1, 2t + 3, 2t + 5$
i odd, $2t - 1 \leq i \leq 4t - 3$	$2t + 7, 2t + 9, \dots, 4t + 5$
i even	$4t + 7, 4t + 9, \dots, 8t + 5.$

Next we show that each vertex label appear only once

Vertex Number	Range of labels
1	$4t + 2$
$2t - 1 \leq i \leq 4t - 2$	2, 4, 6, ..., $4t - 2, 4t$
$4t$	$4t + 4$
$4t - 1$	$4t + 6$
$2 \leq i \leq 2t - 2$	$4t + 8, 4t + 10, \dots, 8t - 2, 8t.$
C_3 vertex	$8t + 2, 8t + 4, 8t + 6$

$\therefore C_3 \cup C_{4t}, t > 1$ is an even vertex magic graph with magic constant $k = 12t + 10$.



□




FIGURE 1. $C_3 \cup C_{16}$, Magic constant $k = 58$

Theorem 2.2. *The graph $C_3 \cup C_{4t+2}$, $t \geq 1$ is an even vertex magic graph.*

Proof. For $t = 1$ and $t = 2$ the labelings are

[16, 3, 18, 7, 12, 9] [10, 1, 14, 13, 4, 11, 2, 15, 8, 5, 6, 17] and

[24, 5, 26, 9, 20, 11] [14, 1, 22, 17, 8, 15, 6, 19, 18, 3, 16, 21, 12, 7, 10, 23, 4, 13, 2, 25]

For $t \geq 3$, we label the vertex and edge of C_3 with $[8t + 8, 2t + 1, 8t + 10, 2t + 5, 8t + 4, 2t + 7]$ and then vertex label C_{4t+2} as follows.

$$f(v_i) = \begin{cases} 4t + 6 & \text{if } i = 1 \\ 8t + 6 & \text{if } i = 2 \\ 4t & \text{if } i = 3 \\ 4t - 2 & \text{if } i = 4 \\ 8t + 12 - 2i & \text{if } 5 \leq i \leq 2t + 2 \\ 4t + 4 & \text{if } i = 2t + 3 \\ 4t + 2 & \text{if } i = 2t + 4 \\ 8t + 6 - 2i & \text{if } 2t + 5 \leq i \leq 4t + 2. \end{cases}$$

The edge label C_{4t+2} as follows

$$f(e_i) = \begin{cases} 1 & \text{if } i = 1 \\ 4t + 7 & \text{if } i = 3 \\ i - 2 & \text{if } i \text{ odd}, 5 \leq i \leq 2t + 1 \\ 2t + 3 & \text{if } i = 2t + 3 \\ i + 4 & \text{if } i \text{ odd}, 2t + 5 \leq i \leq 4t + 1 \\ i + 4t + 7 & \text{if } i \text{ even.} \end{cases}$$

$\therefore C_3 \cup C_{4t+2}, t \geq 1$ is an even vertex magic graph magic constant $k = 12t + 16$.




FIGURE 2. $C_3 \cup C_{18} \quad k = 64$

□

Theorem 2.3. *The graph $C_4 \cup C_{4t-1}$ for $t > 1$ is an even vertex magic graph.*

Proof. We label the vertex and edges of C_4 consecutively as $[2, 4t+3, 4, 8t+3, 8, 4t-1, 6, 8t+5]$

We label the edges of C_{4t-1} consecutively as follows

$$f(e_i) = \begin{cases} i + 2 & \text{if } i \equiv 1 \pmod{4}, i < 4t - 3 \\ 4t + 5 & \text{if } i = 2 \\ i + 4t + 1 & \text{if } i \equiv 2 \pmod{4}, i > 2 \\ i - 2 & \text{if } i \equiv 3 \pmod{4}, i < 4t - 1 \\ i + 4t + 5 & \text{if } i \equiv 0 \pmod{4} \\ 4t - 3 & \text{if } i = 4t - 3 \\ 4t - 1 & \text{if } i = 4t - 1 \end{cases}$$

We label the vertex of C_{4t-1} consecutively as follows

$$f(v_i) = \begin{cases} 8t + 6 & \text{if } i = 1 \\ 8t - 6 & \text{if } i = 5, t \neq 2 \\ 12 & \text{if } t = 2, i = 5 \\ 8t + 2 & \text{if } i = 2 \\ 8t - 2i + 8 & \text{if } i \equiv 6 \pmod{4} \\ 14 & \text{if } i = 4t - 2 \\ 10 & \text{if } i = 4t - 1 \\ 8t + 4 & \text{if } i = 3 \\ 8t - 2i + 12 & \text{if } 7 \leq i \leq 4t - 5, i \equiv 3 \pmod{4} \\ 8t - 2i + 8 & \text{if } i \equiv 0 \pmod{4} \\ 8t - 2i + 4 & \text{if } i \equiv 5 \pmod{4}, i \leq 4t - 7 \\ 12 & \text{if } i = 4t - 3 \end{cases}$$

\therefore The graph $C_4 \cup C_{4t-1}, t > 1$ is an even vertex magic graph with magic constant $k = 12t + 10$.




FIGURE 3. $C_4 \cup C_{15}$ $k = 58$

□

Theorem 2.4. *The graph $C_4 \cup C_{4t-3}, t > 1$ is an even vertex magic graph.*

Proof. We label the edges of C_4 consecutively as $[2, 4t + 1, 8t - 1, 8, 4t - 3, 6, 8t + 1]$ and then edge label C_{4t-3} as follows

$$f(e_i) = \begin{cases} i + 2 & \text{if } i \equiv 1 \pmod{4} \\ i + 4t + 3 & \text{if } i \equiv 2 \pmod{4} \\ i - 2 & \text{if } i \equiv 3 \pmod{4} \\ i + 4t - 1 & \text{if } i \equiv 0 \pmod{4} \end{cases}$$

The vertex label C_{4t-3} as follows

$$f(v_i) = \begin{cases} 8t - 2i + 4 & \text{if } i \equiv 1, 3 \pmod{4} \\ 8t - 2i & \text{if } i \equiv 2 \pmod{4} \\ 8t - 2i + 8 & \text{if } i \equiv 0 \pmod{4} \end{cases}$$




FIGURE 4. $C_4 \cup C_{13} \quad k = 52$

□

REFERENCES

- [1] J.A. Gallian, *A dynamic survey of graph labeling electronic*, J.Combinatorics 5(2002) # D56.
- [2] J.A. MacDougall, M.Miller, Slamin, W.D.Wallis, *Vertex magic total labeling of graphs*, Util.Math 61 (2002) 3-21.

- [3] CT.Nagaraj, C.Y.Ponnappan, G.Prabakaran, *Even vertex magic total labeling of Trees*, *International Journal of pure and Applied Mathematics*, Volume 115, No.9, (2017), 363 - 375.
- [4] J.A.MacDougall, M.Miller, K.A.Sugeng, *Super vertex magic total labelling of graphs*, in: Proc.of the 15th Australian Workshop on Combinatorial Algorithms, 2004, pp.222-229.