

A theorem concerning a product of polynomials and multivariable Gimel-function

Frédéric Ayant

Teacher in High School , France

ABSTRACT

The object of this paper is to establish a general theorem pertaining to a product of class of polynomials and multivariable Gimel-function. Certain integrals are also obtained by application of the theorem. The theorem is quite general nature and capable of yielding a number of new, interesting and useful integrals as its special cases.

KEYWORDS : Multivariable Gimel-function, multiple integral contours, class of polynomials, hypergeometric function.

2010 Mathematics Subject Classification. 33C99, 33C60, 44A20

1. Introduction and preliminaries.

Throughout this paper, let \mathbb{C}, \mathbb{R} and \mathbb{N} be set of complex numbers, real numbers and positive integers respectively. Also $\mathbb{N}_0 = \mathbb{N} \cup \{0\}$.

We define a generalized transcendental function of several complex variables noted \mathfrak{J} .

$$\mathfrak{J}(z_1, \dots, z_r) = \mathfrak{J}_{p_{i_2}, q_{i_2}, \tau_{i_2}; R_2; p_{i_3}, q_{i_3}, \tau_{i_3}; R_3; \dots; p_{i_r}, q_{i_r}, \tau_{i_r}; R_r; p_{i(1)}, q_{i(1)}, \tau_{i(1)}; R^{(1)}; \dots; p_{i(r)}, q_{i(r)}, \tau_{i(r)}; R^{(r)}} \left(\begin{matrix} z_1 \\ \cdot \\ \cdot \\ \cdot \\ z_r \end{matrix} \right)$$

$$[(a_{2j}; \alpha_{2j}^{(1)}, \alpha_{2j}^{(2)}; A_{2j})]_{1, n_2}, [\tau_{i_2}(a_{2ji_2}; \alpha_{2ji_2}^{(1)}, \alpha_{2ji_2}^{(2)}; A_{2ji_2})]_{n_2+1, p_{i_2}}; [(a_{3j}; \alpha_{3j}^{(1)}, \alpha_{3j}^{(2)}, \alpha_{3j}^{(3)}; A_{3j})]_{1, n_3},$$

$$[\tau_{i_2}(b_{2ji_2}; \beta_{2ji_2}^{(1)}, \beta_{2ji_2}^{(2)}; B_{2ji_2})]_{1, q_{i_2}};$$

$$[\tau_{i_3}(a_{3ji_3}; \alpha_{3ji_3}^{(1)}, \alpha_{3ji_3}^{(2)}, \alpha_{3ji_3}^{(3)}; A_{3ji_3})]_{n_3+1, p_{i_3}}; \dots; [(a_{rj}; \alpha_{rj}^{(1)}, \dots, \alpha_{rj}^{(r)}; A_{rj})]_{1, n_r},$$

$$[\tau_{i_3}(b_{3ji_3}; \beta_{3ji_3}^{(1)}, \beta_{3ji_3}^{(2)}, \beta_{3ji_3}^{(3)}; B_{3ji_3})]_{1, q_{i_3}}; \dots;$$

$$[\tau_{i_r}(a_{rji_r}; \alpha_{rji_r}^{(1)}, \dots, \alpha_{rji_r}^{(r)}; A_{rji_r})]_{n_r+1, p_{i_r}}; [(c_j^{(1)}, \gamma_j^{(1)}; C_j^{(1)})]_{1, n^{(1)}}, [\tau_{i(1)}(c_{ji(1)}^{(1)}, \gamma_{ji(1)}^{(1)}; C_{ji(1)}^{(1)})]_{n^{(1)}+1, p_i^{(1)}}]$$

$$[\tau_{i_r}(b_{rji_r}; \beta_{rji_r}^{(1)}, \dots, \beta_{rji_r}^{(r)}; B_{rji_r})]_{1, q_r}; [(d_j^{(1)}, \delta_j^{(1)}; D_j^{(1)})]_{1, m^{(1)}}, [\tau_{i(1)}(d_{ji(1)}^{(1)}, \delta_{ji(1)}^{(1)}; D_{ji(1)}^{(1)})]_{m^{(1)}+1, q_i^{(1)}}]$$

$$; \dots; [(c_j^{(r)}, \gamma_j^{(r)}; C_j^{(r)})]_{1, n^{(r)}}, [\tau_{i(r)}(c_{ji(r)}^{(r)}, \gamma_{ji(r)}^{(r)}; C_{ji(r)}^{(r)})]_{n^{(r)}+1, p_i^{(r)}}]$$

$$; \dots; [(d_j^{(r)}, \delta_j^{(r)}; D_j^{(r)})]_{1, m^{(r)}}, [\tau_{i(r)}(d_{ji(r)}^{(r)}, \delta_{ji(r)}^{(r)}; D_{ji(r)}^{(r)})]_{m^{(r)}+1, q_i^{(r)}}]$$

$$= \frac{1}{(2\pi\omega)^r} \int_{L_1} \dots \int_{L_r} \psi(s_1, \dots, s_r) \prod_{k=1}^r \theta_k(s_k) z_k^{s_k} ds_1 \dots ds_r \tag{1.1}$$

with $\omega = \sqrt{-1}$

$$\psi(s_1, \dots, s_r) = \frac{\prod_{j=1}^{n_2} \Gamma^{A_{2j}} (1 - a_{2j} + \sum_{k=1}^2 \alpha_{2j}^{(k)} s_k)}{\sum_{i_2=1}^{R_2} [\tau_{i_2} \prod_{j=n_2+1}^{p_{i_2}} \Gamma^{A_{2ji_2}} (a_{2ji_2} - \sum_{k=1}^2 \alpha_{2ji_2}^{(k)} s_k) \prod_{j=1}^{q_{i_2}} \Gamma^{B_{2ji_2}} (1 - b_{2ji_2} + \sum_{k=1}^2 \beta_{2ji_2}^{(k)} s_k)]}$$

$$\frac{\prod_{j=1}^{n_3} \Gamma^{A_{3j}} (1 - a_{3j} + \sum_{k=1}^3 \alpha_{3j}^{(k)} s_k)}{\sum_{i_3=1}^{R_3} [\tau_{i_3} \prod_{j=n_3+1}^{p_{i_3}} \Gamma^{A_{3ji_3}} (a_{3ji_3} - \sum_{k=1}^3 \alpha_{3ji_3}^{(k)} s_k) \prod_{j=1}^{q_{i_3}} \Gamma^{B_{3ji_3}} (1 - b_{3ji_3} + \sum_{k=1}^3 \beta_{3ji_3}^{(k)} s_k)]}$$

$$\frac{\prod_{j=1}^{n_r} \Gamma^{A_{rj}} (1 - a_{rj} + \sum_{k=1}^r \alpha_{rj}^{(k)} s_k)}{\sum_{i_r=1}^{R_r} [\tau_{i_r} \prod_{j=n_r+1}^{p_{i_r}} \Gamma^{A_{rji_r}} (a_{rji_r} - \sum_{k=1}^r \alpha_{rji_r}^{(k)} s_k) \prod_{j=1}^{q_{i_r}} \Gamma^{B_{rji_r}} (1 - b_{rji_r} + \sum_{k=1}^r \beta_{rji_r}^{(k)} s_k)]} \tag{1.2}$$

and

$$\theta_k(s_k) = \frac{\prod_{j=1}^{m^{(k)}} \Gamma^{D_j^{(k)}} (d_j^{(k)} - \delta_j^{(k)} s_k) \prod_{j=1}^{n^{(k)}} \Gamma^{C_j^{(k)}} (1 - c_j^{(k)} + \gamma_j^{(k)} s_k)}{\sum_{i^{(k)}=1}^{R^{(k)}} [\tau_{i^{(k)}} \prod_{j=m^{(k)}+1}^{q_{i^{(k)}}} \Gamma^{D_{ji^{(k)}}^{(k)}} (1 - d_{ji^{(k)}}^{(k)} + \delta_{ji^{(k)}}^{(k)} s_k) \prod_{j=n^{(k)}+1}^{p_{i^{(k)}}} \Gamma^{C_{ji^{(k)}}^{(k)}} (c_{ji^{(k)}}^{(k)} - \gamma_{ji^{(k)}}^{(k)} s_k)]} \tag{1.3}$$

1) $[(c_j^{(1)}; \gamma_j^{(1)})]_{1, n_1}$ stands for $(c_1^{(1)}; \gamma_1^{(1)}), \dots, (c_{n_1}^{(1)}; \gamma_{n_1}^{(1)})$.

2) $n_2, \dots, n_r, m^{(1)}, n^{(1)}, \dots, m^{(r)}, n^{(r)}, p_{i_2}, q_{i_2}, R_2, \tau_{i_2}, \dots, p_{i_r}, q_{i_r}, R_r, \tau_{i_r}, p_{i^{(r)}}, q_{i^{(r)}}, \tau_{i^{(r)}}, R^{(r)} \in \mathbb{N}$ and verify :

$$0 \leq m_2, 0 \leq n_2 \leq p_{i_2}, \dots, 0 \leq m_r, 0 \leq n_r \leq p_{i_r}, 0 \leq m^{(1)} \leq q_{i^{(1)}}, \dots, 0 \leq m^{(r)} \leq q_{i^{(r)}}.$$

3) $\tau_{i_2} (i_2 = 1, \dots, R_2) \in \mathbb{R}^+$; $\tau_{i_r} \in \mathbb{R}^+ (i_r = 1, \dots, R_r)$; $\tau_{i^{(k)}} \in \mathbb{R}^+ (i = 1, \dots, R^{(k)}), (k = 1, \dots, r)$.

4) $\gamma_j^{(k)}, C_j^{(k)} \in \mathbb{R}^+; (j = 1, \dots, n_k); (k = 1, \dots, r); \delta_j^{(k)}, D_j^{(k)} \in \mathbb{R}^+; (j = 1, \dots, m_k); (k = 1, \dots, r)$.

$\alpha_{kj}^{(l)}, A_{kj} \in \mathbb{R}^+; (j = 1, \dots, n_k); (k = 2, \dots, r); (l = 1, \dots, k)$.

$\alpha_{kji_k}^{(l)}, A_{kji_k} \in \mathbb{R}^+; (j = n_k + 1, \dots, p_{i_k}); (k = 2, \dots, r); (l = 1, \dots, k)$.

$\beta_{kji_k}^{(l)}, B_{kji_k} \in \mathbb{R}^+; (j = m_k + 1, \dots, q_{i_k}); (k = 2, \dots, r); (l = 1, \dots, k)$.

$\delta_{ji^{(k)}}^{(k)} \in \mathbb{R}^+; (i = 1, \dots, R^{(k)}); (j = m^{(k)} + 1, \dots, q_{i^{(k)}}); (k = 1, \dots, r)$.

$\gamma_{ji^{(k)}}^{(k)} \in \mathbb{R}^+; (i = 1, \dots, R^{(k)}); (j = n^{(k)} + 1, \dots, p_{i^{(k)}}); (k = 1, \dots, r)$.

5) $c_j^{(k)} \in \mathbb{C}; (j = 1, \dots, n_k); (k = 1, \dots, r); d_j^{(k)} \in \mathbb{C}; (j = 1, \dots, m_k); (k = 1, \dots, r)$.

$a_{kji_k} \in \mathbb{C}; (j = n_k + 1, \dots, p_{i_k}); (k = 2, \dots, r)$.

$b_{kji_k} \in \mathbb{C}; (j = m_k + 1, \dots, q_{i_k}); (k = 2, \dots, r)$.

$d_{ji^{(k)}}^{(k)} \in \mathbb{C}; (i = 1, \dots, R^{(k)}); (j = m^{(k)} + 1, \dots, q_{i^{(k)}}); (k = 1, \dots, r)$.

$\gamma_{ji^{(k)}}^{(k)} \in \mathbb{C}; (i = 1, \dots, R^{(k)}); (j = n^{(k)} + 1, \dots, p_{i^{(k)}}); (k = 1, \dots, r)$.

The contour L_k is in the $s_k(k = 1, \dots, r)$ - plane and run from $\sigma - i\infty$ to $\sigma + i\infty$ where σ is a real number with loop, if necessary to ensure that the poles of $\Gamma^{A_{2j}} \left(1 - a_{2j} + \sum_{k=1}^2 \alpha_{2j}^{(k)} s_k \right) (j = 1, \dots, n_2), \Gamma^{A_{3j}} \left(1 - a_{3j} + \sum_{k=1}^3 \alpha_{3j}^{(k)} s_k \right) (j = 1, \dots, n_3), \dots, \Gamma^{A_{rj}} \left(1 - a_{rj} + \sum_{i=1}^r \alpha_{rj}^{(i)} \right) (j = 1, \dots, n_r), \Gamma^{C_j^{(k)}} \left(1 - c_j^{(k)} + \gamma_j^{(k)} s_k \right) (j = 1, \dots, n^{(k)})(k = 1, \dots, r)$ to the right of the contour L_k and the poles of $\Gamma^{D_j^{(k)}} \left(d_j^{(k)} - \delta_j^{(k)} s_k \right) (j = 1, \dots, m^{(k)})(k = 1, \dots, r)$ lie to the left of the contour L_k . The condition for absolute convergence of multiple Mellin-Barnes type contour (1.1) can be obtained of the corresponding conditions for multivariable H-function given by as :

$$|arg(z_k)| < \frac{1}{2} A_i^{(k)} \pi \text{ where}$$

$$A_i^{(k)} = \sum_{j=1}^{m^{(k)}} D_j^{(k)} \delta_j^{(k)} + \sum_{j=1}^{n^{(k)}} C_j^{(k)} \gamma_j^{(k)} - \tau_{i^{(k)}} \left(\sum_{j=m^{(k)}+1}^{q_i^{(k)}} D_{ji^{(k)}}^{(k)} \delta_{ji^{(k)}}^{(k)} + \sum_{j=n^{(k)}+1}^{p_i^{(k)}} C_{ji^{(k)}}^{(k)} \gamma_{ji^{(k)}}^{(k)} \right) - \tau_{i_2} \left(\sum_{j=n_2+1}^{p_{i_2}} A_{2ji_2} \alpha_{2ji_2}^{(k)} + \sum_{j=1}^{q_{i_2}} B_{2ji_2} \beta_{2ji_2}^{(k)} \right) - \dots - \tau_{i_r} \left(\sum_{j=n_r+1}^{p_{i_r}} A_{rji_r} \alpha_{rji_r}^{(k)} + \sum_{j=1}^{q_{i_r}} B_{rji_r} \beta_{rji_r}^{(k)} \right) \tag{1.4}$$

Following the lines of Braaksma ([2] p. 278), we may establish the the asymptotic expansion in the following convenient form :

$$\aleph(z_1, \dots, z_r) = O(|z_1|^{\alpha_1}, \dots, |z_r|^{\alpha_r}), \max(|z_1|, \dots, |z_r|) \rightarrow 0$$

$$\aleph(z_1, \dots, z_r) = O(|z_1|^{\beta_1}, \dots, |z_r|^{\beta_r}), \min(|z_1|, \dots, |z_r|) \rightarrow \infty \text{ where } i = 1, \dots, r :$$

$$\alpha_i = \min_{1 \leq j \leq m^{(i)}} \operatorname{Re} \left[D_j^{(i)} \left(\frac{d_j^{(i)}}{\delta_j^{(i)}} \right) \right] \text{ and } \beta_i = \max_{1 \leq j \leq n^{(i)}} \operatorname{Re} \left[C_j^{(i)} \left(\frac{c_j^{(i)} - 1}{\gamma_j^{(i)}} \right) \right]$$

Remark 1.

If $n_2 = \dots = n_{r-1} = p_{i_2} = q_{i_2} = \dots = p_{i_{r-1}} = q_{i_{r-1}} = 0$ and $A_{2j} = A_{2ji_2} = B_{2ji_2} = \dots = A_{rj} = A_{rji_r} = B_{rji_r} = 1$ $A_{rj} = A_{rji_r} = B_{rji_r} = 1$, then the multivariable Gimel-function reduces in the multivariable Aleph- function defined by Ayant [1].

Remark 2.

If $n_2 = \dots = n_r = p_{i_2} = q_{i_2} = \dots = p_{i_r} = q_{i_r} = 0$ and $\tau_{i_2} = \dots = \tau_{i_r} = \tau_{i^{(1)}} = \dots = \tau_{i^{(r)}} = R_2 = \dots = R_r = R^{(1)} = \dots = R^{(r)} = 1$, then the multivariable Gimel-function reduces in a multivariable I-function defined by Prathima et al. [4].

Remark 3.

If $A_{2j} = A_{2ji_2} = B_{2ji_2} = \dots = A_{rj} = A_{rji_r} = B_{rji_r} = 1$ and $\tau_{i_2} = \dots = \tau_{i_r} = \tau_{i^{(1)}} = \dots = \tau_{i^{(r)}} = R_2 = \dots = R_r = R^{(1)} = \dots = R^{(r)} = 1$, then the generalized multivariable Gimel-function reduces in multivariable I-function defined by Prasad [3].

Remark 4.

If the three above conditions are satisfied at the same time, then the generalized multivariable Gimel-function reduces in the multivariable H-function defined by Srivastava and Panda [7,8].

In your investigation, we shall use the following notations.

$$\mathbb{A} = [(a_{2j}; \alpha_{2j}^{(1)}, \alpha_{2j}^{(2)}; A_{2j})]_{1, n_2}, [\tau_{i_2}(a_{2ji_2}; \alpha_{2ji_2}^{(1)}, \alpha_{2ji_2}^{(2)}; A_{2ji_2})]_{n_2+1, p_{i_2}}, [(a_{3j}; \alpha_{3j}^{(1)}, \alpha_{3j}^{(2)}, \alpha_{3j}^{(3)}; A_{3j})]_{1, n_3},$$

$$[\tau_{i_3}(a_{3ji_3}; \alpha_{3ji_3}^{(1)}, \alpha_{3ji_3}^{(2)}, \alpha_{3ji_3}^{(3)}; A_{3ji_3})]_{n_3+1, p_{i_3}}; \dots; [(a_{(r-1)j}; \alpha_{(r-1)j}^{(1)}, \dots, \alpha_{(r-1)j}^{(r-1)}; A_{(r-1)j})]_{1, n_{r-1}},$$

$$[\tau_{i_{r-1}}(a_{(r-1)j_{i_{r-1}}}; \alpha_{(r-1)j_{i_{r-1}}}^{(1)}, \dots, \alpha_{(r-1)j_{i_{r-1}}}^{(r-1)}; A_{(r-1)j_{i_{r-1}}}n_{r-1+1, p_{i_{r-1}}}] \tag{1.5}$$

$$\mathbf{A} = [(a_{rj}; \alpha_{rj}^{(1)}, \dots, \alpha_{rj}^{(r)}; A_{rj})_{1, n_r}], [\tau_{i_r}(a_{rj_{i_r}}; \alpha_{rj_{i_r}}^{(1)}, \dots, \alpha_{rj_{i_r}}^{(r)}; A_{rj_{i_r}})_{n+1, p_{i_r}}] \tag{1.6}$$

$$A = [(c_j^{(1)}, \gamma_j^{(1)}; C_j^{(1)})_{1, n^{(1)}}], [\tau_{i^{(1)}}(c_{j_{i^{(1)}}}^{(1)}, \gamma_{j_{i^{(1)}}}^{(1)}; C_{j_{i^{(1)}}}^{(1)})_{n^{(1)}+1, p_i^{(1)}}]; \dots ; \\ [(c_j^{(r)}, \gamma_j^{(r)}; C_j^{(r)})_{1, n^{(r)}}], [\tau_{i^{(r)}}(c_{j_{i^{(r)}}}^{(r)}, \gamma_{j_{i^{(r)}}}^{(r)}; C_{j_{i^{(r)}}}^{(r)})_{n^{(r)}+1, p_i^{(r)}}] \tag{1.7}$$

$$\mathbb{B} = [\tau_{i_2}(b_{2j_{i_2}}; \beta_{2j_{i_2}}^{(1)}, \beta_{2j_{i_2}}^{(2)}; B_{2j_{i_2}})]_{1, q_{i_2}}, [\tau_{i_3}(b_{3j_{i_3}}; \beta_{3j_{i_3}}^{(1)}, \beta_{3j_{i_3}}^{(2)}, \beta_{3j_{i_3}}^{(3)}; B_{3j_{i_3}})]_{1, q_{i_3}}; \dots ;$$

$$[\tau_{i_{r-1}}(b_{(r-1)j_{i_{r-1}}}; \beta_{(r-1)j_{i_{r-1}}}^{(1)}, \dots, \beta_{(r-1)j_{i_{r-1}}}^{(r-1)}; B_{(r-1)j_{i_{r-1}}}]_{1, q_{i_{r-1}}} \tag{1.8}$$

$$\mathbf{B} = [\tau_{i_r}(b_{rj_{i_r}}; \beta_{rj_{i_r}}^{(1)}, \dots, \beta_{rj_{i_r}}^{(r)}; B_{rj_{i_r}})]_{1, q_{i_r}} \tag{1.9}$$

$$B = [(d_j^{(1)}, \delta_j^{(1)}; D_j^{(1)})_{1, m^{(1)}}], [\tau_{i^{(1)}}(d_{j_{i^{(1)}}}^{(1)}, \delta_{j_{i^{(1)}}}^{(1)}; D_{j_{i^{(1)}}}^{(1)})_{m^{(1)}+1, q_i^{(1)}}]; \dots ;$$

$$[(d_j^{(r)}, \delta_j^{(r)}; D_j^{(r)})_{1, m^{(r)}}], [\tau_{i^{(r)}}(d_{j_{i^{(r)}}}^{(r)}, \delta_{j_{i^{(r)}}}^{(r)}; D_{j_{i^{(r)}}}^{(r)})_{m^{(r)}+1, q_i^{(r)}}] \tag{1.10}$$

$$U = 0, n_2; 0, n_3; \dots ; 0, n_{r-1}; V = m^{(1)}, n^{(1)}; m^{(2)}, n^{(2)}; \dots ; m^{(r)}, n^{(r)} \tag{1.11}$$

$$X = p_{i_2}, q_{i_2}, \tau_{i_2}; R_2; \dots ; p_{i_{r-1}}, q_{i_{r-1}}, \tau_{i_{r-1}} : R_{r-1}; Y = p_{i^{(1)}}, q_{i^{(1)}}, \tau_{i^{(1)}}; R^{(1)}; \dots ; p_{i^{(r)}}, q_{i^{(r)}}, \tau_{i^{(r)}}; R^{(r)} \tag{1.12}$$

Srivastava ([6], p. 1, Eq. (1)) have introduced the general class of polynomials :

$$S_N^M(x) = \sum_{K=0}^{[N/M]} \frac{(-N)_{MK}}{K!} A_{N,K} x^K \tag{1.13}$$

where M is an arbitrary positive integer and the coefficients $A_{N,K}$ are arbitrary constants real or complex. On specializing these coefficients $A_{N,K}$, $S_N^M[.]$ yields a number of known polynomials as special cases. These include, among others, Hermite polynomials, Jacobi polynomials, Laguerre polynomials, bessel polynomials and several others([10], p. 158-161).

We shall note

$$a_{NK} = \frac{(-N)_{MK}}{K!} A_{N,K} \tag{1.14}$$

2. Main formula.

Theorem.

$$\text{If } {}_2F_1(a, b; c; z) {}_2F_1(a, b; cdz) = \sum_{l=0}^{\infty} C_l z^l \tag{2.1}$$

then

$$\int_0^1 {}_4F_3 \left[\begin{matrix} a, b, \frac{c+d}{2}, \frac{c+b-1}{2} \\ a+b, c, d \end{matrix} \middle| 4z(1-z) \right] S_N^M(z^h) \mathfrak{J}(a_1 z^{h_1}, \dots, a_r z^{h_r}) dz = \sum_{l=0}^{\infty} \sum_{K=0}^{[N/M]} a_{NK} \frac{(c+d-1)_l C_l}{(a+b)_l}$$

$$\mathfrak{J}_{X;p_i r+1, q_i r+1, \tau_i r; R_r; Y}^{U; 0, n_r+1; V} \left(\begin{matrix} a_1 & \mathbb{A}; (-1-Kh; h_1, \dots, h_r; 1), \mathbf{A} : A \\ \vdots & \vdots \\ a_r & \mathbb{B}; \mathbf{B}, (-1-Kh-1; h_1, \dots, h_r; 1) : B \end{matrix} \right) \tag{2.2}$$

provided

$$h, h_i > 0 (i = 1, \dots, r), \quad 1 + \sum_{i=1}^r h_i \min_{1 \leq j \leq m^{(i)}} \operatorname{Re} \left(D_j^{(i)} \frac{d_j^{(i)}}{\delta_j^{(i)}} \right) > 0$$

$|\arg(a_i z^{h_i})| < \frac{1}{2} A_i^{(k)} \pi$ where $A_i^{(k)}$ is defined by (1.4) and the series on the right-hand side of (2.2) is absolutely convergent.

Proof

To prove the above theorem, we use the following formula due to Slater ([5], p. 79, Eq. (2.5.27))

$${}_4F_3 \left[\begin{matrix} a, b, \frac{c+d}{2}, \frac{c+b-1}{2} \\ a+b, c, d \end{matrix} \middle| 4z(1-z) \right] = \sum_{l=0}^{\infty} \frac{(c+d-1)_l}{(a+b)_l} C_l z^l \tag{2.3}$$

where C_l is given by (2.1). Multiplying the both-sides of (2.3) by $S_N^M(z^h) \mathfrak{J}(a_1 z^{h_1}, \dots, a_r z^{h_r})$ and integrating with respect to z between 0 to 1, we get

$$\int_0^1 {}_4F_3 \left[\begin{matrix} a, b, \frac{c+d}{2}, \frac{c+b-1}{2} \\ a+b, c, d \end{matrix} \middle| 4z(1-z) \right] S_N^M(z^h) \mathfrak{J}(a_1 z^{h_1}, \dots, a_r z^{h_r}) dz = \sum_{l=0}^{\infty} \frac{(c+d-1)_l}{(a+b)_l} C_l z^l \tag{2.4}$$

Substituting the expression of the multivariable Gimel-function in terms of Mellin-Barnes multiple integrals contour with the help of (1.1) and $S_N^M[z^h]$ with the help of (1.13), interchanging the order of integrations and summation (which is permissible under the conditions mentioned in (2.2)), now, evaluating the inner z -integral and interpreting the Mellin-Barnes multiple integrals contour in terms of the multivariable Gimel-function, we get the desired result (2.2).

3. Particular cases.

Taking $b = c = d$ in the theorem, we get the following interesting integral :

Corollary 1.

$$\int_0^1 {}_2F_1 \left[\begin{matrix} a, c-\frac{1}{2} \\ a+c \end{matrix} \middle| 4z(1-z) \right] S_N^M(z^h) \mathfrak{J}(a_1 z^{h_1}, \dots, a_r z^{h_r}) dz = \sum_{l=0}^{\infty} \sum_{K=0}^{[N/M]} a_{NK} \frac{(2c-1)_l (2a)_l}{(a+c)_l l!}$$

$$\mathfrak{J}_{X;p_i r+1, q_i r+1, \tau_i r; R_r; Y}^{U; 0, n_r+1; V} \left(\begin{matrix} a_1 & \mathbb{A}; (-1-Kh; h_1, \dots, h_r; 1), \mathbf{A} : A \\ \vdots & \vdots \\ a_r & \mathbb{B}; \mathbf{B}, (-1-Kh-1; h_1, \dots, h_r; 1) : B \end{matrix} \right) \tag{3.1}$$

under the same existence conditions mentioned in (2.2).

Taking $a = -e$ in the above corollary, it reduces to the interesting integral

Corollary 2.

$$\int_0^1 {}_2F_1 \left[\begin{matrix} -e, c-\frac{1}{2} \\ c-e \end{matrix} \middle| 4z(1-z) \right] S_N^M(z^h) \mathfrak{J}(a_1 z^{h_1}, \dots, a_r z^{h_r}) dz = \sum_{l=0}^{2e} \sum_{K=0}^{[N/M]} a_{NK} \frac{(2c-1)_l (-2e)_l}{(c-e)_l l!}$$

$$\mathfrak{J}_{X;p_i r+1, q_i r+1, \tau_{i_r}: R_r: Y}^{U; 0, n_r+1: V} \left(\begin{matrix} a_1 & \mathbb{A}; (-1-Kh; h_1, \dots, h_r; 1), \mathbf{A} : A \\ \cdot & \cdot \\ \cdot & \cdot \\ a_r & \mathbb{B}; \mathbf{B}, (-1-Kh-1; h_1, \dots, h_r; 1) : B \end{matrix} \right) \quad (3.2)$$

under the same existence conditions mentioned in (2.2).

4. Conclusion.

The main integral (2.2) established here are unified and act as key formulae. Thus the multivariable Gimel-function occurring in these integrals can be suitably specialized to a remarkably wide variety of special functions (or product of several such special functions) which are expressible in terms of E, F, G, H, I, Aleph-functions of one or more variables. Again the class of polynomials involved in the integral (2.2) reduces to a large number of polynomials listed by Srivastava and Singh ([9], p.158-161), therefore, from the integral (2.2) we can further obtain various integrals involving a number of simpler polynomials.

REFERENCES.

- [1] F. Ayant, An integral associated with the Aleph-functions of several variables. *International Journal of Mathematics Trends and Technology (IJMTT)*, 31(3) (2016), 142-154.
- [2] B.L.J. Braaksma, Asymptotics expansions and analytic continuations for a class of Barnes-integrals, *Compositio Math.* 15 (1962-1964), 239-341.
- [3] Y.N. Prasad, Multivariable I-function , *Vijnana Parishad Anusandhan Patrika* 29 (1986) , 231-237.
- [4] J. Prathima, V. Nambisan and S.K. Kurumujji, A Study of I-function of Several Complex Variables, *International Journal of Engineering Mathematics* Vol (2014), 1-12.
- [5] L.J. Slater, *Generalized hypergeometric function*, Cambridge University press, Cambridge, 1966.
- [6] H.M. Srivastava, A contour integral involving Fox's H-function, *Indian. J. Math.* 14(1972), 1-6.
- [7] H.M. Srivastava and R. Panda, Some expansion theorems and generating relations for the H-function of several complex variables. *Comment. Math. Univ. St. Paul.* 24 (1975),119-137.
- [8] H.M. Srivastava and R.Panda, Some expansion theorems and generating relations for the H-function of several complex variables II. *Comment. Math. Univ. St. Paul.* 25 (1976), 167-197.
- [9] H.M. Srivastava and N.P. Singh, The integration of certain products of the multivariable H-function with a general class of polynomials, *Rend. CirC. Mat. Palermo.* 32(2)(1983), 157-187.