

# Integration of certain Gimel-function with respect to their parameters

Frédéric Ayant

Teacher in High School , France

**ABSTRACT**

The object of the present paper is to obtain some interesting results by integrating the multivariable Gimel-function with respect to its parameters. Such integrals are useful in the study of certain boundary value problems.

**KEYWORDS :** Multivariable Gimel-function, multiple integral contours, integration with respect to a parameter.

**2010 Mathematics Subject Classification.** 33C99, 33C60, 44A20

## 1. Introduction and preliminaries.

Throughout this paper, let  $\mathbb{C}, \mathbb{R}$  and  $\mathbb{N}$  be set of complex numbers, real numbers and positive integers respectively. Also  $\mathbb{N}_0 = \mathbb{N} \cup \{0\}$ . We define a generalized transcendental function of several complex variables.

$$\mathfrak{I}(z_1, \dots, z_r) = \mathfrak{I}_{p_{i_2}, q_{i_2}, \tau_{i_2}; R_2; p_{i_3}, q_{i_3}, \tau_{i_3}; R_3; \dots; p_{i_r}, q_{i_r}, \tau_{i_r}; R_r; p_{i(1)}, q_{i(1)}, \tau_{i(1)}; R^{(1)}; \dots; p_{i(r)}, q_{i(r)}, \tau_{i(r)}; R^{(r)}}^{0, n_2; 0, n_3; \dots; 0, n_r; m^{(1)}, n^{(1)}; m^{(2)}, n^{(2)}; \dots; m^{(r)}, n^{(r)}} \left( \begin{matrix} z_1 \\ \cdot \\ \cdot \\ \cdot \\ z_r \end{matrix} \right)$$

$$[(a_{2j}; \alpha_{2j}^{(1)}, \alpha_{2j}^{(2)}; A_{2j})]_{1, n_2}, [\tau_{i_2}(a_{2ji_2}; \alpha_{2ji_2}^{(1)}, \alpha_{2ji_2}^{(2)}; A_{2ji_2})]_{n_2+1, p_{i_2}}; [(a_{3j}; \alpha_{3j}^{(1)}, \alpha_{3j}^{(2)}, \alpha_{3j}^{(3)}; A_{3j})]_{1, n_3},$$

$$[\tau_{i_2}(b_{2ji_2}; \beta_{2ji_2}^{(1)}, \beta_{2ji_2}^{(2)}; B_{2ji_2})]_{1, q_{i_2}};$$

$$[\tau_{i_3}(a_{3ji_3}; \alpha_{3ji_3}^{(1)}, \alpha_{3ji_3}^{(2)}, \alpha_{3ji_3}^{(3)}; A_{3ji_3})]_{n_3+1, p_{i_3}}; \dots; [(a_{rj}; \alpha_{rj}^{(1)}, \dots, \alpha_{rj}^{(r)}; A_{rj})]_{1, n_r},$$

$$[\tau_{i_3}(b_{3ji_3}; \beta_{3ji_3}^{(1)}, \beta_{3ji_3}^{(2)}, \beta_{3ji_3}^{(3)}; B_{3ji_3})]_{1, q_{i_3}}; \dots$$

$$[\tau_{i_r}(a_{rji_r}; \alpha_{rji_r}^{(1)}, \dots, \alpha_{rji_r}^{(r)}; A_{rji_r})]_{n_r+1, p_{i_r}}; [(c_j^{(1)}, \gamma_j^{(1)}; C_j^{(1)})]_{1, n^{(1)}}, [\tau_{i(1)}(c_{ji(1)}^{(1)}, \gamma_{ji(1)}^{(1)}; C_{ji(1)}^{(1)})]_{n^{(1)}+1, p_i^{(1)}}]$$

$$[\tau_{i_r}(b_{rji_r}; \beta_{rji_r}^{(1)}, \dots, \beta_{rji_r}^{(r)}; B_{rji_r})]_{1, q_{i_r}}; [(d_j^{(1)}, \delta_j^{(1)}; D_j^{(1)})]_{1, m^{(1)}}, [\tau_{i(1)}(d_{ji(1)}^{(1)}, \delta_{ji(1)}^{(1)}; D_{ji(1)}^{(1)})]_{m^{(1)}+1, q_i^{(1)}}]$$

$$; \dots; [(c_j^{(r)}, \gamma_j^{(r)}; C_j^{(r)})]_{1, n^{(r)}}, [\tau_{i(r)}(c_{ji(r)}^{(r)}, \gamma_{ji(r)}^{(r)}; C_{ji(r)}^{(r)})]_{n^{(r)}+1, p_i^{(r)}}]$$

$$; \dots; [(d_j^{(r)}, \delta_j^{(r)}; D_j^{(r)})]_{1, m^{(r)}}, [\tau_{i(r)}(d_{ji(r)}^{(r)}, \delta_{ji(r)}^{(r)}; D_{ji(r)}^{(r)})]_{m^{(r)}+1, q_i^{(r)}}]$$

$$= \frac{1}{(2\pi\omega)^r} \int_{L_1} \dots \int_{L_r} \psi(s_1, \dots, s_r) \prod_{k=1}^r \theta_k(s_k) z_k^{s_k} ds_1 \dots ds_r \tag{1.1}$$

with  $\omega = \sqrt{-1}$

$$\psi(s_1, \dots, s_r) = \frac{\prod_{j=1}^{n_2} \Gamma^{A_{2j}} (1 - a_{2j} + \sum_{k=1}^2 \alpha_{2j}^{(k)} s_k)}{\sum_{i_2=1}^{R_2} [\tau_{i_2} \prod_{j=n_2+1}^{p_{i_2}} \Gamma^{A_{2ji_2}} (a_{2ji_2} - \sum_{k=1}^2 \alpha_{2ji_2}^{(k)} s_k) \prod_{j=1}^{q_{i_2}} \Gamma^{B_{2ji_2}} (1 - b_{2ji_2} + \sum_{k=1}^2 \beta_{2ji_2}^{(k)} s_k)]}$$

$$\frac{\prod_{j=1}^{n_3} \Gamma^{A_{3j}} (1 - a_{3j} + \sum_{k=1}^3 \alpha_{3j}^{(k)} s_k)}{\sum_{i_3=1}^{R_3} [\tau_{i_3} \prod_{j=n_3+1}^{p_{i_3}} \Gamma^{A_{3ji_3}} (a_{3ji_3} - \sum_{k=1}^3 \alpha_{3ji_3}^{(k)} s_k) \prod_{j=1}^{q_{i_3}} \Gamma^{B_{3ji_3}} (1 - b_{3ji_3} + \sum_{k=1}^3 \beta_{3ji_3}^{(k)} s_k)]}$$

.

.

.

$$\frac{\prod_{j=1}^{n_r} \Gamma^{A_{rj}} (1 - a_{rj} + \sum_{k=1}^r \alpha_{rj}^{(k)} s_k)}{\sum_{i_r=1}^{R_r} [\tau_{i_r} \prod_{j=n_r+1}^{p_{i_r}} \Gamma^{A_{rji_r}} (a_{rji_r} - \sum_{k=1}^r \alpha_{rji_r}^{(k)} s_k) \prod_{j=1}^{q_{i_r}} \Gamma^{B_{rji_r}} (1 - b_{rji_r} + \sum_{k=1}^r \beta_{rji_r}^{(k)} s_k)]} \tag{1.2}$$

and

$$\theta_k(s_k) = \frac{\prod_{j=1}^{m^{(k)}} \Gamma^{D_j^{(k)}} (d_j^{(k)} - \delta_j^{(k)} s_k) \prod_{j=1}^{n^{(k)}} \Gamma^{C_j^{(k)}} (1 - c_j^{(k)} + \gamma_j^{(k)} s_k)}{\sum_{i^{(k)}=1}^{R^{(k)}} [\tau_{i^{(k)}} \prod_{j=m^{(k)}+1}^{q_{i^{(k)}}} \Gamma^{D_{ji^{(k)}}^{(k)}} (1 - d_{ji^{(k)}}^{(k)} + \delta_{ji^{(k)}}^{(k)} s_k) \prod_{j=n^{(k)}+1}^{p_{i^{(k)}}} \Gamma^{C_{ji^{(k)}}^{(k)}} (c_{ji^{(k)}}^{(k)} - \gamma_{ji^{(k)}}^{(k)} s_k)]} \tag{1.3}$$

- 1)  $[(c_j^{(1)}; \gamma_j^{(1)})]_{1, n_1}$  stands for  $(c_1^{(1)}; \gamma_1^{(1)}), \dots, (c_{n_1}^{(1)}; \gamma_{n_1}^{(1)})$ .
- 2)  $n_2, \dots, n_r, m^{(1)}, n^{(1)}, \dots, m^{(r)}, n^{(r)}, p_{i_2}, q_{i_2}, R_2, \tau_{i_2}, \dots, p_{i_r}, q_{i_r}, R_r, \tau_{i_r}, p_{i^{(r)}}, q_{i^{(r)}}, \tau_{i^{(r)}}, R^{(r)} \in \mathbb{N}$  and verify :  
 $0 \leq m_2, 0 \leq n_2 \leq p_{i_2}, \dots, 0 \leq m_r, 0 \leq n_r \leq p_{i_r}, 0 \leq m^{(1)} \leq q_{i^{(1)}}, \dots, 0 \leq m^{(r)} \leq q_{i^{(r)}}.$
- 3)  $\tau_{i_2} (i_2 = 1, \dots, R_2) \in \mathbb{R}^+; \tau_{i_r} \in \mathbb{R}^+ (i_r = 1, \dots, R_r); \tau_{i^{(k)}} \in \mathbb{R}^+ (i = 1, \dots, R^{(k)}), (k = 1, \dots, r).$
- 4)  $\gamma_j^{(k)}, C_j^{(k)} \in \mathbb{R}^+; (j = 1, \dots, n_k); (k = 1, \dots, r); \delta_j^{(k)}, D_j^{(k)} \in \mathbb{R}^+; (j = 1, \dots, m_k); (k = 1, \dots, r).$   
 $\alpha_{kj}^{(l)}, A_{kj} \in \mathbb{R}^+; (j = 1, \dots, n_k); (k = 2, \dots, r); (l = 1, \dots, k).$   
 $\alpha_{kji_k}^{(l)}, A_{kji_k} \in \mathbb{R}^+; (j = n_k + 1, \dots, p_{i_k}); (k = 2, \dots, r); (l = 1, \dots, k).$   
 $\beta_{kji_k}^{(l)}, B_{kji_k} \in \mathbb{R}^+; (j = m_k + 1, \dots, q_{i_k}); (k = 2, \dots, r); (l = 1, \dots, k).$   
 $\delta_{ji^{(k)}}^{(k)} \in \mathbb{R}^+; (i = 1, \dots, R^{(k)}); (j = m^{(k)} + 1, \dots, q_{i^{(k)}}); (k = 1, \dots, r).$   
 $\gamma_{ji^{(k)}}^{(k)} \in \mathbb{R}^+; (i = 1, \dots, R^{(k)}); (j = n^{(k)} + 1, \dots, p_{i^{(k)}}); (k = 1, \dots, r).$
- 5)  $c_j^{(k)} \in \mathbb{C}; (j = 1, \dots, n_k); (k = 1, \dots, r); d_j^{(k)} \in \mathbb{C}; (j = 1, \dots, m_k); (k = 1, \dots, r).$   
 $a_{kji_k} \in \mathbb{C}; (j = n_k + 1, \dots, p_{i_k}); (k = 2, \dots, r).$   
 $b_{kji_k} \in \mathbb{C}; (j = m_k + 1, \dots, q_{i_k}); (k = 2, \dots, r).$   
 $d_{ji^{(k)}}^{(k)} \in \mathbb{C}; (i = 1, \dots, R^{(k)}); (j = m^{(k)} + 1, \dots, q_{i^{(k)}}); (k = 1, \dots, r).$   
 $\gamma_{ji^{(k)}}^{(k)} \in \mathbb{C}; (i = 1, \dots, R^{(k)}); (j = n^{(k)} + 1, \dots, p_{i^{(k)}}); (k = 1, \dots, r).$

The contour  $L_k$  is in the  $s_k (k = 1, \dots, r)$ - plane and run from  $\sigma - i\infty$  to  $\sigma + i\infty$  where  $\sigma$  if is a real number with loop, if necessary to ensure that the poles of  $\Gamma^{A_{2j}} \left( 1 - a_{2j} + \sum_{k=1}^2 \alpha_{2j}^{(k)} s_k \right) (j = 1, \dots, n_2), \Gamma^{A_{3j}} \left( 1 - a_{3j} + \sum_{k=1}^3 \alpha_{3j}^{(k)} s_k \right) (j = 1, \dots, n_3), \dots, \Gamma^{A_{rj}} \left( 1 - a_{rj} + \sum_{i=1}^r \alpha_{rj}^{(i)} s_k \right) (j = 1, \dots, n_r), \Gamma^{C_j^{(k)}} \left( 1 - c_j^{(k)} + \gamma_j^{(k)} s_k \right) (j = 1, \dots, n^{(k)})(k = 1, \dots, r)$  to the right of the contour  $L_k$  and the poles of  $\Gamma^{D_j^{(k)}} \left( d_j^{(k)} - \delta_j^{(k)} s_k \right) (j = 1, \dots, m^{(k)})(k = 1, \dots, r)$  lie to the left of the contour  $L_k$ . The condition for absolute convergence of multiple Mellin-Barnes type contour (1.1) can be obtained of the corresponding conditions for multivariable H-function given by as :

$$|arg(z_k)| < \frac{1}{2} A_i^{(k)} \pi \text{ where}$$

$$A_i^{(k)} = \sum_{j=1}^{m^{(k)}} D_j^{(k)} \delta_j^{(k)} + \sum_{j=1}^{n^{(k)}} C_j^{(k)} \gamma_j^{(k)} - \tau_{i^{(k)}} \left( \sum_{j=m^{(k)}+1}^{q_i^{(k)}} D_{ji^{(k)}}^{(k)} \delta_{ji^{(k)}}^{(k)} + \sum_{j=n^{(k)}+1}^{p_i^{(k)}} C_{ji^{(k)}}^{(k)} \gamma_{ji^{(k)}}^{(k)} \right) +$$

$$- \tau_{i_2} \left( \sum_{j=n_2+1}^{p_{i_2}} A_{2ji_2} \alpha_{2ji_2}^{(k)} + \sum_{j=1}^{q_{i_2}} B_{2ji_2} \beta_{2ji_2}^{(k)} \right) - \dots - \tau_{i_r} \left( \sum_{j=n_r+1}^{p_{i_r}} A_{rji_r} \alpha_{rji_r}^{(k)} + \sum_{j=1}^{q_{i_r}} B_{rji_r} \beta_{rji_r}^{(k)} \right) \tag{1.4}$$

Following the lines of Braaksma ([2] p. 278), we may establish the the asymptotic expansion in the following convenient form :

$$\aleph(z_1, \dots, z_r) = O(|z_1|^{\alpha_1}, \dots, |z_r|^{\alpha_r}), \max(|z_1|, \dots, |z_r|) \rightarrow 0$$

$$\aleph(z_1, \dots, z_r) = O(|z_1|^{\beta_1}, \dots, |z_r|^{\beta_r}), \min(|z_1|, \dots, |z_r|) \rightarrow \infty \text{ where } i = 1, \dots, r :$$

$$\alpha_i = \min_{1 \leq j \leq m^{(i)}} \operatorname{Re} \left[ D_j^{(i)} \left( \frac{d_j^{(i)}}{\delta_j^{(i)}} \right) \right] \text{ and } \beta_i = \max_{1 \leq j \leq n^{(i)}} \operatorname{Re} \left[ C_j^{(i)} \left( \frac{c_j^{(i)} - 1}{\gamma_j^{(i)}} \right) \right]$$

**Remark 1.**

If  $n_2 = \dots = n_{r-1} = p_{i_2} = q_{i_2} = \dots = p_{i_{r-1}} = q_{i_{r-1}} = 0$  and  $A_{2j} = A_{2ji_2} = B_{2ji_2} = \dots = A_{rj} = A_{rji_r} = B_{rji_r} = 1$   $A_{rj} = A_{rji_r} = B_{rji_r} = 1$ , then the multivariable Gimel-function reduces in the multivariable Aleph- function defined by Ayant [1].

**Remark 2.**

If  $n_2 = \dots = n_r = p_{i_2} = q_{i_2} = \dots = p_{i_r} = q_{i_r} = 0$  and  $\tau_{i_2} = \dots = \tau_{i_r} = \tau_{i^{(1)}} = \dots = \tau_{i^{(r)}} = R_2 = \dots = R_r = R^{(1)} = \dots = R^{(r)} = 1$ , then the multivariable Gimel-function reduces in a multivariable I-function defined by Prathima et al. [4].

**Remark 3.**

If  $A_{2j} = A_{2ji_2} = B_{2ji_2} = \dots = A_{rj} = A_{rji_r} = B_{rji_r} = 1$  and  $\tau_{i_2} = \dots = \tau_{i_r} = \tau_{i^{(1)}} = \dots = \tau_{i^{(r)}} = R_2 = \dots = R_r = R^{(1)} = \dots = R^{(r)} = 1$ , then the generalized multivariable Gimel-function reduces in multivariable I-function defined by Prasad [3].

**Remark 4.**

If the three above conditions are satisfied at the same time, then the generalized multivariable Gimel-function reduces in the multivariable H-function defined by Srivastava and Panda [6,7].

In your investigation, we shall use the following notations.

$$\mathbb{A} = [(a_{2j}; \alpha_{2j}^{(1)}, \alpha_{2j}^{(2)}; A_{2j})]_{1, n_2}, [\tau_{i_2}(a_{2ji_2}; \alpha_{2ji_2}^{(1)}, \alpha_{2ji_2}^{(2)}; A_{2ji_2})]_{n_2+1, p_{i_2}}, [(a_{3j}; \alpha_{3j}^{(1)}, \alpha_{3j}^{(2)}, \alpha_{3j}^{(3)}; A_{3j})]_{1, n_3},$$

$$[\tau_{i_3}(a_{3j_{i_3}}; \alpha_{3j_{i_3}}^{(1)}, \alpha_{3j_{i_3}}^{(2)}, \alpha_{3j_{i_3}}^{(3)}; A_{3j_{i_3}})]_{n_3+1, p_{i_3}}; \dots; [(a_{(r-1)j}; \alpha_{(r-1)j}^{(1)}, \dots, \alpha_{(r-1)j}^{(r-1)}; A_{(r-1)j})_{1, n_{r-1}}],$$

$$[\tau_{i_{r-1}}(a_{(r-1)j_{i_{r-1}}}; \alpha_{(r-1)j_{i_{r-1}}}^{(1)}, \dots, \alpha_{(r-1)j_{i_{r-1}}}^{(r-1)}; A_{(r-1)j_{i_{r-1}}})_{n_{r-1}+1, p_{i_{r-1}}}] \tag{1.5}$$

$$\mathbf{A} = [(a_{rj}; \alpha_{rj}^{(1)}, \dots, \alpha_{rj}^{(r)}; A_{rj})_{1, n_r}], [\tau_{i_r}(a_{rj_{i_r}}; \alpha_{rj_{i_r}}^{(1)}, \dots, \alpha_{rj_{i_r}}^{(r)}; A_{rj_{i_r}})_{n+1, p_{i_r}}] \tag{1.6}$$

$$A = [(c_j^{(1)}, \gamma_j^{(1)}; C_j^{(1)})_{1, n^{(1)}}], [\tau_{i^{(1)}}(c_{j_{i^{(1)}}}^{(1)}, \gamma_{j_{i^{(1)}}}^{(1)}; C_{j_{i^{(1)}}}^{(1)})_{n^{(1)}+1, p_i^{(1)}}]; \dots; [(c_j^{(r)}, \gamma_j^{(r)}; C_j^{(r)})_{1, n^{(r)}}], [\tau_{i^{(r)}}(c_{j_{i^{(r)}}}^{(r)}, \gamma_{j_{i^{(r)}}}^{(r)}; C_{j_{i^{(r)}}}^{(r)})_{n^{(r)}+1, p_i^{(r)}}] \tag{1.7}$$

$$\mathbb{B} = [\tau_{i_2}(b_{2j_{i_2}}; \beta_{2j_{i_2}}^{(1)}, \beta_{2j_{i_2}}^{(2)}; B_{2j_{i_2}})]_{1, q_{i_2}}, [\tau_{i_3}(b_{3j_{i_3}}; \beta_{3j_{i_3}}^{(1)}, \beta_{3j_{i_3}}^{(2)}, \beta_{3j_{i_3}}^{(3)}; B_{3j_{i_3}})]_{1, q_{i_3}}; \dots;$$

$$[\tau_{i_{r-1}}(b_{(r-1)j_{i_{r-1}}}; \beta_{(r-1)j_{i_{r-1}}}^{(1)}, \dots, \beta_{(r-1)j_{i_{r-1}}}^{(r-1)}; B_{(r-1)j_{i_{r-1}}})_{1, q_{i_{r-1}}}] \tag{1.8}$$

$$\mathbf{B} = [\tau_{i_r}(b_{rj_{i_r}}; \beta_{rj_{i_r}}^{(1)}, \dots, \beta_{rj_{i_r}}^{(r)}; B_{rj_{i_r}})_{1, q_{i_r}}] \tag{1.9}$$

$$B = [(d_j^{(1)}, \delta_j^{(1)}; D_j^{(1)})_{1, m^{(1)}}], [\tau_{i^{(1)}}(d_{j_{i^{(1)}}}^{(1)}, \delta_{j_{i^{(1)}}}^{(1)}; D_{j_{i^{(1)}}}^{(1)})_{m^{(1)}+1, q_i^{(1)}}]; \dots;$$

$$[(d_j^{(r)}, \delta_j^{(r)}; D_j^{(r)})_{1, m^{(r)}}], [\tau_{i^{(r)}}(d_{j_{i^{(r)}}}^{(r)}, \delta_{j_{i^{(r)}}}^{(r)}; D_{j_{i^{(r)}}}^{(r)})_{m^{(r)}+1, q_i^{(r)}}] \tag{1.10}$$

$$U = 0, n_2; 0, n_3; \dots; 0, n_{r-1}; V = m^{(1)}, n^{(1)}; m^{(2)}, n^{(2)}; \dots; m^{(r)}, n^{(r)} \tag{1.11}$$

$$X = p_{i_2}, q_{i_2}, \tau_{i_2}; R_2; \dots; p_{i_{r-1}}, q_{i_{r-1}}, \tau_{i_{r-1}} : R_{r-1}; Y = p_{i^{(1)}}, q_{i^{(1)}}, \tau_{i^{(1)}}; R^{(1)}; \dots; p_{i^{(r)}}, q_{i^{(r)}}, \tau_{i^{(r)}}; R^{(r)} \tag{1.12}$$

## 2. Main integrals.

In this section, we evaluate three integrals with respect to their parameters involving the generalized multivariable Gimmel-function.

### Theorem 1.

$$\frac{1}{2\pi\omega} \int_{-\omega}^{\omega} \Gamma(a+x)\Gamma(b-x)\Gamma(c-x)e^{\pm\omega\pi x} \mathfrak{J}_{X; p_{i_r}+1, q_{i_r}, \tau_{i_r}; R_r; Y}^{U; 0, n_r+1; V} \left( \begin{matrix} z_1 & \mathbb{A}; \mathbf{A}: A \\ \cdot & \cdot \\ \cdot & \cdot \\ z_r & \mathbb{B}; \mathbf{B}, (1-d+x; h_1, \dots, h_r; 1) : B \end{matrix} \right) dx =$$

$$e^{\pm\omega\pi a} \Gamma(a+b)\Gamma(a+c) \mathfrak{J}_{X; p_{i_r}+1, q_{i_r}+2, \tau_{i_r}; R_r; Y}^{U; 0, n_r+1; V} \left( \begin{matrix} z_1 & \mathbb{A}; (1+a+b+c-d; h_1, \dots, h_r; 1), \mathbf{A}: A \\ \cdot & \cdot \\ \cdot & \cdot \\ z_r & \mathbb{B}; \mathbf{B}, (1+b-d; h_1, \dots, h_r; 1), (1+c-d; h_1, \dots, h_r; 1) : B \end{matrix} \right) \tag{2.1}$$

provided

$$h_i > 0 (i = 1, \dots, r), \operatorname{Re}(d - a - b - c) + \sum_{i=1}^r h_i \min_{1 \leq j \leq m^{(i)}} \operatorname{Re} \left[ D_j^{(i)} \left( \frac{d_j^{(i)}}{\delta_j^{(i)}} \right) \right] > 0.$$

$$|\operatorname{arg}(z_k)| < \frac{1}{2} A_i^{(k)} \pi \text{ where } A_i^{(k)} \text{ is defined by (1.4).}$$

**Proof**

To prove the theorem 1, we replace the multivariable Gimel-function by this multiple integrals contour with the help of (1.1), change the order of integrations which is justified under the conditions mentioned above. We get

$$\frac{1}{(2\pi\omega)^r} \int_{L_1} \cdots \int_{L_r} \psi(s_1, \dots, s_r) \prod_{k=1}^r \theta_k(s_k) z_k^{s_k} \left[ \frac{1}{2\pi\omega} \int_{-\omega\infty}^{\omega\infty} \frac{\Gamma(a+x)\Gamma(c-x)\Gamma(b-x)}{\Gamma(d-x+\sum_{i=1}^r h_i s_i - x)} e^{\pm\omega\pi x} dx \right] ds_1 \cdots ds_r \quad (2.2)$$

Now, we evaluate the inner integral using by a integral formula ([8], p. 289) and interpreting the resulting expression with the help of (1.1), we obtain the desired result.

**Theorem 2.**

$$\frac{1}{2\pi\omega} \int_{-\omega\infty}^{\omega\infty} \Gamma(b-x)\Gamma(a+x)e^{\pm\omega\pi x} \mathfrak{J}_{X;p_{i_r}+1,q_{i_r}+1,\tau_{i_r};R_r:Y}^{U;0,n_r+1:V} \left( \begin{matrix} z_1 & | & \mathbb{A}; (1-c+x; h_1, \dots, h_r; 1), \mathbf{A} : A \\ \cdot & & \cdot \\ \cdot & & \cdot \\ z_r & | & \mathbb{B}; \mathbf{B}, (1-d+x; k_1, \dots, k_r; 1) : B \end{matrix} \right) dx = \Gamma(a+b)$$

$$e^{\pm\omega\pi a} \mathfrak{J}_{X;p_{i_r}+2,q_{i_r}+2,\tau_{i_r};R_r:Y}^{U;0,n_r+2:V} \left( \begin{matrix} z_1 & | & \mathbb{A}; (1-a-c; h_1, \dots, h_r; 1), (1+a+b+c-d; k_1-h_1, \dots, k_r-h_r; 1), \mathbf{A} : A \\ \cdot & & \cdot \\ \cdot & & \cdot \\ z_r & | & \mathbb{B}; \mathbf{B}, (1+b-d; k_1, \dots, k_r; 1), (1+c-d; k_1-h_1, \dots, k_r-h_r; 1) : B \end{matrix} \right) \quad (2.3)$$

provided

$$h_i, k_i, k_i - h_i > 0 (i = 1, \dots, r), \operatorname{Re}(d - a - b - c) + \sum_{i=1}^r (k_i - h_i) \min_{1 \leq j \leq m^{(i)}} \operatorname{Re} \left[ D_j^{(i)} \left( \frac{d_j^{(i)}}{\delta_j^{(i)}} \right) \right] > 0.$$

$$|\operatorname{arg}(z_k)| < \frac{1}{2} A_i^{(k)} \pi \text{ where } A_i^{(k)} \text{ is defined by (1.4).}$$

**Theorem 3.**

$$\frac{1}{2\pi\omega} \int_{-\omega\infty}^{\omega\infty} \frac{\Gamma(a+x)\Gamma(b-x)}{\Gamma(c-x)} e^{\pm\omega\pi x} \mathfrak{J}_{X;p_{i_r}+1,q_{i_r},\tau_{i_r};R_r:Y}^{U;0,n_r+1:V} \left( \begin{matrix} z_1 & | & \mathbb{A}; (1-d+x; h_1, \dots, h_r; 1), \mathbf{A} : A \\ \cdot & & \cdot \\ \cdot & & \cdot \\ z_r & | & \mathbb{B}; \mathbf{B} : B \end{matrix} \right) dx =$$

$$\frac{\Gamma(a+b)\Gamma(c-a-b-d)}{\Gamma(c-b)} e^{\pm\omega\pi a} \mathfrak{J}_{X;p_{i_r}+2,q_{i_r}+1,\tau_{i_r};R_r:Y}^{U;0,n_r+1:V} \left( \begin{matrix} (-z_1)^{h_1} & | & \mathbb{A}; (1-a-d; h_1, \dots, h_r; 1), \mathbf{A} : A \\ \cdot & & \cdot \\ \cdot & & \cdot \\ (-z_r)^{h_r} & | & \mathbb{B}; \mathbf{B} : B \end{matrix} \right) \quad (2.4)$$

provided

$$h_i > 0 (i = 1, \dots, r), \operatorname{Re}(c - a - b - d) + \sum_{i=1}^r h_i \max_{1 \leq j \leq n^{(i)}} \operatorname{Re} \left[ C_j^{(i)} \left( \frac{c_j^{(i)} - 1}{\gamma_j^{(i)}} \right) \right] > 0.$$

$$|\arg(z_k)| < \frac{1}{2} A_i^{(k)} \pi \text{ where } A_i^{(k)} \text{ is defined by (1.4).}$$

To prove the formulae (3.3) and (3.4), we use the similarly process.

#### 4. Conclusion.

The Gimel-function of several variables presented in this paper, are quite basic in nature. Therefore, on specializing the parameters of this function, we may obtain various known and (news) integrals with respect to parameters concerning the special functions of one variable and several variables, For example, Singh et al. [5] have studied these integrals about the multivariable H-function and they have given several particular cases.

#### REFERENCES.

- [1] F. Ayant, An integral associated with the Aleph-functions of several variables. *International Journal of Mathematics Trends and Technology (IJMTT)*, 31(3) (2016), 142-154.
- [2] B.L.J. Braaksma, Asymptotics expansions and analytic continuations for a class of Barnes-integrals, *Compositio Math.* 15 (1962-1964), 239-341.
- [3] Y.N. Prasad, Multivariable I-function, *Vijnana Parishad Anusandhan Patrika* 29 (1986), 231-237.
- [4] J. Prathima, V. Nambisan and S.K. Kurumujji, A Study of I-function of Several Complex Variables, *International Journal of Engineering Mathematics Vol* (2014), 1-12.
- [5] R; Singh, R.S. Chandel and U.C. Jain, Integration of certain H-functions of several variables with respect to their parameters, *Jnanabha*, 22(1992), 41-47.
- [6] H.M. Srivastava and R. Panda, Some expansion theorems and generating relations for the H-function of several complex variables. *Comment certain of. Math. Univ. St. Paul.* 24 (1975), 119-137.
- [7] H.M. Srivastava and R.Panda, Some expansion theorems and generating relations for the H-function of several complex variables II. *Comment. Math. Univ. St. Paul.* 25 (1976), 167-197.
- [8] E.T. Whittaker and G.N. Watson, *A course of modern analysis*, Cambridge Univ. Press., Cambridge, (1952).